

REFERENCES

Thai

พรพิพย์ นิมนานนิตย์, ภาคภูมิ เต็งอำนวย และ กิตติศักดิ์ ลิขิตวิทยาุณิ. 2543. สมุนไพร
ช่วยให้ผิวขาว. จุฬารัตน์ 19: 15-18.

English

Boylan, J. C., Chowhan, Z. T. and Cooper, J. 1896. Handbook of pharmaceutical excipients (American Pharmaceutical Association). U.S.A.: The American Pharmaceutical Association.

Carter, S. J. 1975. Cooper and Gunn's Dispensing for pharmaceutical students. pp. 678-681. London: Pitman Medical.

Clays, P. and Barel, A. 1998. Efficacy of topical treatment of pigmentation skin disorders with plant hydroquinone glucosides as assessed by quantitative color analysis. J Dermatol 25: 412-414.

Curry, K. V. 1974. Evaluation of skin bleach creams. J. Soc. Cosmet. Chem. 25: 339-354.

Dooley, T. P. 1997. Topical skin depigmentation agents: current products and discovery of novel inhibitors of melanogenesis. Dermatol Treat 8: 275-279.

Farmeco, Co. 1998. New potentials for skin lightening. Newsletter 1:1-2.

Farnsworth, N. R., Bunyaphraphatsara, N. 1992. The medicinal plants Recommended for Primary Health Care System. pp. 67-69 Bangkok: Faculty of Pharmacy, Mahidol University.

Ferrer, A. S., Lopez, J. N., Canovas, F. G. and Carmona, F. G. 1995. Tyrosinase: a comprehensive review of its mechanism. Biochim Biophys Acta 1247: 1-11.

Fitzpatrick, T. B., Eisen, A. L., Wolff, K., Freedberg, I. M. and Austen, K. F. 1993. Dermatology in general medicine: In Skin and hair color and epidermal melanin unit. pp. 62-284. U.S.A.: McGraw-Hill.

Hemsworth, B. N. 1973. Biological changes due to topical application of skin lightening compounds. J. Soc. Cosmet. Chem. 24: 727-733.

- Iida, K., Hase, K., Shimomura, K., Sudo, S., Kadota, S. and Namba, T. 1995. Potent Inhibitors of tyrosinase activity and melanin biosynthesis from *Rheum officinale*. Planta Med. 61: 425-428.
- Imokawa, G., Kawai, M., Mishima, Y. and Motogi, I. 1986. Differential analysis of experimental hypermelanosis induced by UVB, PUVA, and allergic contact dermatitis using a Brownish Guinea Pig model. Arch Dermatol Res 278: 352-362.
- Jang, D. I., et al. 1997. Melanogenesis Inhibitory from paper mulberry. C&T 112: 59-62.
- Kim, J. H. and Lee, K. T. 1998. Inhibitory effects Ramulus Mori extracts on melanogenesis. C&T 113: 65-70.
- Kim, Y. M., Yun, J., Lee, C. K., Min, K. R. and Kim, Y. 2002. Oxyresveratrol and hydroxystibene compounds: Inhibitory effect on tyrosinsae and mechanism of action. J Biol Chem (in press).
- Lee, K. K., and Choi, J. D. 1999. The effect of *Areca Catechu L* extract on anti-inflammatory and antimelanogenesis. Int J Cosmet Sci 21: 275-284.
- Lee, K. T., Kim, B. J. and Kim, J. H. 1997. Biological screening of 100 plant extracts for cosmetic use (I): inhibitory activities of tyrosinase and DOPA auto-oxidation. Int J Cosmet Sci 19: 291-298.
- Lee, O. S. and Kim, E. J. 1995. Skin Lighterning. C&T 110: 51-56.
- Lee, O. S., Kang, H. H. and Han, S. H. 1997. Oriental herbs in cosmetics. C&T 112: 57-64.
- Likhitwitayawuid, K., Sritularak, B. and De-Eknamkul, W. 2000. Tyrosinase inhibitory from *Artocarpus gomezianus*. Planta Med. 66: 275-277.
- Maeda, K. and Fukuda, M. 1991. *In vitro* effectiveness of several whitening cosmetic components in human melanocytes. J. Soc. Cosmet. Chem. 42: 361-368.
- Majmudar, G., Jacob, G., Laboy, Y. and Fisher, L. 1998. An *in vitro* method for screening skin-whitening products. J. cosmet. Sci. 49: 361-367.
- Masamoto, Y., Iida, S. and Kubo, M. 1980. Inhibitory effect of chinese crude drugs on tyrosinase. Planta Med. 40: 361-365.

- Masuda, M., Tejima, T. and Suzuki, T. 1996. Skin Lighteners. C&T 111: 65-67.
- Matsuda, H., Nakamura, S. and Kubo, M. 1994. Studies of cuticle drugs from natural sources. II. Inhibitory effects of prunus plants on melanin biosynthesis. Biol. Pharm. Bull. 17: 1417-1420.
- Mithal, B. M. 1980. Text book of Pharmaceutical formulation. Delhi: Sonu Typoprinters Wazirpur.
- Mongkolsuk S., Robertson, A. and Towers, R. 1957. 2,4,3',5'-tetrahydroxystilbene from *Artocarpus lakoocha*. J.Chem.Soc.: 2231-2233.
- Nichita, N. B., Petrescu, A. J., Dwek, R. A., Wormald, M. R., Platt, F. M. and Petrescu, S. M. 1999. Tyrosinase folding and copper loading in vitro: A crucial role calnexin and α -Glucosidasell. Biochem Biophys Res Commun 261: 720-725.
- Poopyruchpong, N., Rungruangsa, K., Nimmanpisut, S., Panijpan, B. and Ratanabanangkoon, K. 1978. Some physico-chemical properties of 2,4,3',5'-tetrahydroxystilbene. J. Sci. Soc. Thailand 4: 163-167.
- Pope, D. G. 1980. Accelerated testing for prediction of drug product stability. Drug cosm Ind. 48: 116.
- Prota, G. 1996. Melanin and Melanogenesis. C&T 111: 44-45.
- Riley, P. A. 1977. The mechanism of skin pigment production. J. Soc. Cosmet. Chem. 28: 395-401.
- Robinson, N. F., Huxtable, C. R. R. 1988. Clinicopathologic principles for verterinary medicine. New York: Cambridge University.
- Scott, D., Miller, W. H. and Griffin, C. E. 1969. Small animal dermatology. USA: Verterinary Learning System.
- Shimizu, K., Kondo, R. and Sakai, K. 2000. Inhibition of tyrosinase by flavonoids, stilbenes and related 4-substituted resorcinols: structure-activity investigations. Planta Med. 66: 11-19.
- Shimizu, K., Kondo, R., lee, S.H. and Sato, H. 1998. The Inhibitory Componants from *Artocarpus incisus* on Melanin Biosysthesis. Planta Med. 64: 408-412.
- Shin, N. H., et al. 1998. Oxyresveratrol as the potent inhibitor on dopa oxidase activity of mushroom tyrosinase. Biochem Biophys Res Commun 243: 801-803.
- Smith, W. P. 1994. Hydroxy acids and skin aging. C&T 113: 41-48.

- Smith, W. P. 1999. The effects of topical L (+) lactic acid and ascorbic acid on skin whitening. Int J Cosmet Sci 21: 33-40.
- Smolinske, S. C. 1992. Handbook of food, drug and cosmetics excipients. pp. 307-321. USA: CRC press, Inc.
- Sritularak, B. 1998. Chemical constituents of Artocarpus lakoocha and A. gomezianus. Master's Thesis, Department of Pharmacognosy, Graduate School, Chulalongkorn University.
- Sritularak., B., De-Eknamkul, W. and Likhitwitayawuid, K. 1998. Tyrosinase inhibitors from *Artocapus lakoocha*. Thai J. Pharm. Sci. 22: 149-155.
- Tabibian, P. M. 2000. Skin Lightening / Depigmenting Agents [online]. Available from: <http://www.emedicine.com/derm/topic 528.htm> [2001, June, 23]
- Tanunkat, A. 1990. Absorption, metabolism and excretion of 2,4,3',5'-tetrahydroxystilbene in volunteers after oral administration of purified extract of Puag-Haad. Master's thesis, Department of Science (pharmacy), Graduate School, Mahidol University.
- Tiptabiankarn, L. 1967. The antioxidant action of 2,4,3',5'-tetrahydroxystilbene and some of its derivatives. Master's thesis, Department of Pharmaceutical Chemistry, Graduate School, Mahidol University.
- Wallach, J. D. and Boever, W. J. 1993. Disease of exotic animals; Medical and surgical management. Philadelphia: Veterinary Learning System.
- Yodhabandu, C. 1960. A pharmacopoeials study on "Puag-Haad" (2,4,3',5'-tetrahydroxystilbene). Special project for the degree of B. Sc. (Pharm), Chulalongkorn University.
- Yokota, T., Nishio, H. and Kubota, Y. 1998. The inhibitory effect of glabridin from licorice extracts on melagonesis and inflammation. Pigment Cell Res 11: 335-361.
- Yoshimaru, K., Harii, K., Aoyama, T. and Iga, T. 2000. Experience of a strong bleaching treatment for skin hyperpigmentation in orientals [online]. Available from: <http://www.st.rim.or.jp/> [2002, February, 20]
- Zhai, H. and Maibach, H. T. 2001. Skin-Whitening agents. C&T 116: 20-25.

Zuidhoff, H. W. and Rijsbergen, J. M. 2001. Whitening efficacy of frequently used whitening ingredients. C&T 116: 53-59.



ศูนย์วิทยทรัพยากร จุฬาลงกรณ์มหาวิทยาลัย



APPENDICES

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Appendix I

Individual melanin and erythema values in guinea pigs for
Propylene glycol (negative control)

3% Kojic acid (positives control)

3% Haadnun extract

5% Haadnun extract

0.5% Puag-Haad

1.0% Puag-Haad

Skin color was evaluated on guinea pig skin with mexameter MX 16 described in methods.

Table 1 Skin measurement before and after applying propylene glycol (negative control) in guinea pigs ($n = 10$ measurements/ guinea pig). The data show average values (SD in parenthesis).

| Guinea pig | Melanin values | | | | Erythema values | | | |
|------------|-------------------|-------------------|------------------|------------------|------------------|-------------------|------------------|-------------------|
| | Before UVB | Week 0* | Week 2 | Week 4 | Before UVB | Week 0* | Week 2 | Week 4 |
| 1 (SD) | 635.00 (13.50) | 640.00 (18.28) | 655.60 (5.89) | 631.00 (3.89) | 560.00 (9.78) | 550.60 (12.07) | 543.40 (7.65) | 557.70 (3.30) |
| 2 | 642.00 (8.25) | 658.00 (9.90) | 649.50 (7.78) | 635.30 (2.83) | 565.00 (4.22) | 551.50 (4.86) | 548.60 (5.56) | 559.90 (4.01) |
| 3 | 649.00 (9.54) | 663.30 (14.80) | 632.00 (8.47) | 632.90 (7.11) | 566.00 (5.19) | 563.80 (8.42) | 541.00 (7.38) | 552.00 (5.89) |
| 4 | 630.50 (5.12) | 646.20 (6.27) | 645.60 (6.15) | 630.30 (7.50) | 562.10 (8.74) | 554.50 (7.23) | 557.20 (7.96) | 553.60 (14.00) |
| 5 | 634.80 (6.25) | 659.10 (6.19) | 649.20 (8.24) | 634.80 (6.39) | 575.50 (4.98) | 565.60 (5.17) | 568.80 (4.24) | 570.30 (5.96) |
| 6 | 636.00 (8.26) | 657.40 (7.07) | 655.70 (3.92) | 631.20 (9.99) | 557.00 (5.46) | 553.60 (5.06) | 553.60 (5.30) | 559.40 (5.68) |

* The value at week 0 = initial value measured immediately before application of propylene glycol. Application was started on the 18th day following 3-day UVB exposure.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Table 2 Skin measurement before and after applying 3% kojic acid (Positive control) in guinea pigs ($n = 10$ measurements/ guinea pig). The data show average values (SD in parenthesis).

| Guinea pig | Melanin values | | | | Erythema values | | | |
|------------|----------------|---------|--------|--------|-----------------|---------|---------|---------|
| | Before UVB | Week 0* | Week 2 | Week 4 | Before UVB | Week 0* | Week 2 | Week 4 |
| 1 | 623.4 | 678.00 | 630.70 | 641.50 | 570.60 | 558.20 | 550.90 | 575.30 |
| (SD) | (15.21) | (17.80) | (4.62) | (5.95) | (8.99) | (12.99) | (9.02) | (7.56) |
| 2 | 607 | 656.20 | 626.20 | 628.70 | 551.20 | 546.60 | 551.70 | 555.90 |
| | (8.53) | (3.85) | (7.77) | (5.31) | (15.00) | (18.02) | (11.17) | (12.00) |
| 3 | 626 | 666.80 | 623.00 | 625.00 | 574.00 | 551.40 | 540.40 | 557.00 |
| | (4.44) | (10.97) | (9.44) | (3.13) | (4.29) | (6.20) | (8.68) | (4.48) |
| 4 | 614.8 | 654.20 | 639.10 | 635.00 | 567.90 | 558.10 | 547.10 | 554.70 |
| | (5.23) | (4.52) | (9.60) | (4.29) | (7.77) | (9.48) | (8.56) | (15.58) |
| 5 | 650.5 | 662.30 | 635.00 | 619.00 | 566.00 | 554.50 | 563.60 | 554.20 |
| | (6.13) | (5.17) | (6.18) | (6.13) | (9.57) | (10.05) | (3.34) | (7.52) |
| 6 | 640.7 | 668.70 | 638.80 | 622.20 | 568.10 | 556.00 | 551.30 | 540.90 |
| | (4.58) | (6.33) | (4.69) | (3.79) | (5.92) | (5.16) | (7.30) | (7.99) |

* The value at week 0 = initial value measured immediately before application of kojic acid. Application was started on the 18th day following 3-day UVB exposure.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Table 3 Skin measurement before and after applying 3% Haadnun extract in guinea pigs ($n = 10$ measurements/ guinea pig). The data show average values (SD in parenthesis).

| Guinea pig | Melanin values | | | | Erythema values | | | |
|------------|-----------------|-------------------|-------------------|------------------|-------------------|-------------------|-------------------|-------------------|
| | Before UVB | Week 0* | Week 2 | Week 4 | Before UVB | Week 0* | Week 2 | Week 4 |
| 1 (SD) | 645 (13.94) | 656.80 (12.87) | 623.70 (6.18) | 606.20 (6.34) | 556.00 (5.85) | 552.20 (8.75) | 538.90 (6.82) | 532.40 (3.47) |
| 2 | 646.7 (9.31) | 657.30 (9.46) | 632.10 (8.39) | 631.10 (5.07) | 549.20 (3.80) | 552.60 (9.91) | 540.40 (5.50) | 553.70 (4.22) |
| 3 | 614 (11.94) | 658.80 (19.65) | 611.50 (5.91) | 612.70 (3.97) | 559.30 (5.89) | 549.90 (7.45) | 537.90 (4.53) | 556.70 (17.04) |
| 4 | 636.4 (7.12) | 666.00 (5.19) | 663.20 (7.74) | 646.10 (7.17) | 576.80 (5.45) | 566.70 (10.23) | 547.50 (16.43) | 565.30 (8.49) |
| 5 | 628 (6.09) | 663.40 (4.99) | 647.10 (10.67) | 632.70 (5.96) | 564.10 (17.10) | 556.90 (20.84) | 567.80 (5.31) | 556.00 (6.48) |
| 6 | 625.6 (7.78) | 652.00 (6.27) | 656.10 (16.25) | 617.50 (3.78) | 553.40 (3.83) | 548.10 (5.67) | 553.20 (5.94) | 588.00 (37.09) |

* The value at week 0 = initial value measured immediately before application of 3% Haadnun extract. Application was started on the 18th day following 3-day UVB exposure.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Table 4 Skin measurement before and after applying 5% Haadnun extract in guinea pigs ($n = 10$ measurements/ guinea pig). The data show average values (SD in parenthesis).

| Guinea pig | Melanin values | | | | Erythema values | | | |
|------------|-----------------|-------------------|-------------------|------------------|-------------------|-------------------|-------------------|-------------------|
| | Before UVB | Week 0* | Week 2 | Week 4 | Before UVB | Week 0* | Week 2 | Week 4 |
| 1 (SD) | 645 (13.94) | 656.80 (12.87) | 635.60 (11.67) | 621.30 (6.99) | 556.00 (5.85) | 552.20 (8.75) | 553.20 (3.77) | 549.80 (4.08) |
| 2 | 646.7 (9.31) | 657.30 (9.46) | 628.50 (7.62) | 623.00 (3.33) | 549.20 (3.80) | 552.60 (9.91) | 540.70 (5.50) | 545.60 (3.72) |
| 3 | 614 (11.94) | 658.80 (19.65) | 630.30 (11.59) | 616.70 (5.60) | 559.30 (5.89) | 549.90 (7.45) | 539.40 (13.43) | 544.30 (7.50) |
| 4 | 636.4 (7.12) | 666.00 (5.19) | 658.70 (6.38) | 643.00 (5.75) | 576.80 (5.45) | 566.70 (10.23) | 563.50 (17.96) | 572.60 (5.99) |
| 5 | 628 (6.09) | 663.40 (4.99) | 643.10 (8.35) | 624.50 (6.98) | 564.10 (17.10) | 556.90 (20.84) | 579.10 (18.11) | 568.00 (13.54) |
| 6 | 625.6 (7.78) | 652.00 (6.27) | 639.00 (9.58) | 615.90 (6.49) | 553.40 (3.83) | 548.10 (5.67) | 545.10 (2.88) | 554.10 (12.41) |

* The value at week 0 = initial value measured immediately before application of 5% Haadnun extract. Application was started on the 18th day following 3-day UVB exposure.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Table 5 Skin measurement before and after applying 0.5% Puag-Haad in guinea pigs ($n = 10$ measurements/ guinea pig). The data show average values (SD in parenthesis).

| Guinea pig | Melanin values | | | | Erythema values | | | |
|------------|----------------|---------|---------|--------|-----------------|---------|---------|---------|
| | Before UVB | Week 0* | Week 2 | Week 4 | Before UVB | Week 0* | Week 2 | Week 4 |
| 1 | 626.00 | 672.70 | 646.40 | 620.60 | 533.40 | 550.30 | 558.30 | 569.40 |
| (SD) | (8.24) | (8.56) | (7.69) | (8.91) | (5.55) | (6.27) | (18.71) | (8.64) |
| 2 | 635.70 | 659.00 | 619.30 | 622.10 | 557.00 | 552.70 | 546.50 | 573.50 |
| | (7.22) | (12.91) | (11.24) | (5.36) | (7.22) | (4.22) | (9.16) | (11.71) |
| 3 | 659.80 | 673.10 | 625.60 | 625.60 | 565.00 | 552.40 | 555.10 | 570.30 |
| | (10.45) | (11.65) | (4.95) | (4.30) | (5.61) | (4.62) | (13.41) | (7.69) |
| 4 | 636.30 | 670.40 | 625.60 | 601.90 | 567.20 | 552.90 | 555.10 | 540.30 |
| | (6.99) | (11.50) | (4.95) | (7.11) | (6.66) | (4.12) | (13.41) | (3.86) |
| 5 | 634.90 | 669.80 | 634.00 | 618.60 | 556.00 | 551.80 | 542.60 | 553.40 |
| | (6.00) | (5.14) | (10.83) | (6.75) | (8.15) | (8.74) | (6.06) | (6.40) |
| 6 | 635.00 | 663.30 | 651.70 | 617.20 | 561.60 | 546.70 | 554.90 | 556.10 |
| | (5.19) | (5.54) | (5.27) | (3.01) | (6.94) | (3.92) | (5.70) | (4.65) |
| 7 | 634.10 | 664.40 | 644.50 | 611.90 | 571.40 | 562.70 | 548.00 | 556.60 |
| | (5.06) | (5.04) | (8.42) | (3.54) | (6.11) | (3.71) | (11.58) | (14.62) |

* The value at week 0 = initial value measured immediately before application of 0.5% Puag-Haad. Application was started on the 18th day following 3-day UVB exposure.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Table 6 Skin measurement before and after applying 1.0% Puag-Haad in guinea pigs ($n = 10$ measurements/ guinea pig). The data show average values (SD in parenthesis).

| Guinea pig | Melanin values | | | | Erythema values | | | |
|------------|----------------|---------|---------|--------|-----------------|---------|---------|---------|
| | Before UVB | Week 0* | Week 2 | Week 4 | Before UVB | Week 0* | Week 2 | Week 4 |
| 1 | 626.00 | 672.70 | 658.20 | 636.00 | 533.40 | 550.30 | 559.80 | 564.40 |
| (SD) | (8.24) | (8.56) | (10.43) | (6.07) | (5.55) | (6.27) | (5.01) | (6.69) |
| 2 | 635.70 | 659.00 | 634.90 | 621.70 | 557.00 | 552.70 | 551.10 | 584.00 |
| | (7.22) | (12.91) | (7.48) | (8.94) | (7.22) | (4.22) | (13.61) | (5.08) |
| 3 | 659.80 | 673.10 | 648.20 | 643.40 | 565.00 | 552.40 | 537.10 | 586.00 |
| | (10.45) | (11.65) | (19.45) | (4.03) | (5.61) | (4.62) | (8.05) | (7.54) |
| 4 | 636.30 | 670.40 | 626.40 | 615.80 | 567.20 | 552.90 | 568.20 | 547.00 |
| | (6.99) | (11.50) | (5.91) | (3.16) | (6.66) | (4.12) | (6.66) | (3.02) |
| 5 | 634.90 | 669.80 | 654.60 | 622.20 | 556.00 | 551.80 | 555.80 | 546.10 |
| | (6.00) | (5.14) | (5.99) | (8.26) | (8.15) | (8.74) | (12.03) | (10.51) |
| 6 | 635.00 | 663.30 | 654.50 | 623.10 | 561.60 | 546.70 | 568.60 | 557.00 |
| | (5.19) | (5.54) | (8.17) | (5.78) | (6.94) | (3.92) | (14.23) | (6.38) |
| 7 | 634.10 | 664.40 | 655.80 | 615.50 | 571.40 | 562.70 | 578.20 | 561.00 |
| | (5.06) | (5.04) | (4.49) | (5.25) | (6.11) | (3.71) | (13.29) | (6.27) |

* The value at week 0 = initial value measured immediately before application of 1.0% Puag-Haad. Application was started on the 18th day following 3-day UVB exposure.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Appendix IIa

Individual baseline melanin values in female volunteers

- 0.50% Puag-Haad (group A)
- 0.25% Puag-Haad (group B)
- 0.25 % Licorice extract (group C)
- 3.00% Kojic acid (group D)

Skin color was evaluated on outside areas of upper arms with Mexameter MX 16 described in methods.

Table 1 Individual baseline melanin values of subjects in group A

| Volunteer No. | Left arm | | | Right arm | | |
|---------------|----------|---------|---------|-----------|---------|---------|
| | Week -2 | Week -2 | Week -1 | Week -2 | Week -2 | Week -1 |
| A1 | 482.60 | 490.80 | 490.20 | 485.00 | 492.40 | 486.00 |
| A2 | 498.80 | 498.80 | 494.00 | 492.20 | 497.20 | 486.80 |
| A3 | 505.20 | 515.40 | 499.00 | 499.00 | 508.80 | 504.80 |
| A4 | 512.20 | 519.80 | 519.00 | 513.00 | 514.20 | 509.40 |
| A5 | 531.00 | 525.40 | 530.20 | 523.40 | 526.80 | 519.40 |
| A6 | 531.60 | 539.20 | 537.80 | 533.60 | 540.80 | 537.40 |
| A7 | 545.40 | 545.60 | 536.60 | 535.60 | 534.00 | 534.40 |
| A8 | 481.00 | 487.20 | 472.40 | 477.40 | 483.20 | 471.60 |
| A9 | 485.00 | 480.00 | 477.50 | 488.87 | 477.52 | 478.58 |
| A10 | 516.20 | 510.80 | 507.00 | 510.60 | 498.40 | 493.80 |
| A11 | 517.40 | 512.60 | 517.40 | 516.00 | 516.00 | 515.60 |
| A12 | 515.40 | 511.20 | 507.40 | 530.60 | 523.00 | 501.40 |
| A13 | 545.00 | 545.00 | 522.00 | 526.40 | 526.40 | 526.00 |
| A14 | 502.80 | 502.20 | 501.60 | 503.40 | 508.20 | 494.60 |
| A15 | 510.40 | 511.20 | 515.20 | 506.80 | 514.40 | 512.00 |
| A16 | 521.60 | 526.20 | 521.00 | 524.60 | 528.00 | 528.60 |
| A17 | 532.80 | 533.40 | 519.80 | 535.20 | 541.00 | 528.40 |
| A18 | 524.20 | 522.14 | 520.25 | 528.25 | 521.27 | 515.08 |
| A19 | 526.80 | 523.60 | 528.80 | 528.00 | 523.00 | 524.80 |
| A20 | 532.05 | 525.61 | 521.00 | 528.09 | 524.13 | 525.45 |
| Average | 515.87 | 516.31 | 511.91 | 514.30 | 514.94 | 509.71 |
| (SD) | (19.01) | (18.18) | (18.13) | (18.09) | (17.77) | (19.29) |

Table 2 Individual baseline melanin values of subjects in group B

| Volunteer No. | Left arm | | | Right arm | | |
|---------------|----------|---------|---------|-----------|---------|---------|
| | Week -2 | Week -1 | Week 0 | Week -2 | Week -1 | Week 0 |
| B1 | 487.00 | 490.60 | 482.80 | 489.00 | 489.20 | 480.00 |
| B2 | 495.80 | 500.80 | 495.20 | 498.80 | 493.80 | 494.00 |
| B3 | 504.20 | 505.00 | 495.40 | 502.60 | 508.80 | 497.40 |
| B4 | 516.60 | 509.80 | 510.00 | 515.20 | 515.80 | 511.00 |
| B5 | 519.20 | 523.40 | 526.80 | 521.60 | 528.80 | 526.20 |
| B6 | 535.20 | 536.60 | 536.00 | 535.80 | 539.40 | 538.40 |
| B7 | 543.60 | 537.20 | 546.80 | 544.40 | 541.00 | 542.80 |
| B8 | 482.80 | 491.00 | 481.80 | 484.60 | 486.60 | 484.80 |
| B9 | 494.00 | 494.60 | 491.40 | 504.00 | 495.80 | 496.20 |
| B10 | 504.00 | 507.20 | 502.00 | 503.20 | 513.80 | 500.80 |
| B11 | 515.40 | 516.20 | 518.40 | 513.60 | 519.20 | 513.20 |
| B12 | 520.40 | 519.00 | 510.60 | 541.40 | 539.20 | 530.80 |
| B13 | 535.60 | 534.40 | 535.80 | 536.40 | 537.20 | 536.00 |
| B14 | 498.20 | 498.00 | 491.80 | 498.60 | 501.40 | 491.60 |
| B15 | 506.40 | 507.00 | 503.20 | 516.00 | 507.20 | 506.80 |
| B16 | 510.00 | 517.40 | 504.60 | 517.60 | 519.60 | 508.00 |
| B17 | 529.40 | 527.40 | 530.00 | 530.60 | 532.40 | 529.80 |
| B18 | 505.20 | 507.20 | 508.40 | 512.60 | 510.60 | 511.80 |
| B19 | 523.20 | 523.60 | 519.40 | 532.80 | 533.20 | 522.00 |
| B20 | 496.60 | 501.00 | 495.20 | 499.60 | 513.80 | 496.60 |
| Average | 511.14 | 512.37 | 509.28 | 514.92 | 516.34 | 510.91 |
| (SD) | (16.80) | (14.70) | (18.54) | (17.55) | (17.43) | (18.56) |

Table 3 Individual baseline melanin values of subjects in group C

| Volunteer No. | Left arm | | | Right arm | | |
|---------------|----------|---------|---------|-----------|---------|---------|
| | Week -2 | Week -1 | Week 0 | Week -2 | Week -1 | Week 0 |
| C1 | 481.20 | 480.40 | 478.20 | 483.00 | 492.80 | 479.20 |
| C2 | 497.00 | 498.00 | 492.40 | 498.00 | 500.60 | 499.80 |
| C3 | 505.40 | 510.80 | 507.00 | 504.40 | 501.60 | 503.20 |
| C4 | 516.20 | 514.80 | 512.40 | 514.20 | 516.40 | 518.20 |
| C5 | 523.40 | 520.20 | 527.60 | 535.20 | 531.80 | 538.80 |
| C6 | 529.40 | 537.20 | 521.60 | 527.40 | 533.40 | 522.80 |
| C7 | 542.20 | 542.20 | 533.60 | 539.40 | 539.40 | 539.00 |
| C8 | 483.40 | 487.80 | 479.00 | 487.80 | 492.80 | 486.60 |
| C9 | 491.80 | 495.40 | 490.00 | 489.00 | 493.40 | 493.40 |
| C10 | 501.80 | 499.40 | 495.00 | 503.60 | 503.80 | 494.40 |
| C11 | 504.20 | 509.80 | 499.40 | 514.40 | 513.40 | 508.40 |
| C12 | 535.20 | 527.20 | 528.80 | 523.20 | 530.00 | 524.20 |
| C13 | 535.00 | 532.40 | 525.60 | 530.00 | 532.00 | 528.80 |
| C14 | 488.40 | 492.40 | 487.80 | 489.60 | 493.00 | 496.00 |
| C15 | 512.40 | 509.40 | 502.40 | 508.00 | 505.00 | 498.80 |
| C16 | 520.80 | 519.20 | 519.80 | 518.80 | 520.00 | 517.00 |
| C17 | 509.60 | 520.00 | 518.20 | 529.00 | 528.60 | 538.20 |
| C18 | 507.20 | 519.00 | 509.80 | 509.00 | 509.80 | 503.20 |
| C19 | 516.20 | 521.80 | 513.80 | 526.20 | 523.80 | 519.00 |
| C20 | 503.40 | 507.00 | 499.80 | 501.20 | 508.60 | 496.40 |
| Average | 510.21 | 512.22 | 507.11 | 511.57 | 513.51 | 510.27 |
| (SD) | (17.36) | (16.49) | (16.70) | (16.93) | (15.55) | (17.92) |

Table 4 Individual baseline melanin values of subjects in group D

| Volunteer No. | Left arm | | | Right arm | | |
|---------------|----------|---------|---------|-----------|---------|---------|
| | Week -2 | Week -1 | Week 0 | Week -2 | Week -1 | Week 0 |
| D1 | 491.60 | 488.40 | 480.60 | 485.20 | 487.40 | 480.80 |
| D2 | 499.20 | 499.00 | 494.00 | 500.00 | 494.60 | 492.80 |
| D3 | 510.80 | 508.60 | 509.80 | 508.00 | 513.00 | 507.80 |
| D4 | 515.80 | 515.20 | 510.00 | 510.20 | 519.80 | 515.60 |
| D5 | 512.60 | 522.60 | 512.00 | 523.60 | 520.80 | 519.20 |
| D6 | 536.20 | 537.80 | 530.20 | 535.40 | 531.60 | 534.20 |
| D7 | 501.00 | 502.00 | 498.00 | 495.80 | 493.80 | 488.80 |
| D8 | 502.80 | 509.60 | 501.20 | 505.20 | 511.60 | 503.60 |
| D9 | 515.40 | 521.60 | 512.20 | 518.60 | 523.20 | 525.60 |
| D10 | 524.40 | 520.00 | 522.40 | 524.80 | 524.40 | 526.20 |
| D11 | 530.40 | 538.80 | 531.80 | 524.80 | 536.40 | 529.20 |
| D12 | 496.60 | 501.00 | 500.40 | 502.20 | 507.40 | 500.20 |
| D13 | 552.80 | 541.80 | 542.80 | 539.20 | 535.40 | 529.20 |
| D14 | 526.60 | 525.60 | 523.60 | 521.20 | 526.40 | 522.20 |
| D15 | 524.69 | 525.10 | 527.21 | 541.12 | 535.89 | 533.11 |
| D16 | 504.80 | 504.80 | 501.40 | 508.80 | 512.80 | 507.40 |
| D17 | 506.20 | 528.60 | 510.40 | 512.40 | 530.00 | 510.40 |
| D18 | 509.80 | 509.20 | 497.60 | 505.20 | 504.60 | 503.60 |
| D19 | 529.60 | 525.20 | 527.20 | 534.60 | 529.00 | 535.40 |
| D20 | 506.00 | 502.40 | 501.40 | 502.80 | 509.20 | 498.40 |
| Average | 514.86 | 516.37 | 511.71 | 514.96 | 517.36 | 513.19 |
| (SD) | (15.29) | (14.68) | (15.52) | (15.37) | (14.66) | (16.21) |

Appendix IIb

Individual melanin (M) and erythema (E) values in female volunteers for
0.50% Puag-Haad (group A)
0.25% Puag-Haad (group B)
0.25 % Licorice extract (group C)
3.00% Kojic acid (group D)

Skin color was evaluated on outside areas of upper arms with Mexameter MX 16 described in methods.

Table 5 Skin measurement before and after applying propylene glycol (control) in volunteers group A. The data show average values ($n = 5$ measurements).

| Volunteer No. | Control group (left arm) | | | | | | | | | | | | | |
|------------------|--------------------------|--------|---------|--------|--------|--------|--------|--------|---------|--------|---------|--------|--------|--------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | | |
| | M | E | M | E | M | E | M | E | M | E | M | E | | |
| A 1 | 490.20 | 588.80 | 485.00 | 608.60 | 486.20 | 575.40 | 487.60 | 591.80 | 492.60 | 591.80 | 492.00 | 583.60 | 487.20 | 575.00 |
| (SD) | (1.64) | (5.26) | (1.87) | (5.90) | (1.79) | (6.23) | (1.52) | (5.85) | (2.19) | (7.63) | (4.06) | (9.18) | (1.92) | (6.20) |
| A 2 | 494.00 | 586.20 | 495.80 | 566.00 | 495.60 | 570.60 | 495.20 | 577.80 | 495.00 | 574.60 | 501.60 | 575.40 | 501.40 | 576.80 |
| (1.00) | (4.32) | (1.79) | (4.30) | (2.88) | (3.85) | (1.64) | (4.55) | (2.00) | (6.39) | (1.67) | (5.22) | (1.14) | (4.97) | |
| A 3 | 499.00 | 589.80 | 497.60 | 571.60 | 500.40 | 571.40 | 500.80 | 586.80 | 500.20 | 581.60 | 501.00 | 570.60 | 500.60 | 580.20 |
| (2.45) | (2.86) | (1.82) | (3.78) | (2.51) | (4.39) | (2.39) | (5.67) | (1.64) | (6.58) | (0.71) | (1.14) | (1.52) | (7.89) | |
| A 4 | 519.00 | 607.20 | 512.00 | 590.80 | 502.60 | 580.60 | 504.80 | 594.80 | 508.40 | 594.00 | 510.60 | 591.00 | 510.40 | 598.20 |
| (1.73) | (6.69) | (1.58) | (4.09) | (2.41) | (5.13) | (3.27) | (6.69) | (1.67) | (6.56) | (2.30) | (6.44) | (1.34) | (3.03) | |
| A 5 | 530.20 | 609.80 | 528.80 | 598.40 | 527.80 | 603.80 | 521.60 | 591.00 | 523.00 | 585.40 | 522.20 | 583.40 | 520.60 | 596.60 |
| (3.11) | (6.83) | (3.27) | (8.29) | (4.60) | (5.63) | (5.77) | (6.82) | (3.00) | (3.65) | (2.05) | (7.89) | (2.07) | (6.58) | |
| A 6 | 537.80 | 611.60 | 535.20 | 609.20 | 532.80 | 598.40 | 531.40 | 603.60 | 533.00 | 587.80 | 534.40 | 599.00 | 532.40 | 598.00 |
| (1.10) | (3.65) | (1.10) | (10.89) | (1.48) | (2.61) | (0.89) | (7.27) | (2.92) | (11.19) | (1.95) | (3.24) | (2.19) | (4.30) | |
| A 7 | 536.60 | 610.20 | 546.00 | 610.80 | 543.00 | 611.60 | 542.80 | 613.40 | 542.60 | 606.60 | 544.00 | 608.60 | 545.20 | 610.00 |
| (1.14) | (8.87) | (4.00) | (5.50) | (5.05) | (8.32) | (3.56) | (4.10) | (1.67) | (4.51) | (1.22) | (2.61) | (3.56) | (8.15) | |
| A 8 | 472.40 | 556.20 | 477.80 | 558.80 | 479.60 | 551.20 | 481.80 | 558.20 | 478.40 | 558.60 | 480.60 | 546.00 | 481.00 | 548.80 |
| (3.13) | (7.22) | (1.30) | (3.63) | (1.52) | (2.95) | (3.11) | (4.32) | (1.67) | (5.86) | (3.36) | (2.00) | (3.32) | (8.26) | |
| A 10 | 507.00 | 614.60 | 510.40 | 612.60 | 495.40 | 594.80 | 493.40 | 604.40 | 491.80 | 597.40 | 495.60 | 587.60 | 494.40 | 594.60 |
| (2.35) | (14.52) | (2.70) | (7.23) | (8.32) | (3.70) | (5.94) | (6.80) | (4.21) | (4.72) | (4.39) | (10.16) | (1.95) | (3.58) | |

Table 5 Skin measurement before and after applying propylene glycol (control) in volunteers group A. The data show average values (n = 5 measurements).

| Volunteer No. | Control group (right arm) | | | | | | Week 12 | | | | | | |
|------------------|---------------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|------------------|-----------------|-----------------|-----------------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | |
| | M | E | M | E | M | E | M | E | M | E | M | E | |
| A11 | 515.6 (1.82) | 617.8 (3.03) | 511.2 (3.49) | 610.2 (4.09) | 507.2 (2.05) | 600.2 (7.66) | 502.8 (2.28) | 617.2 (3.03) | 507.2 (1.64) | 608.4 (10.50) | 506.4 (1.52) | 605.0 (2.92) | 508.0 (0.71) |
| A12 | 501.4 (1.82) | 607.2 (5.07) | 503.4 (2.41) | 595.0 (5.83) | 506.4 (2.30) | 591.6 (8.79) | 504.0 (3.81) | 609.0 (8.94) | 503.0 (1.00) | 594.0 (5.10) | 506.2 (2.59) | 588.4 (4.34) | 507.0 (1.58) |
| A13 | 526.0 (1.00) | 621.6 (6.80) | 521.2 (2.17) | 606.2 (8.20) | 523.8 (1.79) | 615.8 (5.67) | 521.6 (2.88) | 622.4 (5.08) | 524.8 (1.79) | 609.6 (10.53) | 523.6 (1.67) | 605.8 (5.93) | 516.6 (1.52) |
| A14 | 494.6 (1.14) | 606.6 (2.61) | 495.8 (0.84) | 590.6 (7.96) | 494.2 (3.11) | 594.0 (8.72) | 497.2 (1.79) | 602.8 (8.98) | 496.8 (1.92) | 591.4 (4.45) | 499.2 (0.84) | 498.0 (5.70) | 597.6 (8.08) |
| A15 | 512.0 (2.35) | 620.0 (2.68) | 503.6 (1.67) | 596.4 (3.36) | 510.4 (3.21) | 609.8 (6.06) | 502.4 (3.21) | 620.0 (2.95) | 504.6 (2.07) | 596.4 (6.47) | 504.2 (2.77) | 591.6 (8.71) | 504.2 (1.87) |
| A16 | 528.6 (1.52) | 604.6 (4.10) | 518.2 (1.92) | 589.2 (5.89) | 520.2 (1.48) | 591.6 (1.34) | 514.4 (2.19) | 593.2 (3.03) | 518.2 (0.45) | 591.4 (3.85) | 515.2 (6.87) | 504.2 (6.30) | 597.8 (1.64) |
| A17 | 528.4 (1.67) | 617.2 (7.26) | 536.2 (2.59) | 603.0 (4.12) | 533.8 (1.79) | 607.8 (5.93) | 531.8 (3.63) | 606.0 (3.94) | 530.2 (1.79) | 600.4 (2.61) | 531.4 (1.34) | 603.0 (6.24) | 531.4 (0.89) |
| A19 | 524.8 (6.50) | 615.4 (3.29) | 516.4 (2.07) | 599.8 (1.30) | 522.0 (2.35) | 592.6 (5.46) | 522.0 (3.08) | 608.0 (4.58) | 521.2 (1.30) | 587.0 (2.24) | 521.2 (1.30) | 587.0 (2.24) | 512.6 (2.61) |

Table 6 Skin measurement before and after applying 0.5 % w/v Puag-Haad in volunteers group A. The data show average values ($n = 5$ measurements).

| Volunteer | 0.5 % w/v Puag-Haad group (right arm) | | | | | | | | | | | | | |
|-----------|---------------------------------------|--------|---------|--------|--------|--------|---------|---------|--------|---------|---------|--------|---------|--------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | Week 12 | |
| No. | M | E | M | E | M | E | M | E | M | E | M | E | M | E |
| A 1 | 486.0 | 588.8 | 481.4 | 607.6 | 485.4 | 582.2 | 482.6 | 584.6 | 483.2 | 585.2 | 490.0 | 570.8 | 484.4 | 576.6 |
| (SD) | (1.00) | (4.09) | (2.19) | (4.16) | (1.82) | (6.14) | (1.95) | (12.68) | (1.30) | (13.14) | (2.00) | (7.01) | (0.89) | (9.10) |
| A 2 | 486.8 | 574.0 | 495.4 | 572.8 | 489.0 | 575.2 | 488.0 | 567.2 | 494.2 | 568.4 | 491.0 | 569.6 | 490.2 | 562.0 |
| (1.92) | (4.80) | (2.70) | (3.96) | (3.74) | (5.36) | (0.71) | (5.12) | (2.28) | (5.86) | (1.73) | (8.17) | (0.84) | (6.75) | |
| A 3 | 504.8 | 588.0 | 499.2 | 582.8 | 501.8 | 576.2 | 494.2 | 582.8 | 496.2 | 577.0 | 500.2 | 571.0 | 498.2 | 576.4 |
| (2.95) | (4.12) | (2.28) | (4.76) | (2.77) | (3.83) | (1.30) | (7.69) | (1.30) | (6.20) | (1.30) | (6.20) | (4.64) | (0.45) | (5.46) |
| A 4 | 509.4 | 595.6 | 507.2 | 597.2 | 502.4 | 585.8 | 500.4 | 581.0 | 500.4 | 598.0 | 498.0 | 570.6 | 497.4 | 577.2 |
| (3.51) | (4.93) | (1.30) | (10.33) | (1.14) | (4.32) | (0.89) | (10.32) | (0.89) | (9.41) | (2.45) | (7.50) | (1.82) | (5.26) | |
| A 5 | 519.4 | 590.6 | 520.8 | 587.0 | 515.2 | 591.4 | 519.0 | 588.8 | 524.6 | 587.2 | 514.0 | 568.0 | 515.6 | 584.6 |
| (2.79) | (6.73) | (2.39) | (3.67) | (2.39) | (7.20) | (1.87) | (5.81) | (2.30) | (7.36) | (1.41) | (3.74) | (2.30) | (3.65) | |
| A 6 | 537.4 | 612.4 | 536.0 | 610.2 | 527.6 | 597.4 | 523.6 | 602.4 | 523.2 | 581.0 | 524.0 | 583.0 | 522.2 | 590.2 |
| (1.34) | (9.81) | (2.35) | (3.96) | (1.14) | (3.44) | (1.52) | (4.28) | (3.27) | (6.63) | (1.22) | (10.49) | (3.03) | (6.94) | |
| A 7 | 534.4 | 604.8 | 531.2 | 597.0 | 530.2 | 593.6 | 526.2 | 595.0 | 533.0 | 596.2 | 520.6 | 585.4 | 520.2 | 582.2 |
| (4.04) | (7.05) | (1.92) | (4.30) | (2.59) | (4.93) | (3.63) | (6.24) | (1.87) | (8.07) | (3.21) | (8.26) | (1.48) | (10.76) | |
| A 8 | 471.6 | 557.0 | 477.2 | 564.2 | 475.2 | 554.4 | 469.0 | 551.2 | 475.8 | 545.4 | 475.2 | 547.2 | 474.0 | 547.4 |
| (0.55) | (3.81) | (1.10) | (3.49) | (2.86) | (2.30) | (2.00) | (3.11) | (1.48) | (9.99) | (1.48) | (7.53) | (4.58) | (4.04) | |
| A 10 | 493.8 | 591.2 | 497.8 | 602.2 | 488.4 | 580.4 | 481.4 | 595.2 | 488.6 | 580.8 | 490.2 | 582.6 | 488.0 | 581.8 |
| (1.30) | (5.07) | (2.95) | (5.81) | (1.34) | (7.73) | (2.70) | (6.26) | (0.55) | (6.76) | (4.60) | (12.62) | (3.00) | (1.64) | |

Table 6 Skin measurement before and after applying 0.5 % w/v Puag-Haad in volunteers group A. The data show average values ($n = 5$ measurements).

Table 7 Skin measurement before and after applying propylene glycol (control) in volunteers group B. The data show average values ($n = 5$ measurements).

Table 7 Skin measurement before and after applying propylene glycol (control) in volunteers group 8. The data show average values ($n = 5$ measurements).

| Volunteer No. | Week 0 | | | | Week 2 | | | | Week 4 | | | | Week 6 | | | | Week 8 | | | | Week 10 | | | | |
|------------------|---------|--------|---------|--------|--------|--------|---------|--------|---------|---------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|---------|--------|--------|--------|--------|
| | M | E | M | E | M | E | M | E | M | E | M | E | M | E | M | E | M | E | M | E | M | E | M | E | |
| B11 | 513.2 | 601.8 | 505.6 | 599.6 | 509.6 | 591.4 | 509.2 | 599.8 | 508.6 | 586.6 | 510.0 | 598.4 | 510.6 | 598.4 | 510.6 | 598.4 | 510.6 | 598.4 | 510.6 | 598.4 | 510.6 | 598.4 | 510.6 | 598.4 | |
| (SD) | (0.84) | (2.28) | (2.19) | (3.91) | (2.79) | (7.23) | (3.63) | (3.90) | (2.07) | (5.41) | (1.41) | (2.51) | (1.14) | (1.14) | (1.14) | (1.14) | (1.14) | (1.14) | (1.14) | (1.14) | (1.14) | (1.14) | (1.14) | (1.14) | |
| B12 | 530.8 | 617.4 | 539.4 | 612.0 | 537.2 | 614.8 | 538.0 | 606.4 | 535.6 | 606.2 | 533.2 | 606.8 | 531.4 | 602.6 | 531.4 | 602.6 | 531.4 | 602.6 | 531.4 | 602.6 | 531.4 | 602.6 | 531.4 | 602.6 | 531.4 |
| (5.54) | (5.86) | (1.52) | (6.20) | (4.32) | (7.01) | (1.87) | (8.29) | (1.34) | (3.70) | (4.49) | (6.18) | (2.30) | (6.11) | (2.30) | (6.11) | (2.30) | (6.11) | (2.30) | (6.11) | (2.30) | (6.11) | (2.30) | (6.11) | (2.30) | |
| B13 | 536.0 | 609.4 | 536.8 | 613.6 | 532.2 | 606.8 | 529.2 | 614.0 | 522.4 | 596.8 | 530.0 | 601.6 | 531.0 | 598.8 | 531.0 | 598.8 | 531.0 | 598.8 | 531.0 | 598.8 | 531.0 | 598.8 | 531.0 | 598.8 | 531.0 |
| (2.55) | (9.37) | (2.05) | (8.08) | (4.38) | (8.84) | (6.02) | (8.00) | (4.16) | (4.15) | (4.15) | (5.20) | (6.58) | (0.71) | (2.17) | (6.58) | (0.71) | (2.17) | (6.58) | (0.71) | (2.17) | (6.58) | (0.71) | (2.17) | (6.58) | (0.71) |
| B14 | 491.6 | 593.6 | 494.0 | 594.4 | 492.0 | 580.6 | 488.0 | 581.4 | 484.0 | 560.0 | 483.2 | 573.2 | 484.0 | 584.0 | 484.0 | 584.0 | 484.0 | 584.0 | 484.0 | 584.0 | 484.0 | 584.0 | 484.0 | 584.0 | 484.0 |
| (2.30) | (5.03) | (2.00) | (1.67) | (1.87) | (5.77) | (1.00) | (5.08) | (1.22) | (1.07) | (1.07) | (1.07) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) |
| B15 | 506.8 | 594.4 | 508.8 | 585.2 | 512.0 | 596.8 | 497.8 | 585.6 | 501.6 | 580.4 | 503.8 | 584.0 | 501.2 | 583.6 | 501.2 | 583.6 | 501.2 | 583.6 | 501.2 | 583.6 | 501.2 | 583.6 | 501.2 | 583.6 | 501.2 |
| (1.64) | (5.22) | (1.10) | (3.96) | (3.24) | (5.97) | (1.48) | (5.81) | (2.41) | (9.74) | (9.74) | (9.74) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) | (1.79) |
| B16 | 508.0 | 612.6 | 504.6 | 626.0 | 511.2 | 625.4 | 504.2 | 621.0 | 501.8 | 617.2 | 509.0 | 641.6 | 507.2 | 624.8 | 507.2 | 624.8 | 507.2 | 624.8 | 507.2 | 624.8 | 507.2 | 624.8 | 507.2 | 624.8 | 507.2 |
| (1.00) | (4.98) | (2.30) | (5.79) | (0.45) | (2.70) | (2.59) | (10.68) | (2.17) | (9.93) | (9.93) | (9.93) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) | (3.24) |
| B17 | 529.8 | 612.4 | 525.0 | 613.4 | 526.2 | 610.2 | 525.4 | 615.4 | 525.8 | 619.8 | 523.6 | 606.0 | 521.6 | 602.0 | 521.6 | 602.0 | 521.6 | 602.0 | 521.6 | 602.0 | 521.6 | 602.0 | 521.6 | 602.0 | 521.6 |
| (1.30) | (3.71) | (3.39) | (5.13) | (1.64) | (7.60) | (3.51) | (6.88) | (1.10) | (4.76) | (4.76) | (4.76) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) |
| B18 | 511.8 | 611.0 | 506.2 | 598.0 | 511.0 | 596.6 | 509.8 | 611.4 | 509.2 | 604.8 | 511.8 | 608.2 | 509.6 | 600.6 | 509.6 | 600.6 | 509.6 | 600.6 | 509.6 | 600.6 | 509.6 | 600.6 | 509.6 | 600.6 | 509.6 |
| (3.35) | (10.05) | (3.90) | (12.71) | (1.58) | (3.91) | (0.45) | (6.88) | (1.30) | (6.57) | (6.57) | (6.57) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) | (1.92) |
| B19 | 522.0 | 632.4 | 528.0 | 628.6 | 521.4 | 623.0 | 522.0 | 615.0 | 525.0 | 611.8 | 520.8 | 623.6 | 520.8 | 614.8 | 520.8 | 614.8 | 520.8 | 614.8 | 520.8 | 614.8 | 520.8 | 614.8 | 520.8 | 614.8 | 520.8 |
| (2.30) | (4.28) | (5.20) | (10.57) | (4.56) | (6.00) | (2.55) | (9.38) | (3.67) | (12.26) | (12.26) | (12.26) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) | (5.73) |
| B20 | 496.6 | 580.4 | 502.4 | 569.4 | 501.6 | 576.8 | 497.2 | 575.4 | 494.6 | 568.2 | 502.6 | 574.2 | 502.4 | 576.0 | 502.4 | 576.0 | 502.4 | 576.0 | 502.4 | 576.0 | 502.4 | 576.0 | 502.4 | 576.0 | 502.4 |
| (1.67) | (3.05) | (2.70) | (4.10) | (2.70) | (2.39) | (2.95) | (2.97) | (3.85) | (2.97) | (2.97) | (2.97) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) | (1.95) |

Table 8 Skin measurement before and after applying 0.25 % w/v Puag-Haad in volunteers group B. The data show average values (n = 5 measurements).

| Volunteer No. | 0.25 % w/v Puag-Haad group (right arm) | | | | | | | | | | | | | | | | | | | | |
|------------------|--|--------|---------|--------|---------|--------|---------|--------|---------|--------|---------|--------|--------|---------|---------|---------|---|---|---------|---|---|
| | Week 0 | | | Week 2 | | | Week 4 | | | Week 6 | | | Week 8 | | | Week 10 | | | Week 12 | | |
| | M | E | M | M | E | M | M | E | M | M | E | M | E | M | E | M | E | M | E | M | E |
| B 1 | 480.0 | 566.2 | 481.4 | 577.0 | 481.4 | 562.0 | 475.8 | 571.4 | 473.8 | 570.6 | 476.0 | 565.6 | 474.2 | 556.0 | | | | | | | |
| (SD) | (2.00) | (4.92) | (2.07) | (5.24) | (1.52) | (6.63) | (1.30) | (5.18) | (0.84) | (6.58) | (0.71) | (9.02) | (0.45) | (4.42) | | | | | | | |
| B 2 | 494.0 | 572.8 | 493.0 | 576.4 | 483.2 | 567.6 | 484.6 | 589.6 | 484.6 | 560.4 | 484.6 | 576.8 | 483.0 | 562.4 | | | | | | | |
| (1.00) | (9.18) | (3.08) | (7.16) | (2.77) | (6.07) | (1.14) | (9.84) | (0.55) | (9.24) | (0.55) | (9.55) | (6.57) | (2.55) | (7.60) | | | | | | | |
| B 3 | 497.4 | 586.2 | 497.6 | 589.2 | 494.4 | 575.8 | 493.6 | 574.4 | 487.6 | 564.6 | 491.8 | 566.4 | 488.2 | 573.6 | | | | | | | |
| (1.14) | (5.31) | (1.34) | (1.92) | (1.82) | (4.76) | (1.52) | (7.50) | (1.67) | (4.51) | (1.67) | (4.51) | (1.92) | (1.15) | (2.17) | (3.51) | | | | | | |
| B 4 | 511.0 | 583.2 | 512.4 | 584.2 | 508.4 | 595.4 | 508.0 | 573.8 | 508.8 | 575.6 | 504.2 | 580.8 | 505.4 | 577.4 | | | | | | | |
| (0.71) | (1.48) | (2.07) | (2.19) | (2.19) | (10.14) | (0.00) | (5.54) | (0.84) | (6.54) | (0.84) | (6.54) | (1.92) | (1.14) | (8.17) | | | | | | | |
| B 5 | 526.2 | 630.2 | 522.2 | 609.4 | 518.2 | 610.2 | 507.8 | 605.2 | 506.8 | 580.4 | 506.0 | 592.0 | 505.4 | 581.4 | | | | | | | |
| (3.27) | (5.63) | (1.64) | (4.83) | (1.79) | (9.50) | (4.44) | (11.12) | (1.10) | (8.62) | (1.10) | (8.62) | (9.19) | (2.97) | (7.16) | | | | | | | |
| B 6 | 538.4 | 626.2 | 535.4 | 629.8 | 522.4 | 603.6 | 522.0 | 610.8 | 518.0 | 611.4 | 509.2 | 604.2 | 507.6 | 601.2 | | | | | | | |
| (2.41) | (4.66) | (2.07) | (1.64) | (6.58) | (9.24) | (2.35) | (4.49) | (1.41) | (3.85) | (1.41) | (3.85) | (3.77) | (1.14) | (11.73) | | | | | | | |
| B 7 | 542.8 | 625.8 | 537.0 | 598.8 | 538.0 | 598.6 | 531.8 | 610.0 | 530.0 | 587.2 | 525.6 | 590.6 | 524.0 | 596.8 | | | | | | | |
| (2.59) | (5.67) | (3.16) | (3.42) | (2.45) | (2.07) | (2.39) | (4.58) | (1.41) | (4.49) | (1.41) | (4.49) | (0.55) | (2.07) | (4.44) | | | | | | | |
| B 8 | 484.8 | 589.4 | 485.8 | 580.8 | 476.6 | 567.4 | 475.6 | 566.8 | 472.0 | 558.8 | 475.6 | 557.4 | 471.8 | 567.4 | | | | | | | |
| (4.49) | (11.44) | (5.67) | (15.96) | (2.41) | (8.96) | (2.07) | (9.23) | (1.00) | (5.81) | (1.00) | (5.81) | (2.07) | (9.26) | (1.79) | (11.93) | | | | | | |
| B 9 | 496.2 | 610.6 | 497.0 | 593.2 | 494.0 | 588.2 | 492.2 | 594.4 | 492.4 | 582.0 | 497.4 | 590.8 | 492.6 | 587.0 | | | | | | | |
| (0.84) | (6.43) | (1.58) | (5.97) | (1.87) | (6.26) | (1.92) | (4.39) | (0.89) | (10.51) | (0.89) | (10.51) | (3.29) | (8.87) | (1.67) | (7.68) | | | | | | |
| B 10 | 500.8 | 592.0 | 503.2 | 581.4 | 502.8 | 593.0 | 501.4 | 585.6 | 503.6 | 578.8 | 498.8 | 577.4 | 499.0 | 581.2 | | | | | | | |
| (1.10) | (4.06) | (1.30) | (4.98) | (3.11) | (4.18) | (1.95) | (5.03) | (2.07) | (10.18) | (2.59) | (10.18) | (2.59) | (2.92) | (6.26) | | | | | | | |

Table 8 Skin measurement before and after applying 0.25 % w/v Puag-Haad in volunteers group B. The data show average values ($n = 5$ measurements).

| No. | Volunteer | | 0.25 % w/v Puag-Haad group (left arm) | | | | | | | | | | | |
|-----|-----------------|-----------------|---------------------------------------|------------------|-----------------|------------------|------------------|------------------|-----------------|------------------|-----------------|------------------|------------------|---------------------------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | Week 12 | |
| | M | E | M | E | M | E | M | E | M | E | M | E | M | E |
| B11 | 518.4 (2.61) | 606.2 (3.70) | 516.4 (1.34) | 607.0 (4.53) | 510.4 (2.97) | 601.8 (10.50) | 504.4 (2.41) | 593.0 (2.55) | 503.4 (0.89) | 589.0 (5.57) | 503.0 (1.00) | 602.2 (5.76) | 498.4 (5.05) | 585.2 (9.07) |
| B12 | 510.6 (1.14) | 606.2 (6.34) | 515.0 (2.00) | 591.2 (2.28) | 512.2 (0.84) | 597.0 (5.43) | 504.4 (3.36) | 587.0 (5.79) | 506.2 (5.45) | 593.0 (6.96) | 503.8 (3.27) | 576.4 (11.01) | 502.8 (11.01) | 573.6 (1.10) (5.13) |
| B13 | 535.8 (1.10) | 621.2 (4.76) | 531.8 (3.11) | 608.4 (9.34) | 527.4 (3.21) | 606.8 (4.60) | 521.2 (3.49) | 612.6 (1.14) | 516.8 (3.42) | 587.6 (7.30) | 519.2 (1.30) | 587.2 (9.73) | 514.6 (2.51) | 585.6 (7.99) |
| B14 | 491.8 (3.27) | 580.4 (3.21) | 484.2 (2.17) | 575.8 (10.13) | 490.2 (1.10) | 570.8 (4.44) | 483.4 (2.88) | 575.4 (5.13) | 483.2 (0.84) | 562.6 (5.55) | 482.8 (0.45) | 566.8 (1.92) | 473.8 (2.59) | 576.2 (6.06) |
| B15 | 503.2 (1.64) | 604.4 (2.07) | 504.4 (1.82) | 586.0 (3.67) | 506.0 (1.58) | 594.0 (4.69) | 495.6 (2.88) | 577.0 (4.53) | 494.6 (0.55) | 564.4 (10.83) | 492.2 (1.10) | 579.0 (5.05) | 490.4 (2.07) | 561.2 (8.56) |
| B16 | 504.6 (2.07) | 617.2 (1.52) | 503.6 (5.94) | 610.6 (2.55) | 505.0 (8.47) | 628.6 (1.79) | 495.2 (13.59) | 611.2 (1.14) | 491.4 (6.08) | 618.0 (3.91) | 492.6 (3.85) | 604.4 (0.84) | 492.8 (8.23) | 604.4 (5.07) |
| B17 | 530.0 (2.00) | 617.2 (3.35) | 521.4 (1.82) | 610.8 (4.32) | 523.4 (2.70) | 600.6 (6.35) | 520.2 (3.35) | 604.0 (12.08) | 519.8 (1.30) | 601.2 (9.52) | 515.8 (1.64) | 602.4 (3.65) | 512.8 (3.91) | 591.8 (1.79) (5.07) |
| B18 | 508.4 (1.34) | 599.6 (2.97) | 509.0 (2.24) | 588.0 (5.70) | 502.0 (2.00) | 603.2 (5.02) | 496.6 (2.07) | 603.6 (9.15) | 493.8 (0.84) | 586.6 (7.64) | 494.4 (1.67) | 598.2 (10.33) | 494.0 (1.22) | 587.8 (4.49) |
| B19 | 519.4 (2.88) | 637.0 (1.73) | 525.2 (3.35) | 626.6 (5.13) | 504.2 (4.66) | 619.6 (14.60) | 507.4 (1.67) | 609.8 (11.21) | 503.2 (0.84) | 602.0 (3.54) | 490.4 (4.83) | 590.0 (6.56) | 489.6 (2.30) | 615.4 (8.29) |
| B20 | 495.2 (1.79) | 577.2 (2.49) | 493.4 (1.67) | 569.2 (1.48) | 490.2 (1.92) | 572.0 (3.39) | 490.0 (2.24) | 567.0 (11.83) | 491.4 (1.34) | 577.8 (4.55) | 486.0 (1.87) | 561.4 (6.88) | 482.0 (1.00) | 569.0 (13.25) |

Table 9 Skin measurement before and after applying propylene glycol (control) in volunteers group C. The data show average values ($n = 5$ measurements).

| Volunteer No. | Control group (left arm) | | | | | | | | | | Week 12 | | | | | | | | | |
|------------------|--------------------------|--------|---------|--------|---------|--------|---------|---------|---------|--------|---------|--------|---------|--------|---|--------|---|---|---|---|
| | Week 0 | | | | | Week 2 | | | | | Week 4 | | | | | Week 6 | | | | |
| | M | E | M | E | M | M | E | M | E | M | M | E | M | E | M | E | M | E | M | E |
| C 1 | 478.2 | 567.0 | 483.0 | 572.2 | 477.2 | 563.4 | 479.0 | 578.0 | 478.8 | 580.6 | 481.4 | 563.8 | 481.6 | 578.2 | | | | | | |
| (SD) | (2.77) | (6.89) | (0.71) | (2.59) | (2.95) | (9.56) | (3.74) | (11.38) | (1.48) | (5.41) | (3.05) | (8.73) | (2.07) | (6.22) | | | | | | |
| C 2 | 492.4 | 600.4 | 491.6 | 585.0 | 490.8 | 584.8 | 493.0 | 596.0 | 492.8 | 591.2 | 494.8 | 578.4 | 493.8 | 588.0 | | | | | | |
| (1.52) | (8.44) | (2.88) | (6.75) | (1.48) | (10.35) | (1.73) | (7.97) | (1.92) | (10.18) | (2.49) | (6.19) | (1.48) | (6.52) | | | | | | | |
| C 3 | 507.0 | 599.2 | 496.4 | 576.6 | 498.6 | 589.8 | 498.2 | 600.4 | 502.6 | 583.2 | 508.8 | 593.8 | 507.6 | 585.0 | | | | | | |
| (3.00) | (5.26) | (2.88) | (5.41) | (1.95) | (15.47) | (5.07) | (7.47) | (2.70) | (11.32) | (3.03) | (5.40) | (2.61) | (5.15) | | | | | | | |
| C 4 | 512.4 | 591.4 | 513.8 | 583.4 | 518.4 | 589.4 | 512.2 | 597.8 | 512.0 | 583.6 | 512.2 | 593.0 | 509.6 | 589.0 | | | | | | |
| (2.30) | (4.39) | (1.64) | (3.36) | (1.14) | (8.82) | (6.14) | (7.56) | (1.00) | (5.90) | (1.10) | (8.00) | (1.14) | (3.39) | | | | | | | |
| C 5 | 527.6 | 610.0 | 517.0 | 606.2 | 516.4 | 599.6 | 519.6 | 610.2 | 520.2 | 604.8 | 526.4 | 602.2 | 525.2 | 601.8 | | | | | | |
| (2.07) | (2.55) | (2.55) | (4.38) | (2.61) | (4.77) | (1.67) | (5.17) | (1.30) | (4.82) | (0.55) | (10.50) | (2.86) | (7.56) | | | | | | | |
| C 6 | 521.6 | 602.6 | 526.8 | 593.0 | 522.0 | 602.0 | 523.6 | 606.2 | 519.8 | 610.2 | 527.4 | 608.8 | 527.2 | 609.6 | | | | | | |
| (2.70) | (2.19) | (3.96) | (3.94) | (1.87) | (7.11) | (1.82) | (2.39) | (1.30) | (8.58) | (1.14) | (3.90) | (1.92) | (13.30) | | | | | | | |
| C 7 | 533.6 | 630.6 | 528.4 | 615.6 | 524.8 | 603.2 | 518.6 | 609.6 | 526.0 | 605.6 | 524.6 | 602.2 | 523.4 | 610.4 | | | | | | |
| (1.34) | (2.07) | (1.34) | (7.92) | (5.40) | (4.82) | (3.36) | (6.73) | (1.87) | (4.45) | (1.34) | (4.09) | (1.52) | (5.68) | | | | | | | |
| C 8 | 479.0 | 555.2 | 480.4 | 557.2 | 479.6 | 557.0 | 478.6 | 571.2 | 474.8 | 558.4 | 479.4 | 563.0 | 479.6 | 562.6 | | | | | | |
| (1.22) | (7.33) | (0.55) | (7.46) | (1.34) | (4.30) | (1.14) | (3.63) | (2.77) | (3.13) | (1.82) | (6.08) | (1.82) | (8.85) | | | | | | | |
| C 9 | 490.0 | 574.4 | 502.0 | 576.4 | 486.0 | 559.2 | 492.6 | 571.2 | 494.4 | 568.0 | 495.4 | 563.2 | 495.4 | 563.2 | | | | | | |
| (1.73) | (1.52) | (1.41) | (4.16) | (3.16) | (5.93) | (1.52) | (6.38) | (3.13) | (2.12) | (3.05) | (4.38) | (0.89) | (6.30) | | | | | | | |
| C 10 | 495.0 | 598.6 | 496.8 | 585.2 | 499.4 | 593.2 | 493.6 | 580.6 | 495.8 | 588.0 | 495.8 | 578.2 | 496.8 | 585.2 | | | | | | |
| (0.71) | (8.32) | (2.59) | (10.31) | (1.95) | (4.55) | (2.61) | (12.72) | (1.48) | (8.57) | (1.30) | (5.12) | (2.49) | (6.61) | | | | | | | |

Table 9 Skin measurement before and after applying propylene glycol (control) in volunteers group C. The data show average values ($n = 5$ measurements).

| Volunteer No. | Control group (right arm) | | | | | | | | | | | | | |
|------------------|---------------------------|-----------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | Week 12 | |
| | M | E | M | E | M | E | M | E | M | E | M | E | M | E |
| C 11 | 508.4 (SD) (1.14) | 589.4 (5.94) | 507.8 (1.30) | 584.8 (4.66) | 511.6 (2.19) | 584.8 (8.41) | 508.4 (1.82) | 509.4 (4.04) | 576.6 (2.07) | 581.2 (4.60) | 515.0 (1.58) | 573.2 (4.02) | 514.0 (1.22) | 586.0 (4.06) |
| C 12 | 524.2 (0.45) | 633.8 (6.57) | 525.4 (3.51) | 629.8 (10.06) | 521.8 (4.49) | 615.4 (9.63) | 516.6 (2.70) | 624.6 (4.22) | 518.6 (2.61) | 617.8 (6.38) | 516.2 (2.17) | 626.8 (11.52) | 515.8 (0.84) | 613.2 (6.02) |
| C 13 | 528.8 (2.17) | 616.2 (2.59) | 522.8 (1.30) | 608.6 (3.51) | 518.2 (2.17) | 614.2 (3.35) | 513.2 (4.15) | 604.8 (3.05) | 517.6 (9.68) | 606.2 (2.74) | 522.0 (7.27) | 601.4 (3.49) | 522.8 (5.40) | 614.8 |
| C 14 | 496.0 (1.58) | 609.4 (1.95) | 483.2 (1.10) | 581.0 (3.67) | 484.8 (4.76) | 562.8 (10.43) | 484.4 (4.39) | 576.0 (12.94) | 488.6 (2.88) | 585.2 (4.09) | 490.6 (0.55) | 587.0 (6.82) | 490.2 (2.28) | 575.4 (7.30) |
| C 15 | 498.8 (2.59) | 591.8 (5.85) | 498.8 (1.30) | 586.6 (6.50) | 497.2 (9.96) | 601.4 (9.45) | 493.2 (3.49) | 586.2 (4.66) | 495.0 (3.61) | 594.4 (10.33) | 496.4 (2.51) | 592.2 (2.17) | 498.2 (1.10) | 569.4 (7.30) |
| C 16 | 517.0 (1.58) | 587.8 (4.82) | 518.6 (1.52) | 600.4 (3.65) | 521.8 (1.30) | 590.8 (4.32) | 522.0 (1.00) | 590.0 (8.89) | 527.4 (3.58) | 596.0 (5.70) | 529.0 (1.22) | 592.2 (4.83) | 528.2 (1.30) | 590.2 (7.29) |
| C 17 | 538.2 (1.30) | 609.6 (6.84) | 531.0 (1.22) | 601.2 (3.27) | 517.4 (4.22) | 587.0 (5.70) | 512.4 (5.41) | 596.2 (4.55) | 518.4 (1.52) | 594.6 (7.16) | 525.0 (3.00) | 599.8 (8.56) | 523.4 (2.70) | 610.2 (10.83) |
| C 18 | 503.2 (1.64) | 591.6 (2.61) | 501.4 (5.18) | 614.2 (7.92) | 498.6 (2.07) | 589.6 (6.99) | 501.6 (1.52) | 593.2 (8.70) | 497.4 (1.67) | 577.4 (9.99) | 505.6 (2.79) | 588.0 (11.90) | 501.4 (2.19) | 577.0 (4.64) |
| C 19 | 519.0 (6.40) | 603.6 (5.50) | 517.2 (1.14) | 597.4 (3.67) | 511.0 (8.38) | 598.6 (1.82) | 510.4 (4.55) | 601.2 (4.55) | 502.2 (1.79) | 573.0 (8.03) | 501.4 (4.56) | 586.4 (9.02) | 502.2 (1.79) | 589.8 (1.64) |
| C 20 | 496.4 (3.51) | 595.6 (4.98) | 496.8 (5.07) | 600.4 (7.83) | 498.4 (5.03) | 583.8 (5.72) | 491.0 (1.87) | 586.0 (6.89) | 491.8 (1.64) | 578.0 (4.69) | 497.2 (3.11) | 580.8 (7.29) | 498.2 (3.83) | 592.6 (7.92) |

Table 10 Skin measurement before and after applying 0.25 % w/v Licorice extract in volunteers group C. The data show average values ($n = 5$ measurements).

| Volunteer No. | 0.25 % w/v Licorice extract group (right arm) | | | | | | | | | | | | | |
|------------------|---|--------|---------|--------|---------|---------|---------|--------|---------|--------|---------|---------|---------|--------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | Week 12 | |
| | M | E | M | E | M | E | M | E | M | E | M | E | M | E |
| C 1 | 479.2 | 582.0 | 477.4 | 571.2 | 473.6 | 575.8 | 473.4 | 579.8 | 476.8 | 574.0 | 472.4 | 557.2 | 470.6 | 573.2 |
| (SD) | (1.30) | (3.81) | (2.07) | (6.91) | (2.41) | (10.11) | (1.95) | (9.23) | (2.17) | (6.16) | (1.52) | (10.99) | (0.55) | (5.54) |
| C 2 | 499.8 | 602.0 | 494.6 | 582.0 | 499.0 | 586.4 | 496.0 | 591.2 | 491.4 | 582.6 | 490.4 | 584.0 | 490.4 | 590.2 |
| (0.45) | (5.05) | (0.89) | (6.75) | (4.00) | (4.56) | (1.73) | (7.22) | (3.29) | (14.06) | (1.95) | (4.95) | (0.55) | (9.65) | |
| C 3 | 503.2 | 603.4 | 492.4 | 578.8 | 494.6 | 592.4 | 489.2 | 592.0 | 494.0 | 580.6 | 501.2 | 578.6 | 498.2 | 576.6 |
| (0.45) | (5.18) | (2.07) | (11.71) | (2.88) | (15.19) | (2.39) | (14.00) | (1.58) | (2.88) | (1.48) | (11.26) | (2.49) | (11.97) | |
| C 4 | 518.2 | 584.6 | 515.0 | 582.2 | 513.2 | 577.0 | 510.2 | 587.0 | 510.0 | 578.0 | 511.6 | 577.6 | 506.2 | 573.4 |
| (3.83) | (4.51) | (0.71) | (3.77) | (2.39) | (4.42) | (3.35) | (8.63) | (1.00) | (4.64) | (2.07) | (6.66) | (1.30) | (6.11) | |
| C 5 | 538.8 | 625.6 | 528.4 | 601.8 | 526.2 | 609.2 | 521.0 | 606.2 | 527.4 | 607.8 | 519.8 | 597.4 | 518.6 | 593.8 |
| (3.96) | (7.47) | (3.29) | (3.96) | (1.79) | (5.89) | (1.22) | (6.98) | (0.89) | (8.35) | (1.64) | (5.90) | (1.14) | (5.89) | |
| C 6 | 522.8 | 600.0 | 531.6 | 604.8 | 522.2 | 591.6 | 524.2 | 603.2 | 516.6 | 600.0 | 512.6 | 591.4 | 513.8 | 589.8 |
| (1.10) | (4.95) | (2.07) | (6.14) | (3.70) | (6.80) | (1.64) | (7.60) | (1.82) | (4.74) | (3.51) | (8.20) | (1.79) | (5.72) | |
| C 7 | 539.0 | 637.0 | 532.6 | 621.0 | 524.0 | 614.0 | 508.2 | 608.0 | 517.2 | 607.2 | 517.0 | 611.4 | 512.4 | 598.2 |
| (1.58) | (7.65) | (3.05) | (8.46) | (2.24) | (8.22) | (3.03) | (7.18) | (1.64) | (7.16) | (0.71) | (3.05) | (4.93) | (8.50) | |
| C 8 | 486.6 | 568.6 | 484.0 | 563.8 | 483.6 | 558.0 | 481.2 | 575.2 | 482.8 | 572.8 | 481.6 | 563.6 | 479.6 | 561.6 |
| (1.14) | (6.19) | (1.00) | (8.35) | (2.30) | (4.74) | (0.84) | (6.34) | (1.79) | (4.09) | (1.95) | (3.78) | (0.55) | (10.04) | |
| C 9 | 493.4 | 561.8 | 490.2 | 576.8 | 488.6 | 554.4 | 485.0 | 563.2 | 484.6 | 555.6 | 488.4 | 556.8 | 487.6 | 553.2 |
| (1.95) | (6.94) | (0.84) | (3.03) | (2.41) | (4.67) | (3.81) | (11.19) | (1.52) | (3.36) | (0.55) | (6.26) | (0.55) | (7.19) | |
| C 10 | 494.4 | 606.8 | 498.0 | 587.0 | 502.4 | 594.4 | 494.2 | 589.8 | 495.6 | 583.8 | 493.6 | 568.4 | 491.6 | 575.2 |
| (0.89) | (7.60) | (2.00) | (6.63) | (2.19) | (7.16) | (2.68) | (7.66) | (0.89) | (7.12) | (2.30) | (7.47) | (2.41) | (4.02) | |

Table 10 Skin measurement before and after applying 0.25 % w/v Licorice extract in volunteers group C. The data show average values ($n = 5$ measurements).

Table 11 Skin measurement before and after applying propylene glycol (control) in volunteers group D. The data show average values ($n = 5$ measurements).

| Volunteer No. | Control group (left arm) | | | | | | | | | | | | | |
|------------------|--------------------------|--------|--------|--------|---------|---------|---------|--------|---------|--------|---------|--------|---------|--------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | Week 12 | |
| | M | E | M | E | M | E | M | E | M | E | M | E | M | E |
| D 1 | 480.6 | 597.8 | 485.0 | 563.8 | 484.6 | 587.4 | 479.0 | 581.6 | 479.2 | 579.0 | 484.4 | 587.8 | 482.8 | 584.4 |
| (SD) | (4.77) | (3.96) | (1.22) | (3.42) | (1.95) | (12.50) | (1.73) | (5.27) | (1.92) | (5.34) | (1.34) | (7.98) | (1.79) | (1.52) |
| D 2 | 494.0 | 601.4 | 494.4 | 598.2 | 491.0 | 579.4 | 492.6 | 591.6 | 498.6 | 585.8 | 497.4 | 589.4 | 497.4 | 582.2 |
| (1.58) | (8.20) | (1.14) | (4.66) | (1.87) | (11.46) | (1.52) | (9.29) | (2.70) | (8.79) | (4.16) | (9.81) | (1.52) | (9.86) | |
| D 3 | 509.8 | 600.0 | 508.2 | 593.6 | 504.4 | 592.8 | 499.6 | 593.4 | 500.0 | 585.8 | 501.0 | 574.6 | 498.8 | 587.2 |
| (2.28) | (7.58) | (1.79) | (3.29) | (1.14) | (3.96) | (0.55) | (2.07) | (1.00) | (5.26) | (1.22) | (12.42) | (5.54) | (7.16) | |
| D 4 | 510.0 | 614.6 | 510.2 | 595.4 | 503.0 | 599.6 | 503.4 | 603.0 | 507.0 | 600.0 | 507.4 | 596.4 | 508.2 | 595.2 |
| (2.35) | (5.03) | (1.79) | (4.16) | (2.00) | (4.51) | (3.13) | (6.24) | (1.41) | (9.06) | (1.82) | (2.88) | (3.96) | (3.96) | (8.35) |
| D 5 | 512.0 | 603.0 | 513.4 | 611.2 | 510.0 | 578.4 | 509.4 | 595.0 | 507.6 | 582.4 | 509.4 | 593.6 | 508.4 | 582.8 |
| (1.73) | (4.53) | (2.30) | (1.92) | (4.36) | (4.98) | (3.85) | (7.42) | (1.67) | (6.80) | (1.14) | (4.98) | (2.70) | (4.87) | |
| D 6 | 530.2 | 602.4 | 531.0 | 624.6 | 523.2 | 593.4 | 523.2 | 603.2 | 518.2 | 615.4 | 532.6 | 600.4 | 531.8 | 623.0 |
| (1.64) | (1.14) | (1.41) | (4.39) | (0.84) | (6.77) | (2.17) | (5.76) | (3.90) | (10.41) | (2.19) | (3.44) | (2.17) | (7.42) | |
| D 7 | 498.0 | 595.6 | 490.4 | 592.8 | 490.0 | 589.6 | 489.0 | 592.6 | 494.8 | 594.4 | 492.0 | 589.0 | 489.4 | 582.0 |
| (2.92) | (6.99) | (2.07) | (2.59) | (1.22) | (5.03) | (2.00) | (11.93) | (0.84) | (12.54) | (1.22) | (4.06) | (1.14) | (4.53) | |
| D 8 | 501.2 | 582.8 | 503.4 | 580.6 | 502.8 | 575.8 | 498.8 | 580.8 | 499.4 | 558.6 | 500.8 | 567.0 | 504.0 | 570.8 |
| (1.92) | (4.02) | (2.70) | (9.99) | (0.84) | (1.79) | (2.59) | (3.56) | (0.89) | (7.92) | (1.92) | (3.24) | (3.32) | (4.32) | |
| D 9 | 512.2 | 583.2 | 506.0 | 585.6 | 506.6 | 579.4 | 510.6 | 591.4 | 511.0 | 592.2 | 511.4 | 584.4 | 512.8 | 580.4 |
| (1.30) | (1.64) | (1.87) | (3.97) | (2.41) | (6.11) | (3.21) | (5.77) | (2.35) | (3.11) | (0.89) | (2.19) | (1.10) | (3.21) | |
| D 10 | 522.4 | 627.4 | 517.2 | 627.4 | 509.2 | 601.8 | 512.0 | 602.6 | 513.8 | 598.4 | 512.0 | 608.4 | 511.8 | 606.2 |
| (0.55) | (2.61) | (2.28) | (5.64) | (2.95) | (5.72) | (0.71) | (3.13) | (2.49) | (8.50) | (1.22) | (5.98) | (2.17) | (2.49) | |

Table 11 Skin measurement before and after applying propylene glycol (control) in volunteers group D. The data show average values ($n = 5$ measurements).

| No. | Volunteer | | Control group (right arm) | | | | | | | | | | | |
|--------|-----------|--------|---------------------------|--------|---------|--------|--------|--------|---------|---------|---------|--------|---------|--------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | Week 12 | |
| | M | E | M | E | M | E | M | E | M | E | M | E | M | E |
| D11 | 529.2 | 628.6 | 529.6 | 627.0 | 528.0 | 624.0 | 529.8 | 620.6 | 525.2 | 618.0 | 528.2 | 616.8 | 522.4 | 618.8 |
| (SD) | (1.79) | (7.23) | (1.82) | (5.66) | (2.65) | (3.08) | (3.11) | (2.88) | (2.17) | (6.60) | (1.64) | (2.39) | (4.67) | (4.76) |
| D12 | 500.2 | 573.8 | 501.8 | 581.4 | 507.2 | 581.0 | 505.2 | 582.2 | 504.8 | 573.4 | 504.8 | 568.2 | 505.4 | 573.0 |
| (1.30) | (2.95) | (2.17) | (6.23) | (2.59) | (5.70) | (1.64) | (2.77) | (2.59) | (5.59) | (1.48) | (5.59) | (5.81) | (2.97) | (2.55) |
| D13 | 529.2 | 619.2 | 537.2 | 615.6 | 536.4 | 621.4 | 536.4 | 620.0 | 531.6 | 606.2 | 533.8 | 614.2 | 533.2 | 605.0 |
| (2.17) | (3.42) | (3.27) | (6.02) | (3.36) | (8.44) | (0.55) | (3.46) | (3.46) | (2.79) | (17.95) | (1.30) | (3.19) | (3.11) | (5.39) |
| D14 | 522.2 | 626.0 | 520.8 | 611.2 | 515.6 | 601.6 | 513.2 | 613.8 | 520.4 | 611.6 | 520.6 | 606.8 | 521.0 | 615.8 |
| (4.09) | (4.24) | (2.39) | (7.22) | (2.19) | (2.61) | (2.28) | (2.28) | (2.28) | (1.67) | (5.27) | (2.30) | (6.57) | (1.41) | (6.30) |
| D16 | 507.4 | 567.4 | 506.0 | 565.6 | 512.0 | 573.6 | 511.2 | 567.8 | 509.0 | 566.8 | 508.6 | 562.8 | 507.2 | 563.2 |
| (1.52) | (5.37) | (2.65) | (5.46) | (2.65) | (3.65) | (3.70) | (5.12) | (2.35) | (5.07) | (1.67) | (3.42) | (1.64) | (3.42) | |
| D17 | 510.4 | 612.6 | 517.2 | 613.2 | 521.8 | 621.4 | 517.0 | 610.2 | 514.2 | 591.6 | 520.4 | 607.4 | 519.4 | 604.6 |
| (1.52) | (6.23) | (2.77) | (2.59) | (4.44) | (1.67) | (1.87) | (8.53) | (3.77) | (9.45) | (2.51) | (2.07) | (2.70) | (5.32) | |
| D18 | 503.6 | 601.8 | 502.2 | 588.6 | 495.0 | 589.4 | 501.6 | 582.8 | 504.2 | 589.4 | 505.4 | 589.8 | 503.8 | 599.2 |
| (2.41) | (4.09) | (0.45) | (6.07) | (8.57) | (9.04) | (3.05) | (5.54) | (2.28) | (5.59) | (2.07) | (11.03) | (2.49) | (5.54) | |
| D19 | 535.4 | 617.8 | 529.4 | 611.2 | 523.0 | 603.2 | 524.2 | 603.2 | 523.4 | 582.2 | 518.8 | 600.6 | 520.8 | 593.6 |
| (2.19) | (8.67) | (4.77) | (11.50) | (1.22) | (10.40) | (1.30) | (8.04) | (2.41) | (10.11) | (1.10) | (6.99) | (2.68) | (5.08) | |
| D20 | 498.4 | 621.8 | 501.0 | 610.2 | 499.2 | 601.4 | 494.6 | 611.6 | 494.0 | 602.2 | 498.8 | 611.0 | 497.6 | 608.4 |
| (2.07) | (6.22) | (1.22) | (5.81) | (2.28) | (9.32) | (2.19) | (1.14) | (1.58) | (9.98) | (2.59) | (5.57) | (0.89) | (7.60) | |

Table 12 Skin measurement before and after applying 3.00 % w/v Kojic acid in volunteers group D. The data show average values ($n = 5$ measurements).

| Volunteer No. | 3.00 % w/v Kojic acid group (right arm) | | | | | | | | | | | | | |
|------------------|---|--------|--------|--------|--------|---------|---------|--------|---------|---------|---------|--------|---------|--------|
| | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | Week 12 | |
| | M | E | M | E | M | E | M | E | M | E | M | E | M | E |
| D 1 | 480.8 | 593.4 | 477.4 | 578.0 | 481.2 | 580.4 | 476.6 | 580.0 | 479.2 | 580.0 | 483.0 | 585.8 | 482.2 | 580.0 |
| (SD) | (0.84) | (1.14) | (2.51) | (4.00) | (3.56) | (11.13) | (2.41) | (6.00) | (0.45) | (13.66) | (1.87) | (5.26) | (1.92) | (9.97) |
| D 2 | 492.8 | 594.8 | 497.2 | 609.2 | 488.2 | 578.8 | 488.8 | 590.6 | 493.4 | 578.4 | 492.4 | 613.6 | 489.0 | 572.8 |
| (1.64) | (5.89) | (3.35) | (4.82) | (0.84) | (5.89) | (1.10) | (8.32) | (1.52) | (1.95) | (1.34) | (7.23) | (1.87) | (9.71) | |
| D 3 | 507.8 | 604.2 | 507.4 | 593.4 | 505.2 | 592.6 | 500.0 | 608.0 | 499.0 | 587.6 | 497.6 | 588.4 | 494.0 | 588.8 |
| (2.17) | (10.40) | (3.13) | (7.09) | (1.30) | (5.27) | (3.08) | (6.36) | (1.58) | (8.53) | (1.14) | (9.56) | (2.74) | (7.05) | |
| D 4 | 515.6 | 616.4 | 506.4 | 598.0 | 507.4 | 602.8 | 503.4 | 597.8 | 499.4 | 587.2 | 495.6 | 594.2 | 499.6 | 597.4 |
| (3.36) | (5.90) | (2.70) | (6.20) | (1.95) | (8.79) | (3.29) | (4.76) | (2.41) | (10.52) | (2.51) | (5.93) | (3.29) | (5.98) | |
| D 5 | 519.2 | 608.6 | 517.4 | 610.4 | 512.0 | 587.8 | 505.0 | 589.2 | 508.8 | 587.4 | 508.8 | 589.8 | 505.2 | 575.0 |
| (1.30) | (4.16) | (3.65) | (7.60) | (2.92) | (3.56) | (4.06) | (4.09) | (0.84) | (5.90) | (0.45) | (2.86) | (2.49) | (1.73) | |
| D 6 | 534.2 | 613.8 | 526.2 | 603.4 | 522.6 | 597.6 | 514.4 | 602.8 | 513.4 | 601.8 | 512.6 | 588.2 | 510.4 | 585.4 |
| (1.64) | (3.27) | (3.83) | (6.80) | (1.52) | (6.58) | (2.88) | (2.95) | (0.89) | (2.86) | (1.14) | (2.77) | (1.67) | (2.07) | |
| D 7 | 488.8 | 593.8 | 489.6 | 584.4 | 488.2 | 576.0 | 482.8 | 593.6 | 487.0 | 579.6 | 481.0 | 588.0 | 480.2 | 580.0 |
| (1.64) | (9.98) | (2.07) | (3.51) | (2.39) | (6.63) | (1.64) | (5.13) | (0.71) | (3.85) | (1.58) | (2.00) | (0.84) | (6.40) | |
| D 8 | 503.6 | 583.2 | 504.0 | 581.8 | 504.6 | 580.4 | 500.6 | 580.8 | 501.4 | 566.4 | 506.4 | 580.0 | 501.6 | 570.6 |
| (1.52) | (3.56) | (1.58) | (8.01) | (0.89) | (6.95) | (1.82) | (8.38) | (1.95) | (7.27) | (1.14) | (3.61) | (4.72) | (9.50) | |
| D 9 | 525.6 | 595.8 | 511.4 | 582.8 | 514.4 | 581.6 | 513.6 | 585.2 | 511.2 | 582.0 | 508.0 | 583.6 | 506.4 | 564.8 |
| (2.07) | (5.63) | (3.58) | (2.49) | (3.21) | (6.43) | (4.04) | (3.96) | (2.39) | (6.32) | (1.41) | (6.27) | (1.67) | (2.17) | |
| D 10 | 526.2 | 636.2 | 513.8 | 626.0 | 516.2 | 601.6 | 513.2 | 602.4 | 508.6 | 593.4 | 508.2 | 589.6 | 505.2 | 594.2 |
| (0.84) | (2.59) | (4.15) | (4.85) | (3.83) | (5.73) | (3.96) | (10.55) | (2.88) | (5.59) | (1.30) | (7.40) | (4.27) | (6.38) | |

Table 12 Skin measurement before and after applying 3.00 % w/v Kojic acid in volunteers group A. The data show average values (n = 5 measurements).

| No. | Volunteer | 3.00 % w/v Kojic acid group (left arm) | | | | | | | | | | Week 12 | | | |
|--------|-----------|--|--------|--------|--------|--------|--------|--------|---------|--------|--------|---------|--------|---------|---|
| | | Week 0 | | Week 2 | | Week 4 | | Week 6 | | Week 8 | | Week 10 | | Week 12 | |
| | | M | E | M | E | M | E | M | E | M | E | M | E | M | E |
| D11 | 531.8 | 629.0 | 534.4 | 632.6 | 523.6 | 622.6 | 523.6 | 614.4 | 522.4 | 609.6 | 517.6 | 604.4 | 517.6 | 624.4 | |
| (SD) | (2.28) | (7.68) | (1.82) | (2.07) | (2.30) | (5.03) | (1.67) | (6.47) | (1.14) | (5.90) | (2.19) | (9.71) | (1.14) | (8.62) | |
| D12 | 500.4 | 580.2 | 496.6 | 580.4 | 499.0 | 573.2 | 493.4 | 573.8 | 492.6 | 553.0 | 490.0 | 552.8 | 490.0 | 555.0 | |
| (2.88) | (3.35) | (4.04) | (9.96) | (1.87) | (5.54) | (2.51) | (6.14) | (1.82) | (10.42) | (2.55) | (4.97) | (1.41) | (3.00) | | |
| D13 | 542.8 | 629.8 | 552.8 | 630.4 | 544.6 | 616.8 | 540.0 | 622.0 | 546.4 | 604.8 | 534.8 | 608.2 | 530.4 | 598.8 | |
| (2.77) | (3.56) | (3.49) | (1.67) | (4.34) | (4.97) | (2.74) | (4.85) | (2.74) | (4.85) | (2.30) | (5.45) | (3.11) | (3.70) | (5.12) | |
| D14 | 523.6 | 628.2 | 522.6 | 608.6 | 511.8 | 600.6 | 512.8 | 597.4 | 511.8 | 595.8 | 512.0 | 592.0 | 502.2 | 581.6 | |
| (2.88) | (5.93) | (1.95) | (1.95) | (2.59) | (7.09) | (2.95) | (7.09) | (2.95) | (7.09) | (1.10) | (6.06) | (1.00) | (9.27) | (2.59) | |
| D16 | 501.4 | 571.0 | 506.0 | 566.4 | 504.8 | 567.2 | 498.6 | 565.6 | 498.4 | 559.8 | 497.8 | 564.2 | 494.0 | 568.8 | |
| (1.95) | (4.85) | (1.41) | (3.91) | (2.86) | (3.49) | (1.14) | (1.14) | (1.14) | (1.14) | (2.07) | (3.35) | (1.64) | (2.68) | (9.12) | |
| D17 | 510.4 | 601.0 | 514.8 | 610.6 | 517.8 | 600.0 | 502.6 | 597.4 | 507.8 | 595.0 | 506.8 | 609.4 | 505.4 | 587.8 | |
| (2.30) | (6.04) | (4.21) | (6.95) | (2.95) | (9.41) | (2.30) | (6.39) | (3.11) | (10.70) | (1.64) | (9.40) | (0.55) | (6.53) | | |
| D18 | 497.6 | 593.4 | 500.2 | 587.0 | 500.6 | 583.8 | 499.8 | 583.4 | 497.6 | 579.6 | 497.8 | 579.0 | 495.6 | 580.0 | |
| (3.65) | (3.91) | (4.60) | (3.54) | (4.39) | (9.28) | (1.64) | (2.19) | (1.34) | (7.96) | (0.84) | (9.62) | (0.89) | (3.08) | | |
| D19 | 527.2 | 618.6 | 529.6 | 607.4 | 523.0 | 608.0 | 515.8 | 590.0 | 518.2 | 590.4 | 514.4 | 589.6 | 511.8 | 593.4 | |
| (3.90) | (3.78) | (4.39) | (5.13) | (3.46) | (7.11) | (3.83) | (5.34) | (1.64) | (4.93) | (1.82) | (6.84) | (3.03) | (8.08) | | |
| D20 | 501.4 | 613.8 | 502.2 | 618.2 | 498.6 | 604.8 | 496.0 | 602.2 | 494.2 | 605.8 | 492.2 | 603.8 | 490.2 | 603.8 | |
| (0.89) | (9.58) | (0.45) | (1.64) | (2.88) | (8.35) | (1.22) | (7.56) | (2.77) | (14.10) | (1.10) | (5.26) | (2.05) | (4.21) | | |

Appendix IIc

Individual %Whitening of female volunteers for

0.50% Puag-Haad (group A)

0.25% Puag-Haad (group B)

0.25 % Licorice extract (group C)

3.00% Kojic acid (group D)

Skin color was evaluated on outside areas of upper arms with Mexameter MX 16 described in methods.

Table 13 The % whitening values in the upper arms of volunteers (group A) treated with 0.5 % Puag-Haad and Propylene glycol for 12 weeks.

| Volunteer No. | % Whitening (0.5 % Puag-Haad) | | | | | | % Whitening (Propylene glycol) | | | | | |
|---------------|-------------------------------|--------|--------|--------|---------|---------|--------------------------------|--------|--------|--------|---------|---------|
| | week 2 | week 4 | week 6 | week 8 | week 10 | week 12 | week 2 | week 4 | week 6 | week 8 | week 10 | week 12 |
| A1 | 0.95 | 0.12 | 0.70 | 0.58 | -0.82 | 0.33 | 1.06 | 0.82 | 0.53 | -0.49 | -0.37 | 0.61 |
| A2 | -1.77 | -0.45 | -0.25 | -1.52 | -0.86 | -0.70 | -0.36 | -0.32 | -0.24 | -0.20 | -1.54 | -1.50 |
| A3 | 1.11 | 0.59 | 2.10 | 1.70 | 0.91 | 1.31 | 0.28 | -0.28 | -0.36 | -0.24 | -0.40 | -0.32 |
| A4 | 0.43 | 1.37 | 1.77 | 1.77 | 2.24 | 2.36 | 1.35 | 3.16 | 2.74 | 2.04 | 1.62 | 1.66 |
| A5 | -0.27 | 0.81 | 0.08 | -1.00 | 1.04 | 0.73 | 0.26 | 0.45 | 1.62 | 1.36 | 1.51 | 1.81 |
| A6 | 0.26 | 1.82 | 2.57 | 2.64 | 2.49 | 2.83 | 0.48 | 0.93 | 1.19 | 0.89 | 0.63 | 1.00 |
| A7 | 0.60 | 0.79 | 1.53 | 0.26 | 2.58 | 2.66 | -1.75 | -1.19 | -1.16 | -1.12 | -1.38 | -1.60 |
| A8 | -1.19 | -0.76 | 0.55 | -0.89 | -0.76 | -0.51 | -1.14 | -1.52 | -1.99 | -1.27 | -1.74 | -1.82 |
| A10 | -0.81 | 1.09 | 2.51 | 1.05 | 0.73 | 1.17 | -0.67 | 2.29 | 2.68 | 3.00 | 2.25 | 2.49 |
| A11 | 0.27 | 1.97 | 2.51 | 3.36 | 2.78 | 2.63 | 0.85 | 1.63 | 2.48 | 1.63 | 1.78 | 1.47 |
| A12 | 1.46 | 0.75 | 1.81 | 2.64 | 2.21 | 2.68 | -0.40 | -1.00 | -0.52 | -0.32 | -0.96 | -1.12 |
| A13 | 0.27 | 0.04 | 0.96 | 1.00 | 1.46 | 2.30 | 0.91 | 0.42 | 0.84 | 0.23 | 0.46 | 1.79 |
| A14 | 0.32 | -0.24 | -0.32 | 0.76 | 0.68 | 1.75 | -0.24 | 0.08 | -0.53 | -0.44 | -0.93 | -0.69 |
| A15 | -0.16 | 0.70 | 1.86 | 2.17 | 3.65 | 4.11 | 1.64 | 0.31 | 1.88 | 1.45 | 1.52 | 1.52 |
| A16 | -0.50 | 0.50 | 1.15 | 2.19 | 2.73 | 4.53 | 1.97 | 1.59 | 2.69 | 1.97 | 2.53 | 2.35 |
| A17 | -1.73 | -1.96 | -0.88 | -1.81 | -1.19 | -0.73 | -1.48 | -1.02 | -0.64 | -0.34 | -0.57 | -0.57 |
| A19 | 1.59 | 1.21 | 1.93 | 1.85 | 1.85 | 2.34 | 1.60 | 0.53 | 0.53 | 0.69 | 0.69 | 2.32 |
| Average | 0.05 | 0.49 | 1.21 | 0.99 | 1.28 | 1.75 | 0.26 | 0.40 | 0.69 | 0.52 | 0.30 | 0.55 |
| (SD) | (1.00) | (0.98) | (1.08) | (1.54) | (1.49) | (1.56) | (1.12) | (1.27) | (1.48) | (1.22) | (1.39) | (1.53) |

Table 14 The % whitening values in the upper arms of volunteers (group B) treated with 0.25 % Puag-Haad and Propylene glycol for 12 weeks.

| Volunteer No. | % Whitening (0.25 % Puag-Haad) | | | | | | % Whitening (Propylene glycol) | | | | | |
|------------------|--------------------------------|--------|--------|--------|---------|---------|--------------------------------|--------|--------|--------|---------|---------|
| | week 2 | week 4 | week 6 | week 8 | week 10 | week 12 | week 2 | week 4 | week 6 | week 8 | week 10 | week 12 |
| B1 | -0.29 | -0.29 | 0.87 | 1.29 | 0.83 | 1.21 | 0.33 | -1.08 | 0.04 | 0.12 | -0.29 | -0.17 |
| B2 | 0.20 | 2.19 | 1.90 | 1.90 | 2.23 | -0.12 | 1.41 | 1.82 | 1.41 | 1.74 | 1.29 | |
| B3 | -0.04 | 0.60 | 0.76 | 1.97 | 1.13 | 1.85 | -0.24 | 0.04 | 0.12 | -0.28 | -0.20 | -0.12 |
| B4 | -0.27 | 0.51 | 0.59 | 0.43 | 1.33 | 1.10 | -0.35 | 1.10 | 0.63 | 0.43 | 0.43 | 0.63 |
| B5 | 0.76 | 1.52 | 3.50 | 3.69 | 3.84 | 3.95 | 0.87 | 1.78 | 2.77 | 2.24 | 2.51 | 2.81 |
| B6 | 0.56 | 2.97 | 3.05 | 3.79 | 5.42 | 5.72 | 1.16 | 1.08 | 1.72 | 1.57 | 2.16 | 2.28 |
| B7 | 1.07 | 0.88 | 2.03 | 2.36 | 3.17 | 3.46 | 1.43 | 2.12 | 3.37 | 1.79 | 2.60 | 2.93 |
| B8 | -0.21 | 1.69 | 1.90 | 2.64 | 1.90 | 2.68 | -0.25 | 0.75 | -0.46 | -0.04 | -0.71 | -0.50 |
| B9 | -0.16 | 0.44 | 0.81 | 0.77 | -0.24 | 0.73 | -0.20 | -0.77 | -0.57 | -0.53 | -1.06 | -0.94 |
| B10 | -0.48 | -0.40 | -0.12 | -0.56 | 0.40 | 0.36 | -1.71 | -1.39 | -0.88 | -0.76 | -0.92 | -0.48 |
| B11 | 0.39 | 1.54 | 2.70 | 2.89 | 2.97 | 3.86 | 1.48 | 0.70 | 0.78 | 0.90 | 0.62 | 0.51 |
| B12 | -0.86 | -0.31 | 1.21 | 0.86 | 1.33 | 1.53 | -1.62 | -1.21 | -1.36 | -0.90 | -0.45 | -0.11 |
| B13 | 0.75 | 1.57 | 2.72 | 3.55 | 3.10 | 3.96 | -0.15 | 0.71 | 1.27 | 2.54 | 1.12 | 0.93 |
| B14 | 1.55 | 0.33 | 1.71 | 1.75 | 1.83 | 3.66 | -0.49 | -0.08 | 0.73 | 1.55 | 1.71 | 1.55 |
| B15 | -0.24 | -0.56 | 1.51 | 1.71 | 2.19 | 2.54 | -0.39 | -1.03 | 1.78 | 1.03 | 0.59 | 1.10 |
| B16 | 0.20 | -0.08 | 1.86 | 2.62 | 2.38 | 2.34 | 0.67 | -0.63 | 0.75 | 1.22 | -0.20 | 0.16 |
| B17 | 1.62 | 1.25 | 1.85 | 1.92 | 2.68 | 3.25 | 0.91 | 0.68 | 0.83 | 0.76 | 1.17 | 1.55 |
| B18 | -0.12 | 1.26 | 2.32 | 2.87 | 2.75 | 2.83 | 1.09 | 0.16 | 0.39 | 0.51 | 0.00 | 0.43 |
| B19 | -1.12 | 2.93 | 2.31 | 3.12 | 5.58 | 5.74 | -1.15 | 0.11 | 0.00 | -0.57 | 0.23 | 0.23 |
| B20 | 0.36 | 1.01 | 1.05 | 0.77 | 1.86 | 2.67 | -1.17 | -1.01 | -0.12 | 0.40 | -1.21 | -1.17 |
| Average | 0.18 | 0.95 | 1.73 | 2.02 | 2.32 | 2.78 | 0.00 | 0.17 | 0.68 | 0.67 | 0.49 | 0.65 |
| (SD) | (0.72) | (1.04) | (0.91) | (1.17) | (1.47) | (1.47) | (0.97) | (1.05) | (1.19) | (1.01) | (1.18) | (1.16) |

Table 15 The % whitening values in the upper arms of volunteers (group C) treated with 0.25 % Licorice extract and Propylene glycol for 12 weeks.

| Volunteer No. | % Whitening (0.25 % Licorice extract) | | | | | | % Whitening (Propylene glycol) | | | | | |
|---------------|---------------------------------------|-------------|-------------|-------------|-------------|-------------|--------------------------------|-------------|-------------|-------------|-------------|-------------|
| | week 2 | week 4 | week 6 | week 8 | week 10 | week 12 | week 2 | week 4 | week 6 | week 8 | week 10 | week 12 |
| C1 | 0.38 | 1.17 | 1.21 | 0.50 | 1.42 | 1.79 | -1.00 | 0.21 | -0.17 | -0.13 | -0.67 | -0.71 |
| C2 | 1.04 | 0.16 | 0.76 | 1.68 | 1.88 | 0.88 | 0.16 | 0.32 | -0.12 | -0.08 | -0.49 | -0.28 |
| C3 | 2.15 | 1.71 | 2.78 | 1.83 | 0.40 | 0.99 | 2.09 | 1.66 | 1.74 | 0.87 | -0.36 | -0.12 |
| C4 | 0.62 | 0.96 | 1.54 | 1.58 | 1.27 | 2.32 | -0.27 | -1.17 | 0.04 | 0.08 | 0.04 | 0.55 |
| C5 | 1.93 | 2.34 | 3.30 | 2.12 | 3.53 | 3.75 | 2.01 | 2.12 | 1.52 | 1.40 | 0.23 | 0.45 |
| C6 | -1.68 | 0.11 | -0.27 | 1.19 | 1.95 | 1.72 | -1.00 | -0.08 | -0.38 | 0.35 | -1.11 | -1.07 |
| C7 | 1.19 | 2.78 | 5.71 | 4.04 | 4.08 | 4.94 | 0.97 | 1.65 | 2.81 | 1.42 | 1.69 | 1.91 |
| C8 | 0.53 | 0.62 | 1.11 | 0.78 | 1.03 | 1.44 | -0.29 | -0.13 | 0.08 | 0.88 | -0.08 | -0.13 |
| C9 | 0.65 | 0.97 | 1.70 | 1.78 | 1.01 | 1.18 | -2.45 | 0.82 | -0.53 | -0.90 | -1.10 | -1.10 |
| C10 | -0.73 | -1.62 | 0.04 | -0.24 | 0.16 | 0.57 | -0.36 | -0.89 | 0.28 | -0.16 | -0.16 | -0.36 |
| C11 | -0.80 | -0.52 | 0.16 | -0.84 | -1.76 | -0.92 | 0.12 | -0.63 | 0.00 | -0.20 | -1.30 | -1.10 |
| C12 | 1.63 | 0.87 | 2.57 | 3.06 | 3.06 | 3.25 | -0.23 | 0.46 | 1.45 | 1.07 | 1.53 | 1.60 |
| C13 | -0.27 | 1.26 | 1.45 | 1.33 | 1.45 | 1.41 | 1.13 | 2.00 | 2.95 | 2.12 | 1.29 | 1.13 |
| C14 | 0.21 | 0.98 | 0.82 | 1.11 | 1.35 | 2.17 | 2.58 | 2.26 | 2.34 | 1.49 | 1.09 | 1.17 |
| C15 | 0.12 | 0.04 | 0.72 | 1.63 | 1.55 | 2.19 | 0.00 | 0.32 | 1.12 | 0.76 | 0.48 | 0.12 |
| C16 | -1.04 | -0.85 | -0.31 | -2.08 | -0.69 | -0.85 | -0.31 | -0.93 | -0.97 | -2.01 | -2.32 | -2.17 |
| C17 | 0.93 | 2.05 | 1.85 | 2.28 | 1.89 | 2.28 | 1.34 | 3.86 | 4.79 | 3.68 | 2.45 | 2.75 |
| C18 | 0.59 | 0.31 | -0.12 | 0.04 | 0.08 | -0.16 | 0.36 | 0.91 | 0.32 | 1.15 | -0.48 | 0.36 |
| C19 | 1.83 | 1.99 | 2.10 | 2.65 | 2.26 | 2.57 | 0.35 | 1.54 | 1.66 | 3.24 | 3.39 | 3.24 |
| C20 | 0.40 | 0.08 | 2.24 | 2.64 | 2.28 | 1.56 | -0.08 | -0.40 | 1.09 | 0.93 | -0.16 | -0.36 |
| Average (SD) | 0.48 (1.02) | 0.77 (1.10) | 1.47 (1.44) | 1.35 (1.41) | 1.41 (1.37) | 1.70 (1.42) | 0.26 (1.18) | 0.70 (1.30) | 1.00 (1.44) | 0.80 (1.31) | 0.20 (1.37) | 0.29 (1.36) |

Table 16 The % whitening values in the upper arms of volunteers (group D) treated with 3.00 % Kojic acid and Propylene glycol for 12 weeks.

| Volunteer No. | % Whitening (3 % Kojic acid) | | | | | | % Whitening (Propylene glycol) | | | | | |
|---------------|------------------------------|--------|--------|--------|---------|---------|--------------------------------|--------|--------|--------|---------|---------|
| | week 2 | week 4 | week 6 | week 8 | week 10 | week 12 | week 2 | week 4 | week 6 | week 8 | week 10 | week 12 |
| D1 | 0.71 | -0.08 | 0.87 | 0.33 | -0.46 | -0.29 | -0.92 | -0.83 | 0.33 | 0.29 | -0.79 | -0.46 |
| D2 | -0.89 | 0.93 | 0.81 | -0.12 | 0.08 | 0.77 | -0.08 | 0.61 | 0.28 | -0.93 | -0.69 | -0.69 |
| D3 | 0.08 | 0.51 | 1.54 | 1.73 | 2.01 | 2.72 | 0.31 | 1.06 | 2.00 | 1.92 | 1.73 | 2.16 |
| D4 | 1.78 | 1.59 | 2.37 | 3.14 | 3.88 | 3.10 | -0.04 | 1.37 | 1.29 | 0.59 | 0.51 | 0.35 |
| D5 | 0.35 | 1.39 | 2.73 | 2.00 | 2.00 | 2.70 | -0.27 | 0.39 | 0.51 | 0.86 | 0.51 | 0.70 |
| D6 | 1.50 | 2.17 | 3.71 | 3.89 | 4.04 | 4.46 | -0.15 | 1.32 | 1.32 | 2.26 | -0.45 | -0.30 |
| D7 | -0.16 | 0.12 | 1.23 | 0.37 | 1.60 | 1.76 | 1.53 | 1.61 | 1.81 | 0.64 | 1.20 | 1.73 |
| D8 | -0.08 | -0.20 | 0.60 | 0.44 | -0.56 | 0.40 | -0.44 | -0.32 | 0.48 | 0.36 | 0.08 | -0.56 |
| D9 | 2.70 | 2.13 | 2.28 | 2.74 | 3.35 | 3.65 | 1.21 | 1.09 | 0.31 | 0.23 | 0.16 | -0.12 |
| D10 | 2.36 | 1.90 | 2.47 | 3.34 | 3.42 | 3.99 | 1.00 | 2.53 | 1.99 | 1.65 | 1.99 | 2.03 |
| D11 | -0.49 | 1.54 | 1.54 | 1.77 | 2.67 | 2.67 | -0.08 | 0.23 | -0.11 | 0.76 | 0.19 | 1.28 |
| D12 | 0.76 | 0.28 | 1.40 | 1.56 | 2.08 | 2.08 | -0.32 | -1.40 | -1.00 | -0.92 | -0.92 | -1.04 |
| D13 | -1.84 | -0.33 | 0.52 | -0.66 | 1.47 | 2.28 | -1.51 | -1.36 | -0.45 | -0.87 | -0.76 | |
| D14 | 0.19 | 2.25 | 2.06 | 2.25 | 2.22 | 4.09 | 0.27 | 1.26 | 1.72 | 0.34 | 0.31 | 0.23 |
| D16 | -0.92 | -0.68 | 0.56 | 0.60 | 0.72 | 1.48 | 0.28 | -0.91 | -0.75 | -0.32 | -0.24 | 0.04 |
| D17 | -0.86 | -1.45 | 1.53 | 0.51 | 0.71 | 0.98 | -1.33 | -2.23 | -1.29 | -0.74 | -1.96 | -1.76 |
| D18 | -0.52 | -0.60 | -0.44 | 0.00 | -0.04 | 0.40 | 0.28 | 1.71 | 0.40 | -0.12 | -0.36 | -0.04 |
| D19 | -0.46 | 0.80 | 2.16 | 1.71 | 2.43 | 2.92 | 1.12 | 2.32 | 2.09 | 2.24 | 3.10 | 2.73 |
| D20 | -0.16 | 0.56 | 1.08 | 1.44 | 1.83 | 2.23 | -0.52 | -0.16 | 0.76 | 0.88 | -0.08 | 0.16 |
| Average | 0.21 | 0.68 | 1.53 | 1.42 | 1.76 | 2.23 | 0.02 | 0.44 | 0.57 | 0.50 | 0.18 | 0.30 |
| (SD) | (1.18) | (1.09) | (0.97) | (1.28) | (1.40) | (1.35) | (0.81) | (1.34) | (1.12) | (0.99) | (1.18) | (1.19) |

Appendix III

Individual % tyrosinase inhibitory activity of

0.25% Puag-Haad solution in various types of antioxidants were prepared in the same solvent (20% propylene glycol in water) and coded as follows:

P = 0.25% Puag-Haad

P+A1 = 0.25% Puag-Haad + 0.15% sodium metabisulfite

P+A2 = 0.25% Puag-Haad + 0.10% butylated hydroxy anisole (BHA)

P+A3 = 0.25% Puag-Haad + 0.05% EDTA

P+A4 = 0.25% Puag-Haad + 0.15% sodium metabisulfite + 0.10% BHA

= P + A1 + A2

P+A5 = 0.25% Puag-Haad + 0.15% sodium metabisulfite + 0.05% EDTA

= P + A1 + A3

P+A6 = 0.25% Puag-Haad + 0.10% BHA + 0.05% EDTA

= P + A2 + A3

P+A7 = 0.25% Puag-Haad + 0.15% sodium metabisulfite + 0.10% BHA + 0.05% EDTA

= P + A1 + A2 + A3

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Table 1 Stability of Puag-Haad with and without antioxidants as determined from % tyrosinase inhibitory activity at room temperature (ambient 27 °C)

Week 0

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 75.10 | 73.88 | 67.70 | 69.81 | 71.57 | 72.39 | 63.43 | 82.09 |
| 2 | 71.46 | 71.76 | 66.97 | 66.45 | 69.03 | 73.13 | 63.43 | 79.85 |
| 3 | 72.23 | 74.10 | 64.82 | 70.91 | 70.15 | 74.63 | 69.40 | 84.70 |
| Mean | 72.93 | 73.24 | 66.50 | 69.05 | 70.25 | 73.38 | 65.42 | 82.21 |
| (SD) | (1.92) | (1.29) | (1.50) | (2.32) | (1.27) | (1.14) | (3.45) | (2.43) |

Week 4

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 60.07 | 68.60 | 62.47 | 65.60 | 66.71 | 77.45 | 63.01 | 84.47 |
| 2 | 63.03 | 69.33 | 68.58 | 61.30 | 68.62 | 72.26 | 65.90 | 78.56 |
| 3 | 58.23 | 74.14 | 60.36 | 64.97 | 69.84 | 77.30 | 64.43 | 83.73 |
| Mean | 60.44 | 70.69 | 63.80 | 63.96 | 68.39 | 75.67 | 64.45 | 82.26 |
| (SD) | (2.42) | (3.01) | (4.27) | (2.32) | (1.58) | (2.96) | (1.45) | (3.22) |

Week 8

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 49.47 | 66.62 | 63.74 | 65.91 | 69.11 | 64.41 | 54.96 | 75.84 |
| 2 | 53.74 | 69.04 | 61.30 | 57.60 | 67.33 | 60.14 | 56.52 | 75.84 |
| 3 | 53.02 | 67.74 | 58.60 | 61.60 | 66.62 | 66.90 | 56.80 | 78.11 |
| Mean | 52.08 | 67.80 | 61.21 | 61.70 | 67.69 | 63.82 | 56.09 | 76.59 |
| (SD) | (2.29) | (1.21) | (2.57) | (4.15) | (1.28) | (3.42) | (0.99) | (1.31) |

Week 12

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 51.98 | 62.82 | 54.20 | 58.43 | 66.44 | 63.34 | 53.17 | 73.86 |
| 2 | 45.44 | 64.20 | 57.59 | 56.61 | 65.58 | 59.38 | 56.27 | 74.53 |
| 3 | 45.44 | 65.92 | 55.86 | 54.05 | 66.78 | 59.21 | 52.48 | 68.28 |
| Mean | 47.62 | 64.31 | 55.88 | 56.36 | 66.27 | 60.64 | 53.97 | 72.22 |
| (SD) | (3.78) | (1.55) | (1.69) | (2.20) | (0.62) | (2.34) | (2.02) | (3.43) |

Table 1 Stability of Puag-Haad with and without antioxidants as determined from % tyrosinase inhibitory activity at room temperature (ambient 27 °C)

Week 16

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 51.25 | 58.27 | 54.26 | 59.49 | 64.71 | 62.55 | 54.29 | 66.55 |
| 2 | 49.25 | 56.21 | 58.54 | 54.99 | 63.51 | 60.13 | 57.03 | 67.26 |
| Mean | 50.25 | 57.24 | 56.40 | 57.24 | 64.11 | 61.34 | 55.66 | 66.90 |
| (SD) | (1.41) | (1.45) | (3.03) | (3.18) | (0.85) | (1.71) | (1.93) | (0.50) |

Week 24

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 38.19 | 45.83 | 46.68 | 41.62 | 57.06 | 52.20 | 50.17 | 56.33 |
| 2 | 35.23 | 46.27 | 44.37 | 39.03 | 56.44 | 50.15 | 48.86 | 57.36 |
| Mean | 36.71 | 46.05 | 45.53 | 40.33 | 56.75 | 51.18 | 49.52 | 56.85 |
| (SD) | (2.09) | (0.31) | (1.63) | (1.83) | (0.44) | (1.45) | (0.93) | (0.73) |

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Table 2 Stability of Puag-Haad with and without antioxidants as determined from % tyrosinase inhibitory activity at 45 °C

Week 0

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 75.10 | 73.88 | 67.70 | 69.81 | 71.57 | 72.39 | 63.43 | 82.09 |
| 2 | 71.46 | 71.76 | 66.97 | 66.45 | 69.03 | 73.13 | 63.43 | 79.85 |
| 3 | 72.23 | 74.10 | 64.82 | 70.91 | 70.15 | 74.63 | 69.40 | 84.70 |
| Mean | 72.93 | 73.24 | 66.50 | 69.05 | 70.25 | 73.38 | 65.42 | 82.21 |
| (SD) | (1.92) | (1.29) | (1.50) | (2.32) | (1.27) | (1.14) | (3.45) | (2.43) |

Week 2

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 63.26 | 62.89 | 59.67 | 64.26 | 67.54 | 70.95 | 65.91 | 81.00 |
| 2 | 62.86 | 58.76 | 59.76 | 63.57 | 67.57 | 68.98 | 61.25 | 79.25 |
| 3 | 64.30 | 60.48 | 62.08 | 60.48 | 69.26 | 72.63 | 58.32 | 83.60 |
| Mean | 63.47 | 60.71 | 60.50 | 62.77 | 68.13 | 70.85 | 61.83 | 81.29 |
| (SD) | (0.74) | (2.07) | (1.36) | (2.01) | (0.98) | (1.83) | (3.83) | (2.19) |

Week 4

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 53.70 | 58.78 | 58.35 | 66.67 | 71.95 | 75.58 | 62.71 | 81.95 |
| 2 | 51.39 | 56.67 | 60.96 | 61.39 | 64.69 | 64.69 | 55.45 | 78.98 |
| 3 | 56.01 | 59.65 | 62.05 | 66.01 | 68.65 | 64.69 | 67.00 | 79.31 |
| Mean | 53.70 | 58.36 | 60.45 | 64.69 | 68.43 | 68.32 | 61.72 | 80.08 |
| (SD) | (2.31) | (1.53) | (1.90) | (2.88) | (3.64) | (6.29) | (5.84) | (1.63) |

Week 6

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 50.44 | 55.19 | 56.07 | 65.56 | 63.70 | 68.22 | 61.15 | 67.74 |
| 2 | 53.33 | 59.07 | 60.30 | 64.44 | 64.26 | 68.52 | 63.56 | 73.89 |
| 3 | 54.21 | 58.52 | 57.44 | 60.00 | 64.96 | 70.89 | 62.89 | 74.44 |
| Mean | 52.66 | 57.59 | 57.94 | 63.33 | 64.31 | 69.21 | 62.53 | 72.02 |
| (SD) | (1.97) | (2.10) | (2.15) | (2.94) | (0.63) | (1.46) | (1.24) | (3.72) |

Table 2 Stability of Puag-Haad with and without antioxidants as determined from % tyrosinase inhibitory activity at 45 °C

Week 8

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 51.62 | 44.78 | 50.24 | 56.98 | 57.95 | 55.06 | 61.23 | 60.56 |
| 2 | 53.55 | 48.64 | 52.17 | 55.70 | 59.87 | 61.16 | 57.44 | 59.92 |
| 3 | 52.91 | 48.64 | 47.67 | 56.98 | 58.27 | 55.06 | 59.65 | 60.56 |
| Mean | 52.69 | 47.35 | 50.03 | 56.55 | 58.69 | 57.09 | 59.44 | 60.35 |
| (SD) | (0.98) | (2.22) | (2.25) | (0.74) | (1.03) | (3.52) | (1.90) | (0.37) |

Week 10

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 55.17 | 48.38 | 42.91 | 53.45 | 55.36 | 54.98 | 49.27 | 52.45 |
| 2 | 57.09 | 47.79 | 45.40 | 49.23 | 52.87 | 52.11 | 46.97 | 56.30 |
| 3 | 55.75 | 49.34 | 47.32 | 52.87 | 55.56 | 53.07 | 50.87 | 57.47 |
| Mean | 56.00 | 48.50 | 45.21 | 51.85 | 54.60 | 53.38 | 49.04 | 55.41 |
| (SD) | (0.98) | (0.78) | (2.21) | (2.29) | (1.50) | (1.46) | (1.96) | (2.63) |

Week 12

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 44.74 | 46.76 | 45.19 | 47.35 | 56.82 | 48.99 | 43.20 | 50.14 |
| 2 | 43.18 | 51.45 | 45.41 | 43.80 | 56.38 | 48.77 | 43.20 | 51.91 |
| 3 | 40.94 | 46.53 | 45.41 | 46.59 | 55.70 | 51.01 | 43.57 | 51.15 |
| Mean | 42.95 | 48.25 | 45.34 | 45.91 | 56.30 | 49.59 | 43.33 | 51.07 |
| (SD) | (1.91) | (2.78) | (0.13) | (1.87) | (0.56) | (1.23) | (0.21) | (0.89) |

Week 16

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 44.90 | 50.32 | 52.14 | 47.27 | 53.15 | 48.55 | 47.03 | 52.07 |
| 2 | 45.87 | 51.01 | 44.58 | 50.85 | 55.70 | 52.09 | 44.26 | 49.20 |
| Mean | 45.39 | 50.67 | 48.36 | 49.06 | 54.42 | 50.32 | 45.64 | 50.63 |
| (SD) | (0.69) | (0.49) | (5.35) | (2.53) | (1.81) | (2.50) | (1.96) | (2.03) |

Table 2 Stability of Puag-Haad with and without antioxidants as determined from % tyrosinase inhibitory activity at 45 °C

Week 24

| Test | P | P+A1 | P+A2 | P+A3 | P+A4 | P+A5 | P+A6 | P+A7 |
|------|--------|--------|--------|--------|--------|--------|--------|--------|
| 1 | 28.31 | 38.24 | 35.33 | 35.61 | 42.56 | 40.85 | 36.38 | 40.29 |
| 2 | 28.00 | 36.55 | 37.75 | 36.01 | 40.83 | 42.19 | 34.92 | 43.48 |
| Mean | 28.16 | 37.40 | 36.54 | 35.81 | 41.70 | 41.52 | 35.65 | 41.89 |
| (SD) | (0.22) | (1.20) | (1.71) | (0.28) | (1.22) | (0.95) | (1.03) | (2.26) |

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

VITA

Miss Koomkhwan Pengrungrangwong was born on 26th April 1976, in Nakornpathom, Thailand. She got Bachelor of Science in Pharmacy (2nd class Honor) in 1998 from Faculty of Pharmaceutical Science, Naresuan University, Thailand. She has been working as a pharmacist in pharmacy department, Makarak Hospital since 1999.

