



References

- Ahmad, R. and Cava, M.P. Leucoxylonine and Ocoxylonine-Hexaoxygenated Aporphines from *Ocotea leucoxylon*. Heterocycles 79(1977): 927-931.
- Alfred, G., Dieter, L. and Robert, R. Composition of *Avensara aromatica*, One of the Lauraceae found on Madagascar. Planta Med. 18(1969): 66-72.
- Arbain, D., Dasman, Ibrahim, S., and Sargent, M.V. The constituents of the bark of *Litsea elliptica* (Lauraceae). Aust. J. Chem. 43(1990): 1949-1952.
- Babcock, P.P., and Segalmen, A.B. Alkaloids of *Lindera benzoin* (Lauraceae): Isolation and Identification of Laurotetanine. J. Pharm. Sci. 63(1974): 1495. (Through, C.A. 82: 13999a)
- Backer, C.A., and Bakhuizen Van Den Brink, R.C., Jr. Flora of Java. Vol I: 117-135, Netherlands: N.V.P. Noordhoff-Groningen, 1965.
- Baralle, F., Busch, A., Vernengo, M.J., and Kuck, A.M. Isolation of Ocoteine from *Nectandra saligna* and *Phoebe porfiria*. J. Nat. Prod. 35(1972): 300-302.
- _____, Schvarzberg, N., Vernengo, M., and Comin, J. Dehydroocoteine and Dihydroocoteine from *Ocotea puberula*. Experientia 28(1972): 875-876.
- _____, Schvarzberg, N., Vernengo, M., Moltrasio, G.Y., and Giacopello, D. Thalycinine from *Ocotea puberula*. Phytochemistry 12(1973): 948-949.
- Bhakuni, D.S., and Dhar, M.M. Aporphine Alkaloids of *Litsea sebifera*, *L. wightiana*, and *Actinodaphne obovata*. Phytochemistry 11(1972): 3057-3059.
- _____, and Gupta, S. Alkaloids of *Litsea wightiana*. Planta Med. 48(1983): 52-54.
- Bick, I.R.C. and Douglas, G.K. Austral. J. Chem. 18(1965): 1997
- _____, Preston, N.W., and Potier, P. The alkaloids of *Cryptocarya odorata* (Lauraceae). Bull. Soc. Chim. Fr. 12(1972): 4596. (Through, C.A. 78:94832 h)

- Bick, I.R.C., et al. Alcalodes du *Cryptocarya phyllostemon* Kostermans. Planta Med. 39 (1980): 205.
- Borthakur, N., and Rastogi, R.C. Laetanine, a new Noraporphine alkaloid from *Litsea laeta*. Phytochemistry 18(1979): 910-911.
- Bandara, B.M.R., et al. Aporphine Alkaloids from *Litsea gardneri* and *Actinodaphne speciosa*. Planta Med. 55(1989): 393.
- _____, Cortes, D., Jayasingha, U.L.B., Karunaratne, V., Sotheewaran, S., and Winnigama, G.P. Isoboldine and Legenone from *Neolitsea fuscata*. Planta Med. 43(1981): 309.
- _____, Cortes, D., Jayasingha, U.L.B., Karunaratne, V., Sotheewaran, S., and Winnigama, G.P. Aporphine Alkaloids from *Litsea gardneri* and *Actinodaphne speciosa*. Planta Med. 55(1989): 393.
- Casagrande, C., and Ferrari, G. Proaporphine and Aporphine Alkaloids IV Minor Alkaloids of *Ocotea glaziovii* Mez. Farmaco. Ed. Sci. 30(1975): 490-497. (Through, C.A. 83: 93873 p.)
- Castrelo, L., Saa, J.M., Suau, R., Villaverde, C., and Potier, P. *Litsea triflora* Alkaloids. An. Quim. Ser. C. 76(1980): 171-174. (Through, C.A. 94: 61694 s)
- Castro, O.C., Lopez, V.J., and Vergara, G.A. Aporphine Alkaloids from *Phoebe pittieri* Phytochemistry 24(1985): 203-204.
- _____, Lopez, J., and Stermitz, F.R. New Aporphine Alkaloids from *Phoebe valeriana*. J. of Nat. Prod. 49(1986): 1036.
- _____, Lopez, J., and Venkatewarlu, A. Dehydroocopodine, Dicentrinone, and other Alkaloids from *Ocotea macropoda* and *Hernandida jamaicensis*. Tetrahedron 22 (1971): 2639-2643.
- Cava, M.P., et al. *Ocotea* Alkaloids: The Characterization and Structures of four new Aporphine alkaloids Bases. Tetrahedron Lett. (1968): 2437.
- _____, Behforouz, M., and Mitchell, M.J. *Ocotea* Alkaloids: Variabiline, a novel Aminoaporphine. Tetrahedron Lett. 46(1972): 4647-4649.

- Cava, M.P., Rao, K.V., Douglas, B., and Weisbush, J.A. The Alkaloids of *Cassytha americana* (*C. filiformis* L.). J. Org. Chem. 33(1968): 2443-2446.
- Chen, I.S. Studies on the Alkaloids of Formosan Lauraceous Plants. IX. Alkaloids of *Lindera oldhamii* Hemsl. J. Chin. Chem. Soc. (Taipei) 24(1977): 41-44. (Through, C.A. 87: 35908 a)
- Chen, C.C., et al. Bioactive Principles from the roots of *Lindera magaphilla*. Planta Med. 57(1991): 406-407.
- Chen, C.H., Lee, S.S., Lai, C.F., Lu, L., and Beal, J.L. A caryachine N-methosalt from *Cryptocarya chinensis* and PMR Spectral Characteristics of Some Quaternary Pavine Alkaloids. J. Nat. Prod. 42(1979): 163-167.
- Chowdhury, B.K., Sethi, M.L., Lloyd, H.A., and Kapadia, G.J. Potential Carcinogen part 6, Aporphine and Tetrahydroisoquinoline Alkaloids in *Sassafras albidum*. Phytochemistry 15(1976): 1803-1804.
- Clezy, P.S., Gellert, E., Lau, D.Y.K., and Nichol, A.W. The alkaloids of *Eilchmiedia elliptica*. Austral. J. Chem. 19(1966): 135-142.
- Cooke, R.G., and Haynes, H.F. The Alkaloids of *Cryptocarya angulata* C.T. White and *Cryptocarya triplinervis* R.Br. Austral. J. Chem. 7(1954): 99.
- Cordell, G.A. Aporphine alkaloids in Introduction to Alkaloids A Biogenetic Approach chap. 8.18. pp 388-410. New York: John Wiley & Sons, 1981a .
- _____. Bisbenzylisoquinoline Alkaloids in Introduction to Alkaloids A Biogenetic Approach chap. 8.13. pp. 349-350. New York: John Willey & Sons, 1981b.
- _____. Morphinandienone Alkaloids in Introduction to Alkaloids A Biogenetic Approach chap. 8.24. pp. 430-431. New York: John Willey & Sons, 1981c.
- _____. Oxoaporphine in Introduction to Alkaloids A Biogenetic Approach, Chap 8.20 . pp. 416,. New York: John Willey & Sons, 1981d.
- _____. Pavine Alkaloids in Introduction to Alkaloids A Biogenetic Approach chap. 8.16. pp. 429-430. New York: John Willey & Sons, 1981e.

- Cordell, G.A. Protoberberine Alkaloids in Introduction to Alkaloids A Biogenetic Approach chap. 8.28. pp. 472-473. New York: John Wiley & Sons, 1981f.
- Correa, D.B., and Gottlieb, O.R. Duckein, an Alkaloid from *Aniba duckeri*. Phytochemistry 14(1975): 271-272.
- David, D.S.F., and Satri, S.J. Alkaloid from *Nectandra megapotamica* (Sprg.) Chodat and Assler: Inhibitory Activity of the Growth of *Crithidia fasciculata* and *Trypanosoma cruzi*. Rev. Fac. Farm. dontol. Ribeirao Preto. 17(1980): 21-24.
- De la Lande, The Alkaloid of *Cryptocarya pleuroperma*. Austral. J. Exptl. Biol. Med. Sci. 26(1948): 181.
- Desai, P.D. Chemical Investigation of Some Indian Plants Part II. Indian J. Chem. 4 (1966): 457.
- Dutta, T. Investigations on *Litsea polyantha* Isolation and Identification of Actinodaphnine. J. Indian Chem. Soc. 45(1968): 987-991. (Through, C.A. 70: 144820 s)
- Ellis, J., Gellert, E., and Summons, R.E. Alkaloids of the Genus *Cinnamomum* III Identification of Reticuline as the major bark Alkaloid of *Cinnamomum laubattii*. Austral. J. Chem. 25(1972): 1829.
- Ewing, J., Hughes, G.K., Ritchie, E., and Taylor, W.C. The Alkaloids of *Cryptocarya bowiei* (Hook.) Druce. Austral. J. Chem. 6(1953): 78.
- Ferrari, G., and Casagrande, C. Extraction of the *Octea* Alkaloid, Glaziovine. Chem. Abs. 78(1967): 108439z.
- _____, Fervidi, O., and Ferrari, M. Occurrence of Isoboldine in *Nectandra pichurim*. Phytochemistry 10(1971): 465-466.
- Filho, D.D.S., and Gilbert, B. Alkaloids of *Nectandra megapotamica*. Phytochemistry 14(1975): 821-822.
- _____, and Silvio, J. Rev. Fac. Farm. Odontol. Ribeirao preto. 17(1), 21-24, 1980. (Through, C.A. 95:200537 d)

- Fraca, N.C., et al. Benzylisoquinolines from *Ocotea* Species. Phytochemistry 14(1975): 1671-1672.
- Gaind, N., and Baveja, S.K. Investigation of *Machilus macrantha* Nees II Pharmacological Action and Chemical Constitution of Machiline. J. Am. Pharm. Assoc. 49(1960): 663-665.
- Gellert, E. The Constituents of *Cryptocarya pleurosperma* White & Francis: I Pleurospermine A New Alkaloid of the Leaves. Austral. J. Chem. 12(1959): 90.
- _____, and Summons, R.E. Alkaloids of the Genus *Cinnamomum*: Isolation Structure and Synthesis of Cinnamolaurine. Tetrahedron Lett. (1969): 5055-5058.
- _____. Alkaloids of the Genus *Cinnamomum* sp. T.G.H. 13077. Austral. J. Chem. 23 (1970): 2095-2099.
- Ghose, T.P., Grishna, S., and Schlittler, E. Constitution of Actinodaphnine. Helv. Chim. Acta 17(1934): 919.
- Gilbert, B., et al. The Aporphine and Isoquinoline dienone Alkaloids of *Ocotea glaziovii*. J. Amer. Chem. Soc. 86(1964): 694.
- Glaser, R., and Bernstein, M.A. ^1H and ^{13}C NMR Studies on the Conformation of *N*-Methyl Diastereomeromers of (+)-Glaucine Hydrofluoroacetate, an Aporphine Alkaloid Salt. J. Chem. Soc. Perkin Trans. 2. (1991): 2047-2053.
- Goodwin, T.W., and Mercer, E.T. Introduction of the Plant Biochemistry 2nd ed. pp. 528-554. Oxford:Bergamon press, 1983.
- Gottlieb, O.R. Chemosystematics of the Lauraceae. Phytochemistry 11(1972): 1537-1570.
- Govindachari, T.R., Viswanathan, N., Mohamed, P.A. Litsomentol, A New tetracyclic triterpene. Chem. Commun. (1971): 665.
- Gozler, B., Lantz, M.S., and Shamma, M. The Pavine and Isopavine Alkaloids J. Nat. Prod. 46(1983): 293-309.

- Greshoff, M. Ber. 23(1890): 3537. in Ralph, I., Bick, C., and Sinchai, W. Alkaloids of the Lauraceae. Heterocycles. 9(1978): 903-945.
- Grieson, A.J.C and Long, D.G. Flora of Bhutan vol .I, part 2 , pp. 39. Edinburgh:Royal Botanic Garden ,1984.
- Groebel, V.A., Lenoir, D., and Pernet, R. Über die Inhaltsstoffe aus *Ravensara aromatica*, einer auf Madagaskar vorkommenden Lauraceae. Planta Med. 18 (1970): 66.
- Grundon, M.F., and McGarvey, J.E.B. Alkaloids from Greenheart Part I. The Isolation of the Alkaloids and the Structure of Sepeerine. J. Chem. Soc. (1960): 2739.
- Guha, K.P. and Mukherjee, B. Bisbenzylisoquinoline Alkaloids A Review. J. Nat. Prod. 42(1979): 1-83.
- Guinadeau, H., Leboeuf, M., and Cave, A. Aporphine Alkaloids. J. Nat. Prod. 38(1975): 276-338.
- . Aporphine Alkaloids II J. Nat. Prod. 42(1979): 325-356.
- . Aporphine alkaloids III. J. Nat. Prod. 46(1983): 761-835.
- Gunatilaka, A.A.L., Sotheeswaran, S., Sriyani, B., Balasubramaniam, S. Isoboldine and Lupenone from *Neolitsea fuscata*. Planta Med. 43(1981): 309-310.
- Gupta, S., and Bhakuni, D. Alkaloids of *Litsea deccadensis*. Planta Med. 55(1989): 197.
- Hart, N.K., et al. Alkaloids of Several *Litsea* species from New Guinea. Austral. J. Chem. 22(1969): 2259-2262. (Through, C.A. 90: 92313p)
- Harrigan, G.G., Gunatilaka, A.A.L., Kingston, D.G.L., Chan, G.W., and Johnson, R.K. Isolation of Bioactive and Other Oxoaporphine Alkaloids from 2 Annonaceous Plants, *Xylopia-Aethiopica* and *Miliusa banacea* C.f. J. Nat. Prod. 57(1994): 68-73.
- Hasso, R., Alfred, P., Klaus, S., and Hilda, F. Structure Elucidation and C-13 NMR of the New Aporphine Alkaloid 3-hydroxyglaucine. Liebigs Ann. Chem. (1983): 744-752. (Through, C.A. 99: 67417 e)

- Herath, H.M.T.B., Kumar, N.S., Wimalasiri, K.M.S. Structural studies of an Arabinoxylan Isolated from *Litsea glutinosa* (Lauraceae). Carbohydr. Res. 198(1990): 343-351.
- Hoffmann, J., Luzbetak, D.J., Torrance, S.J., and Cole, J.R. Cryptopleurine, Cytotoxic Agent from *Boehmeria caudata* (Urticaceae) and *Cryptocarya laevigata* (Lauraceae). Phytochemistry 17(1978): 1448.
- Holloway, D.M., and Scheinmann, A. Co-occurrence of Aporphine and Biphenyl Constituents of *Litsea turfosa*. Phytochemistry 12(1973): 1503-1505.
- Hui, W.H., Loo, S.N., and Arthur, H.R. New Aporphine Alkaloids from *Neolitsea pulchella* J. Chem. Soc. Pak. 2(1965): 157-159. (Through, C.A. 94: 205453n)
- Hussain, S.F., Amin, A., and Shamma, M. The Alkaloids of *Machilus duthei*. J. Chem. Soc. Pak. 2(1980): 157-159. (Through, C.A. 94: 205453n)
- Hutchison, J. The Genera of Flowering Plants (Angiosperm). pp. 128. Oxford: Charendon Press, 1964.
- Ivorra, M.D., Chulia, S., Lugnier, C., and Docon, M.P. Selective Action of 2 Aporphines at Alpha-1-Adrenoreceptors and Potential-Operated Ca^{2+} Channels. European J. of Pharmacology 231(1993): 165-174.
- _____, Martinez, F., Serrano, A., and Docon, P. Different Mechanism of Relaxation Induced by Aporphine Alkaloids in Rat uterus. J. of Pharmacy and Pharmacology 45(1993): 439-443.
- Jian, W.G. et al., Chemical Constituents of the Essential Oil from the Root of *Litsea cubeba* (Lour.) Pers. Hauzhong Shifan Daxue Xuebao, Ziran Kexueban 20 (1986): 169-173. (Through C.A. 106:15731v)
- Johns, S.R., and Lamberton, J.A. New Aporphine Alkaloids from *Cassytha filiformis* L. Austral. J. Chem. 19(1966): 297-302.
- _____. Alkaloids of *Phoebe clemensii* Allen. (Family Lauraceae). Austral. J. Chem. (1967): 1277-1281.

- Johns, S.R., Lamberton, J.A., and Sioumis, A.A. Alkaloids of *Actinodaphne nitida* (Lauraceae). Austral. J. Chem. 22(1969): 2257.
- . 1-Benzyl-1, 2, 3, 4-tetrahydroisoquinoline Alkaloids from *Alseodaphne archboldiana* (Allen) Kostermans (Family Lauraceae). Austral. J. Chem. (1967): 1729-1763.
- . Cassytha alkloid II, Alkaloids of *Cassytha pubescens* R.Br. Austral. J. Chem. 19 (1966): 2331-2338.
- . Cassytha Alkaloid III, Aporphine Alkaloid from *Cassytha malantha* R.Br. Austral. J. Chem. (1966): 2339-2345.
- . Cassytha alkaloid IV, The Alkaloid of *Cassytha racemosa* Nees (Family Lauraceae). Austral. J. Chem. 20(1967): 1427-1462.
- . The alkaloids of *Neolitsea pubescens* (Lauraceae). Austral. J. Chem. (1969): 1311-1312.
- Johns, S.R., Lamberton, A.J., and Tweedale, H.J. Identification of (-) Armeapavine as the Major Leaf Alkaloid of *Cryptocarya archboldiana* (Lauraceae). Austral. J. Chem. 22(1969): 1340.
- , Lamberton, A.J., Sioumis, A.A., and Tweeddale, H.J. New Aporphine Alkaloids from *Beilschmiedia podagraria*. Austral. J. Chem. 22(1969): 1277-1281.
- , Lamberton, A.J., Sioumis, A.A., and Willing, R.I. New Alkaloids From *Cryptocarya pleurosperma* (Lauraceae): The structures of Cryptopleuridine and Cryptopleurospermine. Austral. J. Chem. 23(1970): 353.
- Jones, S.B., Jr., and Luchsinger, A.E. Plant Systematics. pp. 238-239, New York: McGraw-Hill Book Company, 1979.
- Kametanni, T., Satoh, Y., Fukumoto, K., and Pai, B.R. On the Alkaloids from *Litsea xylostea*: Occurrence of (+)-Reticuline, (+)-Isoboldine, (+)-N-norisoboldine. Indian J. Chem. 9(1971): 770.
- Kataoku, H. Alkaloids of *Neolitsea acuminatissima*. Chem. Abs. 63(1965): 12005 g.

- Kerr, K.M., Kook, A.M., and Davis, P.J. High Field and 2D-NMR Studies with The Aporphine Alkaloid Glaucine. J. of Nat. Prod. 49(1986): 576-582.
- Kiang, A.K., and Sim, K.Y. Lindcarpine, An Alkaloid from *Lindera pipericarpa* Boerl (Lauraceae). J. Chem. Soc. (1967): 282.
- Kitagawa, I., et al. Dehatrine, an antimalarial bisbenzylisoquinoline alkaloid from the Indonesian medicinal plant *Beilshmiedia madang*, isolated as a mixture of two rotational isomer. Chem. Pharm. Bull. 41(1993): 997-999.
- Kostermans, A.J.G.H., Pinkley, H.V., and Stern, N.L. New Amazonian Arrow poison, *Ocotea venenosa*. Bot. Mus. Leafl. Harvard Univ. 22(1969): 241-252.
(Through, C.A. 72: 53309 a)
- Koerniati, S. *Litsea cubeba* Pers.:Potential and Prospects . Plant Resources of South East Asia pp. 268-269 Netherlands: Pudoc Wageningen, 1989.
- Kozuka, M. et al. Alkaloids from *Lindera glauca*, *Lindera erythrocarpa*, and *Lindera citridora*. J. Nat. Prod. 17(1984): 1066.
- . The Alkaloids of *Litsea japonica*, J. Pharm. Soc. Japan 82(1962): 1567.
- , Inada, A., Konoshima, T., and Sawada, T. Aporphine Alkaloids from *Parabenzoin praecox* (Sieb & Zucc.) Nakai. Chem. Pharm. Bull. 32(1984): 5055-5058.
- , Miyazawa, S., Yokoyama, K., Odani, T., and Kubo, M. Alkaloids from *Lindera umbellata*, *L. sericea* and their Varieties. J. Nat. Prod. 48(1985): 160-161.
- , Takeuchi, M., and Sawada, T. Alkaloids from *Neolitsea aciculata*. J. Nat. Prod. 47(1984): 1062.
- , Yoshikawa, M., and Sawada, T. Alkaloids from *Lindera strychnifolia* (Sieb & Zucc.) F. Vill. (Lauraceae). J. Nat. Prod. 47(1984): 1063.
- Krane, B.D., and Shamma, M., The Isoquinolone Alkaloids. J. Nat. Prod. 45(1982): 383.

- Krishna, S., and Ghose, T.P. Actinodaphnine, an Alkaloid from *Actinodaphne hookeri* Meissn. J. Indian Chem. Soc. 9(1932): 429.
- Lajis, N.H., and Mahmud, Z. *N*-Methyl-2,3,6-trimethoxymorphinandien-7-one *N*-oxide, a minor alkaloid from *Alseodaphne perakensis*. J. Nat. Prod. 54(1991): 612-614.
- _____, Sharf, A.M., Kiew, R., Khan, M.N., and Samadi, Z. Pertanika 15(1992), 176-177. (Through, C.A. 119:113430)
- Lalbot, W.A. Forest Flora of The Bombay Presidency and Sind vol.2 pp. 393-394. Delhi : Jayyed Press, 1976.
- Lamberton, J.A., and Vashist, V.N. Alkaloids of *Litsea leefeana* and *Cryptocarya foveolata* (Lauraceae). Austral. J. Chem. 25(1972): 2737-2738.
- Leboeuf, M., Cave, A., Provost, J., and Forgacs, P. Alkaloids from *Litsea laurifolata* (Jacq.) Cardemoy, Lauraceae. Planta Med. Phytother. 13(1979): 262-267. (Through, C.A. 92: 194477 d)
- _____, Ranaivo, A., Cave, A., and Moskowitz, H.G. La velucryptine, nouvel alcaloide isoquinolique isole de *cryptocarya velutinosa*. J. Nat. Prod. 52 (1989): 516-521.
- Le Quesne, P.W., Larrahondo, J.E., and Raffauf, R.T. Antitumor Plants X Constituents of *Nectandra rigida*. J. Nat. Prod. 43(1980): 353-359.
- Lee, S.S., et al. Quaternary Alkaloids from *Litsea cubeba* and *Cryptocarya konishii*. J. Nat. Prod. 56(1993): 1971-1976.
- _____, Liu, Y.C., and Chen, C.H. Neocaryachine, A New Pavine Alkaloid from *Cryptocarya chinensis*, and NMR Spectral Properties of Related Alkaloids. J. Nat. Prod. 53(1990): 1267-1271.
- _____, Chen, C.K., Chen, I.S., and Liu, K.C.S. Additional isoquinoline alkaloids from *Litsea cubeba*. J. Chin. Chem. Soc. (Taipei). 39(1992): 453-545.
- Li, H.-L., Liu, T.Sh., Huang, Ts.-Ch., Koyama, T., and, Devol, C.E. Flora of Taiwan. vol.2. pp. 439-440. Taiwan: Epoch Publishing Co., 1976.

- Lu, S.T. Alkaloids of Formosan Lauraceous Plants XII Alkaloids of *Machilus kusanoi*, Isolation of dl-coclaurine. Yakugaku Zasshi 83(1963): 19-21. (Through, C.A. 59: 3974)
- . Alkaloids of Formosan Lauraceous Plants III Alkaloids of *Machilus pseudolongifolia*. Yakugaku Zasshi 83(1963): 214-216. (Through, C.A. 59:308c)
- . Alkaloids of Formosan Lauraceous Plants XII Alkaloids of *Cryptocarya konishii* and *Machilus acuminatissimus*. Yakugaku Zasshi 87(1967): 1278-1281a. (Through, C.A. 68: 59777 r)
- . Alkaloids of Formosan Lauraceous Plants XIII Alkaloids of *Notaphoebe konishii*. Yakugaku Zasshi 87(1967b): 1282-1284. (Through, C.A. 68: 47025 f)
- , and Chen, I.S. Structure of a new Bisbenzylisoquinoline Alkaloid, Linoldhanine. Heterocycles 4(1976): 1073-1076.
- , and Homg, C.J. Studies on the Alkaloids of formosan Lauraceous plants XXI Alkaloids of *Neolitsea daibuensis* Kamikoti. Tai-wan Yao Hsueh Tsa chih. 28(1977): 27-30. (Through, C.A. 67: 117005 e)
- , and Lin, F.M. Alkaloids of Formosan Lauraceous Plants XI Alkaloids of *Litsea cubeba* II. Yakugaku Zasshi 87(1967): 878-879. (Through, C.A. 67: 117005 e)
- , and Su, T.L. Alkaloids of Formosan Lauraceous Plants XVI Alkaloids of *Neolitsea variabilima* J. Chin. Chem. Soc.(Taipei) 20(1973): 75-81. (Through, C.A. 123625 w)
- , and Su, T.L. Alkaloids of formasan Lauraceous Plants XVII Alkaloids of *Phoebe formasana*. J. Chin. Chem. Soc.(Taipei) 20(1973): 87-93. (Through, C.A. 79: 123626 x)

- Lu, S.T., and Tsai, I.T. Two Hexahydroproaporphine Alkaloids, Lauformine and N-Methyllauformine from *Phoebe formosana*. Heterocycles. 22(1984): 1031-1033.
- _____, and Leou, S.P. Alkaloids of *Dehaasia triandra*. Phytochemistry 28(1989): 615-620.
- _____, Su, T.L., and Duh, C.Y. Studies on the Alkaloids of Formosan Lauraceous Plants XXIV Alkaloids of *Litsea kawakamii* Hayata and *Litsea akoensis* Hayata. Taiwan Yao Hsueh Tsa Chih. 31(1979): 23-27. (Through, C.A. 92:194484 d)
- _____, Su, T.L. and Wang, F.C. Studies on the Alkaloids of Formosan Lauraceous Plants XVIII. Alkaloids of *Neolitsea buisanensis* Yamamoto & Kamikoti and *Neolitsea aurata* (Hay) Koidz. J. Chin. Chem. Soc. (Taipei). 22(1975): 349-353. (Through, C.A. 84: 132659 r)
- _____, Wang, S.J., and Lin, F.S. Alkaloids of Formosan Lauraceous Plants XIV Alkaloids of *Actinodaphne acutivena* and *Litsea hayatae*. Yakugaku Zasshi. 89(1969): 1313-1316. (Through, C.A. 72: 6203 b)
- _____, Wang, S.J., Lai, P.H., Lin, C.M., and Lin, L.C. Alkaloids of Formosan Lauraceous Plants XV Alkaloids of *Lindera oldhamii* I. Yakugaku Zasshi. 92 (1972): 910-917. (Through, C.A. 77:101949 n)
- Manandhar, M.D., Shoeb, A., Kapil, R.S., and Popli, S.P. Studies in Medicinal Plants : Part V Bis (2-n-propylpentyl)phthalate and Reticuline from *Cryptocarya amygdalina* Nees. Indian J. Chem. 17(1979): 411-412.(Through, C.A.92: 169113z)
- Manske, R. Phenanthroindolizidine and Phenanthroquinolizidine Alkaloids in The Alkaloids vol.19. pp. 193-219. New York: Academic Press., 1981.
- Martinez, N., de Diaz Aura, M.P., and Joseph-Nathan, P. Oxoaporphine Alkaloids from the Wood of *Phoebe cinnamomifolia*. Planta Med. 54(1988): 361-362.

- Merchant, J.R., and Desai, H.K. Isolation of Nantenine from *Cassytha filiformis* Linn. & Its Synthesis. Indian J. Chem. 11(1973): 342. (Through, C.A. 79: 32149 e)
- Mors, W.B., and Gottlieb, O.R. Anibine, the Alkaloid of Coto Bark. Anais. Assoc. Brasil. Quim. 18(1959): 185. (Through, C.A. 54: 12181 h)
- _____, Gottlieb, O.R., and Djerassi, C. The Chemistry of Rosewood Isolation and Structure of Anibine and 4-Methoxyparacoteine. J. Am. Chem. Soc. 79(1957) : 4507.
- Nakasato, T. and Asada, S. Alkaloids of Lauraceous Plants VI Structure of Litsericine. Yakugaku Zasshi. 86(1966): 134-139.
- Nakasato, T., and Nomura, S. Alkaloids of Lauraceous Plants I Alkaloids of *Neolitsea sericea*. Yakugaku Zasshi. 77(1957): 816-817. (Through, C.A. 51:15893)
- _____. Alkaloids of Lauraceous Plants II Alkaloids of *Neolitsea sericea*. Yakugaku Zasshi 78(1958): 540-542. (Through, C.A. 52:17312 g)
- _____. Alkaloids of Lauraceous Plants III Alkaloids isolated from the leaves of *Neolitsea sericea*. Yakugaku Zasshi 79(1959): 1267-1272. (Through, C.A. 54:4646 f)
- Nakasato, T., Asada, S., and Koezuka,Y. Alkaloids of Lauraceous Plants V Alkaloids Isolated from the Trunk Bark of *Neolitsea sericea*. Yakugaku Zasshi 86(1966) 129-134. (Through, C.A. 54: 4646 f)
- Naruto, S. and Kaneko, H. Constituents of *Corydis stewartii* . Phytochemistry 11(1972): 2644.
- Paulo, M. de Q., et al. Antimicrobial activity of benzylisoquinoline alkaloids from *Annona salzmanii* D.C. J. Ethnopharmacol. 36(1992): 39-41. (Through, C.A. 116: 2319u)
- Patnaik, P.C., and Gopinath, K.W. *Litsea* Alkaloid: Characterization & Structure of a New Aporphine alkaloid. Indian J. chem. 13(1975): 197. (Through, C.A. 83:28394 u)

- Pech, B. and Bruneton, J. Alcaloides De L' *Umbellaria californica*. J.Nat. Prod. 47 (2), 390, 1984.
- Perry, L.M. and Metzger, J. Medicinal plants of East and Southeast Asia. pp. 200-201. Massachusetts: The MIT Press, 1980.
- Ralph, I., Bick, C., and Sinchai, W. Alkaloids of Lauraceae. Heterocycles 9(1978): 903-945.
- Rastogi, R.C., and Borthakur, N. Alkaloids of *Litsea laeta* and *L. salicifolia*. Phytochemistry 19(1980): 998-999.
- Robison, T. The Biochemistry of Alkaloids. (3 vol.) 2nd ed. pp. 76-91. New York: Springer Verlag Inc., 1981.
- Russel, G.R., and Fraser, J.G. Identification of Isoboldine as the Major alkaloid from Berries of *Beilschmiedia tawa*. N.Z.J. Sci. 12(1969): 694-695. (Through, C.A. 72: 51811 j)
- Said, A. M., Latiff, A., Partridge, S.J., and Phillipson, J.D. Alkaloids from *Dehaasia incrassata*. Planta Med. 57(1991): 389.
- Segelman, A.B., Segelman, F.P. and Karliner, J. Chemical constituents of *Sassafras albidum* (Nuttall) Nees. II Isolation and Identification of boldine, reticuline and two new alkaloids, sassafrasine and sassafrasinine from the root bark. Lloydia 39(1976): 473.
- Shamma, M. Proven or Probable Biogenetic Loci for the Formation of the Isoquinoline Alkaloids. in The Isoquinoline Alkaloids, the inner front cover. New York Academic Press, 1972 .
- . The Aporphine in The Isoquinoline Alkaloids. pp.195-225. Academic Press, New York, 1972 .
- , and Guinadeau, H. Aporphinoid Alkaloids. Nat. Prod. Rep. 2(1985): 227-233.
- , and Guinadeau, H.. Biogenetic pathways for the Aporphinoid Alkaloids. Tetrahedron. 40(1984): 4795-4822.

- Siphar, S.A. Antiasthenic Composition from an Alkaloids Extracted from a Plant of the Ocotea Family. Chem. Abs. 75(1971): 40412 r.
- Sivakumaran, M. and Gopinath, K.W. Sebiferine & Litseferine, Two New Alkaloids from *Litsea sebifera*. Indian J. Chem. 14B(1976): 150-151.
- Smitinand, T. Thai Plant Names (Botanical Names-Vernacular Names). p.26. Bangkok: Funny Publishing, 1980.
- Staunton, J. Biosynthesis of Isoquinoline Alkaloids. Planta Med. 36(1979): 1-19.
- Stermitz, F.R., and Castro, C.O. Pentasubstituted Aporphine Alkaloids from *Phoebe molicella*. J. Nat. Prod. 46(1983): 913-916.
- Suarez, C. and Enrique, L. Hydroxytobain and Reticuline from *Ocotea caparrapi*. Rev. atinoam. Quim. 11(1980):110-111. (Through, C.A. 94: 80222 c)
- Suvatti, C. Flora of Thailand. pp. 564-565. Thailand : Royal Institute, 1978.
- Talapatra, B.M., Goswami, S.P. Ghosh, A., and Talapatra, S.K. Alkaloids of *Machilus glaucescens* Wight. "Machigline", a New Phenolic Oxoaporphine Alkaloid. J. Indian. Chem. Soc. 59(1982): 1364-1368. (Through, C.A. 99: 136832 z)
- Tanaka, H., Nakamura, T., Ichino, K., Ito, K., Tanaka, T. Butanolides from *Litsea japonica*. Phytochemistry 29(1990): 857-859.
- Tantisewie, B., and Ruchirawat, S. Alkaloids from the plants of Thailand in Brossi, A., and Cordell, G.A.(eds.) The Alkaloids (vol. 41) pp 1-36. San Diego: Academic Press, Inc., 1992.
- Teixeira, M.I., and Garbarino, J.A. Alcaloides De *L'Umbellolaria californica*. J. Nat. Prod. 47(1984): 390
- Tewari, S., Bhakuni, D.S., and Dhar, M.M. The Aporphine Alkaloids of *Litsea glutinosa*. Phytochemistry 11(1972): 1149-1152. (Through, C.A. 76:138158 h)
- Tillequin, F., and Koch, M. Two New Morphinane Alkaloids from *Beilsenia oreophila* Schlechter (Lauraceae). Heterocycles 23(1985): 1357-1361.
- Tomita, M., and Ozuka, M. The alkaloids of *Cinnamomum camphora*. J. Pharm. Soc. Jap. 84(1964): 365. a

- Tomita, M., and Ozuka, M. Alkaloids of *Machilus thunbergii*. Yakugaku Zasshi 84 (1964): 362-365. (Through, C.A., 61: 4706) b
- Tomita, M., Lu, S.T., Fu, S.C., and Lin, Y.M. Studies on the Alkaloids of Formosan Lauraceous Plants VI Alkakloids of *Neolitsea acuminatissima*. J. Pharm. Soc. Jap. 85(1965): 662. (Through, C.A., 63: 12005 g) a
- _____, Lu, S.T., and Lan, P.K. Alkaloids of Formosan Lauraceous Plants V Alkakloids of *Machilus* genus Plants. J. Pharm. Soc. Jap. 85(1965): 588-593. (Through, C.A. 63: 12005 e) b
- _____, Kozuka, M., Nakagawa, E., and Mitsunori, Y. On the Alkaloids of *Laurus nobilis*. J. Pharm. Soc. Jap. 83(1963): 763.
- _____, Lu, S.T., Fu, S.C., and Lin, Y.M. Alkaloids of Formosan Lauraceous Plants V Alkakloids of *Machilus obovatifolia* and *M. arisanensis*. Yakugaku Zasshi 85 (1965): 588. (Through, C.A. 62: 12005 g) c
- _____, Lu, S.T., Lan, P.K. and Lin, F.M. Studies on the Alkaloids of Formosan Lauraceous Plants V. Alkakloids of *Litsea cubeba*. J. Pharm. Soc. Jap. 85 (1965): 593. d
- _____, Lu, S.T., and Wang, S.J. Studies on the Alkaloid Formosan Lauraceous Plants VII Alkakloids of *Cassytha filiformis* Structures of a New Aporphine-type Alkaloid "Cassyfiline". Yakugaku Zasshi 85(1965): 827-851. (Through, C.A. 63: 18187 g) e
- _____, Sawada, T., Kozuka, M., Hamano, D., and Yoshimura, K. Alkaloids of *Lindera strychnifolia* and *L. umbellata*. Yakugaku Zasshi 89(1969): 737-740. (Through, C.A. 71: 67953 x)
- _____, Yang, T.H., Gaind, K.N., and Bajeja, S.K. Structure of Machiline, An Alkaloid of *Machilus macrantha*. Yakugaku Zasshi 83(1963): 218. (Through, C.A. 59: 2874)
- _____, Yang, S.T., and Lu, S.T. Alkaloids of Formosan Lauraceous Plants I Alkakloids of *Machilus kusanoi*. Isolation of L-(-)-N-norarmepavine.

- Yakugaku Zasshi 83(1963):15-18. (Through, C.A. 59: 3974)
- Ubeda, A., Montesinos, C., Paya, M., Terencio, C., and Alcaraz, M.J. Antioxidant Action of Benzylisoquinoline Alkaloids. Free Radical Research Communications 18(1993): 167-175.
- Upadhyay, H., Bhakuni, D.S., and Dhar, M.M. Aporphine Alkaloids of *L. sebifera*, *L. Wightiana* and *Actinodaphne obovata*. Phytochemistry 1972(11): 3357.
- _____, Urzua, A., Torres, R., and Cassels, B. (+)-Reticuline in *Cryptocarya alba*. Rev. Latinoam. Quim. 6(1975): 102. (Through, C.A. 83: 111149 p)
- Vecchietti, V., Casagrande, C., and Ferrari, G. Ocobotrine and 14-Episinomenine New Trans-Morphinan Alkaloids of *Ocotea brachybotra*. Tetrahedron Lett. 19 (1976): 1631-1634.
- _____, Casagrande and Severini Ricca, G. New aporphine alkaloids of *Ocotea minarum*. Farmaco. ed. Sci. 34(1979): 829-840. (Through, C.A. 92: 19789 y)
- _____, Casagrande, C., Ferrari, G., Danieli, B., and Palmisanno, B. Alkaloids of *Ocotea acutangula*. J. Chem. Soc. Per. Trans. I 2(1981): 378-581.
- Vilegas, J.H.Y., Gottlieb, O.R., Kaplan, M.A.C., and Gottlieb, H.G. Aporphine alkaloids from *Ocotea caesia*. Phytochemistry 28(1989): 3577-3578.
- wang, J.P., Meng, S.J., and Li, J.M. Fatty acid composition of three species of *Litsea*. Chih Wu Hsueh Pao 25(1983): 245-249.
- Weber, J.F., Brunneton, J., and Pusset, J. Plants of New Caledonia Part 103. Alkaloids of *Litsea lecardii*. Planta Med. (1986): 74-75.
- Wu, J., Beal, J.L., Wu, W.N., and Doskotch, R.W. Alkaloids of Thalictrum XXIV Isolation and Identification of Alkaloids from The Tertiary Phenolic Fraction of *Thalictrum revolutum* Tops. J. Nat. Prod. 40(1977): 593-601.
- Wu, Y.C., Liou, J.Y., Duh, C.Y., Lee, S.S., and Lu, S.T. Litebamine, A Novel Phenanthrene Alkaloid from *Litsea cubeba*. Tetrahedron Lett. 32(1991): 4169-4170.

- Yu, S.M., et al. Hemodynamic-Effects of Dicentrine, a Novel Alpha-1-Adrenoreceptor Antagonist-Comparison with Prazosin in Spontaneously Hypertensive and Normotensive Wistar-Kyoto Rats. British J. of Pharmacology 106(1992): 797-801.
- Zhang, F., Lui, M., Li, Y., Mai, L., and Lu, R. Zhiwu Xuebao 30(1988): 183-186.
(Through, C.A. 109:107755 w)
- Zytshi, N.L. Essential Oils-Isolates and Semisynthesis. In Atal, C.K., and Kapur, B.M. (eds.) Cultivation and Utilization of Aromatic Plants. pp. 67. New Delhi: Publications & Information Directorate (CSIR), 1989.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย



APPENDIX

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย




Figure 3 The eims spectrum of compound Z-35




Figure 4 The uv spectrum of compound Z-35




Figure 5 The ir spectrum of compound Z-35




Figure 6 The ^1H NMR spectrum of compound Z-35




Figure 7 The ^1H nmr spectrum of compound Z-35 expanded from δ 2.15-3.25 ppm.




Figure 8 The 125 MHz ^{13}C nmr spectrum of compound Z-35





Figure 9 The ^1H , ^1H COSY spectrum of compound Z-35







Figure 10 The ^1H , ^{13}C COSY spectrum of compound Z-35






Figure 11 The ^1H , ^{13}C NMR spectrum of compound Z-35




Figure 12: The eims spectrum of compound G-47




Figure 13 The uv spectrum of compound G-47




Figure 14 The ir spectrum of compound G-47




Figure 15 The ^1H NMR spectrum of compound G-47





Figure 16 The 125 MHz ^{13}C nmr spectrum of compound G-47






Figure 17 The ^1H , ^1H COSY spectrum of compound G-47





Figure 18 The ¹H, ¹³C COSY spectrum of compound G-47




Figure 19 The ¹H, ¹³C COLOC spectrum of compound G-47





Figure 20 The ^1H NMR spectrum of compound G-47 expanded from δ 2.3–3.3 ppm







Figure 21 The ^1H decoupling experiment of compound G-47



Figure 22 The HMBC spectrum of compound G-47 expanded from δ 2.4-4.0 ppm





Solvent system :- chloroform:hexane: methanol (3:1:0.3)
Figure 23 TLC chromatogram of compound Z-35 and G-47



VITA

NAME: Boonying Tubtim
BORN: August 22, 1963
EDUCATION: B.S. in Pharmacy, Faculty of Pharmaceutical Sciences, Chieng Mai University, Chieng Mai, Thailand, 1987
WORKING: Mae Tha Hospital, Lum Poon, Thailand

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย