

ເອກສາຮອ້າງອີງ

- Allan, G., Brook, C.D., Cambridge, D., and Haldkiwskyj, J.J. (1983). Enhanced responsiveness of vascular smooth muscle to vasoconstrictor agents after removal of endothelial cells. Br. J. Pharmacol. 79 : 334 P.
- Alosachie, I., and Godfraind, T. (1986). Role of cyclic GMP in the modulation by endothelium of the adrenolytic action of prazosin in the rat isolated aorta. Br. J. Pharmacol. 89 : 525-532.
- Angus, J.A., Cocks, T.M., and Sato, K. (1986a). Alpha adrenoceptors on endothelial cells. Fed. Proc. 45 : 2355-2359.
- _____. Cocks, T.M., and Sato, K. (1986b). Alpha-1 adrenoceptors and endothelium - dependent in canine large arteries. Br. J. Pharmacol. 88 : 767-777.
- _____. Cocks, T.M., and Sato, K. (1986c). Depression of contractile response in rat aorta by spontaneously released endothelium-derived relaxing factor. J. Pharmacol. Exp. Ther. 237 : 529-538.

Apperley, E., Feniuk, W., Humphrey, P.P.A., and Levy, G.P. (1980). Evidence for two types of excitatory receptor for 5-hydroxytryptamine in dog isolated vasculature. Br. J. Pharmacol. 68 : 215-224.

Barnden, I.M., and Colson, R., rev. eds. (1980). Experiments on isolated smooth muscle preparations. Hugo Sachs Electronik KG : Germany.

_____. and Lang, W.J. (1973). Neural dopaminergic vasodilator control in the kidney. Nature 246 : 27-29.

_____. and Mya, M.K.K. (1977). Is the vasodilatation induced by β -adrenoceptor stimulants in the dog mediated through dopamine receptor. Experimentia 33 : 638-639.

Bell, C. and Stubbs, A. (1978). Localization of vasodilator dopamine receptors in the canine hindlimb. Br. J. Pharmacol. 64 : 253-257.

Berkowitz, B.A. (1983). Dopamine and dopamine receptors as target site for cardiovascular drug action. Fed. Proc. 42 : 3019-3021.

Berridge, M.J. (1987). Inositol triphosphate and diacylglycerol : two interacing second messengers. Annu. Rev. Biochem. 56 : 159-193.

Bevan. J.A., Bevan, R.D., and Duckles, S.P. (1980).

Adrenergic regulation of vascular smooth muscle.

In D.F. Bohr, A.P. Somlyo, and H.V. Spark Jr.

(eds.), Handbook of Physiology : The CVS, pp. 515-

566. Baltimore : American Physiological Society.

Blackshear, J.L., Orlandi, C., Garnic, J.D., and Hollenberg, N.L. (1985). Differential large and small vessel

responses to serotonin in the dog hindlimb in vivo : role of the 5-HT₂-receptor. J. Cardiovasc. Pharmacol.

7 : 42-49.

Bolton, T.B. (1979a). Cholinergic mechanisms in smooth

muscle. Br. Med. Bull. 35 : 275-283.

_____. (1979b). Mechanisms of action of transmitters

and other substance on smooth muscle. Physiol. Rev.

59 : 606-718.

Brading, A.F. (1981.) How do drugs initiate contraction

in smooth muscles? TIPS : 261-265.

_____. Burdyga, T.V. and Scripnyuk, Z.D. (1983).

The effects of papaverine on the electrical and mechanical activity of the guinea-pig ureter.

J. Physiol. 334 : 79-89.

Bradley, P.B., Humphrey, P.P.A., and Williams, R.H. (1986).

Evidence of 5-hydroxytryptamine receptors, which are not of the 5-HT₂ type mediating contraction of isolated rabbit basilar artery. Br. J. Pharmacol.

87 : 3-4.

Brayden, J.E., and Bevan, J.A. (1985). Neurogenic muscarinic vasodilation in the cat : an example of endothelial cell-independent cholinergic relaxation. Circ. Res. 56 : 205-211.

Brenner, B.M., Troy, J.L., and Ballermann, B.J. (1989).

Endothelium-dependent vascular responses : mediator and mechanisms. J. Clin. Invest. 84 : 1373-1378.

Brodde, O.E. (1982). Vascular dopamine receptors : demonstration and characterization by in vitro studies. Life Sci. 31 : 289-306.

_____. and Meyer, F.J. (1979). On the existence of vascular dopamine receptors on the isolated rabbit mesenteric artery. Naunyn-Schmiedeberg's Arch. Pharmacol. 308 : R 56.

_____. Meyer, F.J., Schemuth, W., and Freistuhler (1981). Demonstration of specific vascular dopamine receptors mediating vasodilation on the isolated rabbit mesenteric artery. Naunyn-Schmiedeberg's Arch. Pharmacol. 316 : 24-30.

_____. and Schemuth, W. (1979). Specific antagonism by metoclopramide of dopamine-induced relaxation on isolated rabbit mesenteric arteries contracted with prostaglandin F₂. Life Sci. 25 : 23-30.

Bullock, G.R., Taylor, S.G., and Weston, A.H. (1986). Influence of the vascular endothelium on agonist-induced contractions and relaxations in rat aorta. Br. J. Pharmacol. 89 : 819-830.

Burgen, A.S.V., and Mitchell, J.F. (1985). Circulation. In Gaddum's Pharmacology, pp. 118-129. 9th ed. New York : Oxford University Press.

Burnstock, G. (1980). Cholinergic and purinergic regulation of blood vessels. In D.F. Bohr, A.P. Somlyo, H.V. Sparks Jr. (eds.), Handbook of Physiology : the CVS, pp. 567-596. Baltimore : American Physiological Society.

Carrier, G.O., and White, R.E. (1985). Enhancement of alpha-1 and alpha-2 adrenergic agonist-induced vasoconstriction by removal of endothelium in rat aorta. J. Pharmacol. Exp. Ther. 232 : 682-687.

Casteels, T., and Suzuki, H. (1980). The effect of histamine on the smooth muscle cells of the ear artery of the rabbit. Pflugers. Arch. 387 : 17-25.

Calvete, J.A., Hayes, R.J., Oates, N.S., Sever, P.S., and Thom, S. (1984). Alpha-1 and alpha-2 adrenoceptor response in human isolated arteries. Br. J. Pharmacol. 83 : 364P.

Cherry, P.D., Furchtgott, R.F., Zawadzki, J.D., and Jothianandan, D. (1982). The role of endothelial cells in the relaxation of isolated arteries by bradykinin. Proc. Natl. Acad. Sci., U.S.A. 79 : 2105-2110.

Clark, B.J. (1985). The role of dopamine in periphery. In E. Fluckiger, E.E. Müller, and M.O. Thorner (eds.), The Dopaminergic System, pp. 27-39. Berlin : Springer-Verlag.

_____. and Meninger, K (1980). Peripheral dopamine receptor. Cir. Res. 46 : 59-63.

Cocks, T.M., Angus, J.A. (1983). Endothelium-dependent relaxation of coronary arteries by noradrenaline and serotonin. Nature 305 : 627-630.

Cohen, M.L., and Berkowitz, B.A. (1975). Differences between the effects of dopamine and apomorphine on rat aortic strips. Eur. J. Pharmacol. 34 : 49-58.

- _____. Fuller, R.W., and Wiley, K.S. (1981). Evidence for 5-HT₂ receptor mediating contraction in vascular smooth muscle. J. Pharmacol. Exp. Ther. 218 : 421-425.
- _____. Mason, N., Wiley, K.S., and Fuller, R.W. (1983a). Further evidence that vascular serotonin receptors are of the 5-HT₂ type. Biochem. Pharmacol. 32 : 567-570.
- Cohen, M.L., and Wittenauer, L.A. (1987). Serotonin receptor activation of phosphoinositide turnover in uterine, fundal, vascular, and tracheal smooth muscle. J. Cardiovascc. Pharmacol. 10 : 176-181.
- Cohen, R.A. (1986). Contractions of isolated canine coronary arteries resistant to S₂-serotonergic blockade. J. Pharmacol. Exp. Ther. 237 : 548-552.
- _____. (1989). Interactions of 5-hydroxytryptamine with endothelial cells. In J.R. Fozard (ed.), The peripheral action of 5-HT, pp. 182-200. New York : Oxford University Press.
- _____. Shepherd, J.T., and Vanhoutte, P.M. (1983). 5-Hydroxytryptamine can mediate endothelium-dependent relaxation of coronary arteries. Am. J. Physiol. 245 : H1077-H1080.

Zitnay, K.M., Weisbrod, R.M., and Tesfmariam, B. (1988). Influence of the endothelium on tone and the response of isolated pig coronary artery to norepinephrine J. Pharmacol. Exp. Ther. 244 : 550-555.

Connor, H.E., and Feniuk, W. (1988). Endothelium removal augments reactivity of canine basilar artery to 5-HT and selective 5-HT agonist. Br. J. Pharmacol. 93 : 267P.

Connor, H.E., Feniuk, W., and Humphrey, P.P.A. (1987). Characterisation of 5-HT receptor mediating contraction in dog and primate isolated basilar artery. Br. J. Pharmacol. 92 : 757P.

Conti, A., Monopoli, A., Forlani, A., Ongini, E., Antona, C., and Bigliolo, P. (1990). Role of 5-HT₂ receptor in serotonin-induced contraction in the human mammary artery. Eur. J. Pharmacol. 176 : 207-212.

Coulson, C.J. (1988). Neurotransmitter action and metabolism. In Molecular of drug action, pp. 127-186. London : Taylor & Francis.

Davies, J.M., and Williams, K.I. (1983). The involvement of the endothelium in the action of substances which relax the rat aorta. Br. J. Pharmacol. 80 : 445P.



DeMey, J.G., and Vanhoutte, P.M. (1981). Role of the intima in cholinergic and purinergic relaxation of isolated canine femoral arteries. J. Physiol. 316 : 347-355.

Dinerstein, R.J., Jones, R.T., and Goldberg, L.I. (1983). Evidence for dopamine-containing renal nerves. Fed. Proc. 42 : 3005-3008.

Duckles, S.P. (1988). Vascular muscarinic receptors : pharmacological characterization in the bovine coronary artery. J. Pharmacol. Exp. Ther. 246 : 929-934.

Edvinsson, L., Hardebo, J.E., and Owman, C. (1978). Pharmacological analysis of 5-hydroxytryptamine receptors in isolated intracranial and extracranial vessels of cat and man. Cir. Res. 35 : 835-849.

_____. and Owman, C.A. (1975). A pharmacological comparison of histamine receptors in isolated extracranial and intracranial arteries in vitro. Neurology 25 : 271-276.

Egleme, C., Godfraind, T., and Miller, R.C. (1984a). Enhanced responsiveness of rat isolated aorta to clonidine after removal of the endothelial cells. Br. J. Pharmacol. 81 : 16-18.

Godfraind, T., and Miller, R.C. (1984b). Role of endothelium in the rat aorta response to vasoconstrictors. Arch. int. Pharmacodyn. Ther. 268 : 116.

Ercan, A.S., and Turker, R.K. (1975). Histamine receptors in the vasculature of the rabbit ear. Naunyn-Schmiedeberg's Arch. Pharmacol. 291 : 23-29.

Feniuk, W., Humphrey, P.P.A., and Watts, A.D. (1983). 5-Hydroxytryptamine-induced relaxation of isolated mammalian smooth muscle. Eur. J. Pharmacol. 96 : 71-78.

Ferguson, D.R., Johnson, B.I., and Price, N. (1985). Responses of the pig isolated renal artery to transmural electrical stimulation and drugs. Br. J. Pharmacol. 84 : 879-882.

Forster, C., Drew, G.M., Hilditch, A., and Whalley, E.T. (1983). Dopamine receptors in human basilar arteries. Eur. J. Pharmacol. 87 : 227-235.

Forstermann, U., Trogisch, G., and Busse, R. (1984). Species-dependent differences in the nature of endothelium-derived vascular relaxing factor. Eur. J. Pharmacol. 106 : 639-643.

Furchtgott, R.F., (1955). The pharmacology of vascular smooth muscle. Pharmacol. Rev. 7 : 183-205

- _____. (1981). The requirement for endothelial cells in the relaxation of artery by acetylcholine and some other vasodilators. TIPS 2: 173-176.
- _____. (1983). Role of endothelium in responses of vascular smooth muscle. Cir. Res. 53 : 557-573.
- _____. (1984). Role of endothelium in response of vascular smooth muscle to drugs. Annu. Rev. Pharmacol. Toxicol. 24 : 175-197.
- _____. Cherry, P.D., Zawadki, J.V.L., and Jothianandan, D. (1984). Endothelial cells as mediators of vasodilation of arteries. J. Cardiovasc. Pharmacol. 6 : S336-S344.
- _____. Jothianandan, D., Martin, W., and Villani, G.M. (1985). Depression of contractile response by spontaneously released EDRF in rat aorta. Br. J. Pharmacol. 86 : 565P.
- _____. and Zawadzki, J.U. (1980). The obligatory role of endothelial cells in relaxation smooth muscle by acetylcholine. Nature 288 : 373-376.
- _____. and Vanhoutte, P.M. (1989). Endothelium derived relaxing and contracting factors. FASEB J. 3 : 2007-2018.

Georgieva, S.A., ed. (1989). Essentials of physiology.
Moscow : Mir Publishers.

Godfraind, T., Egleme, C., and Alosachie, I. (1985).

Role of endothelium in the contractile response of rat aorta to alpha-adrenoceptor agonists. Clin. Sci. 68 : 65s-71s.

_____. Miller, R., and Wibo, M. (1986). Calcium antagonism and calcium entry blockers. Pharmacol. Rev. 38 : 321-416.

Goldberg, L.I. (1972). Cardiovascular and renal actions of dopamine : potential clinical applications. Pharmacol. Rev. 24 : 1-29.

_____. Kohli, J.D., Kotake, A.N., and Volkman, P.H. (1978). Characteristics of the vascular dopamine receptor : comparison with other receptor Fed. Proc. 37 : 2396-2403.

_____. and Toda, N. (1975). Dopamine-induced relaxation of isolated canine renal, mesenteric and femoral arteries contracted with prostaglandin F₂. Circ. Res. 36 & 37 : 197-1102.

Gorman, P. (1988). Vasodilator response to dopamine in the ferret pulmonary circulation. Br. J. Pharmacol. 94 : 212-218.

Gothert, M., Theis, F.K., and Weth, N. (1976). Effect of droperidol on cardiovascular adrenoceptors. Arch int. Pharmacodyn. Ther. 224 : 199-214.

_____. and Schlicker, E. (1987). Classification of serotonin receptors. J. Cardiovasc. Pharmacol. 10 : S3-S7.

Gotoh, S., Ogihara, T., Ohde, H., Nakamura, M., Masvo, K., Saeki, S., and Kumahara, Y. (1982). Release of prostaglandin I₂ from dog kidney by propranolol. Prostagl. Leuk. Med. 8 : 537-544.

Goyal, R.K. (1989). Muscarinic receptor subtypes : physiology and clinical implications. N. Eng. J. Med. 321 : 1022-1029.

Grega, G.J., and Adamski, S.W. (1987). Patterns of constriction produced by vasoactive agents. Fed. Proc. 46 : 270-275.

Griffith, T.M., Edwards, D.H., Lewis, M.J., Newby, A.C., and Henderson, A.H. (1984a). The nature of endothelium-derived vascular relaxant factor. Nature 308 : 645-647.

_____. Henderson, D., Edwards, D.H., and Lewis, M. J. (1984b). Isolated perfused rabbit coronary artery and aortic strip preparations : the role of endothelium-derived relaxant factor. J. Physiol. 351 : 13-24.

Haeusler G, and Osterwalder R. (1984). Different response of pulmonary arterial smooth muscle to alpha-1 and alpha-2 adrenoceptor agonists. Naunyn-Schmiedberg's Arch. Pharmacol. 325 : R64.

Heltianu, C., Simionescu, M., and Simionesco, N. (1984). Detection of histamine at cellular level. Biochem. Pharmacol. 33 : 343-346.

Hickey, K.A., Rubanyi, G., Paul, R.J., and Highsmith, R. F. (1985). Characterization of a coronary vasoconstrictor produced by cultured endothelial cells. Am. J. Physiol. 248 : C550-C556.

Hieble, J.P., and Woodward, D.F. (1984). Different characteristics of postjunctional alpha-adrenoceptors on arterial and venous smooth muscle. Naunyn-Schmiedeberg's Arch. Pharmacol. 328 : 44-50.

Hilditch, A., and Drew, G.M. (1981). Characteristics of the dopamine receptors in the rabbit isolated splenic artery. Eur. J. Pharmacol. 72 : 287-296.

Horn, P.T., Kohli, J.D., Listinsky, J.J., and Goldberg, L.I. (1982). Regional variation in the alpha-adrenergic receptor in the canine resistance vessels. Naunyn-Schmiedeberg's Arch. Pharmacol. 319 : 1-7.

_____. Marcou, M., Munday, K.A., Woodruff, G.N. (1981). Effects of dopamine receptor agonists in the guinea-pig renal vasculature and their antagonism by sulpiride. Br. J. Pharmacol. 72 : 507P-508P.

Hoshino, Y., Obara, H., and Iwai, S. (1986) Relaxant effects of dopamine on isolated rabbit pulmonary artery. Life Sci. 39 : 2525-2531.

Hynes, M.R., Banner, W. Jr., Yamamura, H.T., and Duckles, S.P. (1986). Characterization of muscarinic receptor of the rabbit ear artery smooth muscle and endothelium. J. Pharmacol. Exp. Ther. 238 : 101-105.

Ignarro, L.J., Buga, G.M., Wood, K.S., Byrns, R.E., and Chaudhuri, G. (1987). Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. Proc. Natl. Acad. Sci., U.S.A. 84 : 9215-9269.

_____. and Kadowitz, P.J. (1985). The pharmacological and physiological role of cyclic GMP in vascular smooth muscle relaxation. Annu. Rev. Pharmacol. Toxicol. 25 : 171-191.

Itoh, T., Kanmura, Y., Kuriyama, H., and Sasaguri, T. (1985). Nitroglycerine- and isoprenaline-induced vasodilatation : assessment from the actions of cyclic nucleotides. Br. J. Pharmacol. 84 : 393-406.

Jackson, C.V., Pope, T.K., and Lucchesi, B.R. (1987).

Coronary artery vasodilation in the canine :
physiological and pharmacological roles of
 β -adrenergic receptors. J. Cardiovasc. Pharmacol.
10 : 196-124.

Johansson, B., and Somlyo, A.P. (1980). Electrophysiology
and E-C coupling. In D.F. Bohr, A.P. Somlyo, and
H.V. Sparks Jr. (eds.), Handbook of Physiology :
The CVS, pp. 301-323. Baltimore : American
Physiological Society.

Kamata, K., Miyata, N., and Kasuya, Y. (1988). Involvement
of endothelial cells in relaxation and contraction
responses of the aorta to isoproterenol in naive
and streptozotocin-induced diabetic rats. J.
Pharmacol. Exp. Ther. 249 : 890-894.

Kelly, M.J. (1972). Receptors for dopamine in some
isolated vascular tissues of the dog. Br. J.
Pharmacol. 46 : 575P-576P.

_____. (1981). Effect of isoprenaline on dopamine
receptors in the rabbit isolated renal artery.
Experimentia 37 : 889-890.

Kenakin, T.P. (1984). The classification of drugs and
drug receptors in isolated tissues. Pharmacol.
Rev. 36 : 165-222.

Ketabian, J.W., and Calne, D.B. (1979). Multiple receptors for dopamine. Nature 277 : 93-96.

Khayyal, M., Gross, F., and Kreye, V.A.W. (1981). Studies on the direct vasodilator effect of hydralazine in isolated rabbit renal artery. J. Pharmacol. Exp. Ther. 218 : 390-394.

Kipshidz, N.N., Nresheret, A.P., and Khomaziuk, A.I. (1982). Effect of dopamine on coronary circulation. Kardiologija 22 : 60 (Abstr.).

Kokuvetz, W.R., and Poch, G. (1970). Inhibition of cyclic-3'-5'-nucleotide-phosphodiesterase as a possible mode of action of papaverine and similarly acting drugs. Naunyn-Schmiedeberg's Arch. Pharmacol. 267 : 189-194.

Konishi, M., Toda, N., and Yamamoto, M. (1981). Different mechanisms of action of histamine in isolated artery of the dog. Br. J. Pharmacol. 74 : 111-118.

Kreye, V.A.W. (1984). Direct vasodilators with unknown modes of action : the nitrocompound and hydralazine. J. Cardiovasc. Pharmacol. 6 : S646-S655.

Langer, S.Z., and Hicks, P.E. (1984). Alpha-adrenoceptor subtype in blood vessels : physiology and pharmacology. J. Cardiovasc. Pharmacol. 6 : S547-S558.

Lang, W.J., and Woodman, O.L. (1982). Comparison of the vasodilator action of dopamine and dopamine agonists in the renal and coronary beds of the dog. Br. J. Pharmacol. 77 : 23-28.

Lee, M.R. (1982). Dopamine and kidney. Clin. Sci. 62: 439-448.

Luscher, T.F., Neves, R., and Vanhoutte, P.M. (1986). Endothelium-dependent relaxation to acetylcholine and adenosine diphosphate in human renal arteries Kidney Int. 29 : 29 (Abstr.).

_____. and Vanhoutte, P.M. (1988). Endothelium-dependent responses in human blood vessels. TIPS 9 : 181-184.

Macedo, T.R.A., Fontes Ribeiro, C.A., Profirio, M.H. and Cotrim. (1988). Endothelium-dependent relaxing effect of apomorphine in the human uterine artery. Br. J. Pharmacol. 95 : 719P.

McGrath, J.C. (1982). Evidence for more than one type of postjunctional α -adrenoceptor. Biochem. Pharmacol. 31 : 467-484.



McLean, A.J., Souich., P., Barron, K.W., and Briggs, A.H. (1978). Interaction of hydralazine with tension development and mechanisms of calcium accumulation in K^+ -stimulated rabbit aortic strips. J. Pharmacol. Exp. Ther. 207 : 40-48.

McNay, J.L., and Goldberg, L.I. (1966). Comparison effect of dopamine, isoproterenol, norepinephrine and bradykinin on canine renal blood flow. J. Pharmacol. Exp. Ther. 151 : 23-31.

McNay, J.L., McDonald, R.H., and Goldberg, L.I. (1965). Direct renal vasodilation produced by dopamine in the dog. Cir. Res. 16 : 510-517.

Mellander, S., and Johansson, B. (1968). Control of resistance, exchange, and capacitance functions in the peripheral circulation. Pharmacol. Rev. 20 : 117-196.

Miller, V.M., and Vanhoutte, P.M. (1985). Endothelial alpha-2 adrenoceptors in canine pulmonary and systemic blood vessels. Eur. J. Pharmacol. 118 : 123-129.

Moncada, S., Radomski, M.W., and Palmer, R.M.J. (1988). Endothelium-derived relaxing factor : identification as nitric oxide and role in the control of vascular tone and platelet function. Biochem. Pharmacol. 37 : 2495-2501.

Moritoki, H., Hosoki, E., and Ishida, Y. (1986). Age-related decrease in endothelium-dependent dilator response to histamine in rat mesenteric artery. Eur. J. Pharmacol. 126 : 61-67.

Murad, F. (1986). Cyclic uanosine monophosphate as a mediator of vasodilation. J. Clin. Invest. 78 : 1-5.

Nakaki, T., Roth, B.L., chuang, D.M., and Costa, E. (1985). Phasic and tonic components in 5-HT₂ receptor-mediated rat aorta contraction. Participation of Ca²⁺ channels and phospholipase C. J. Pharmacol. Exp. ther. 243 : 442-446.

Nielsen, H., Mortense, F.V., and Mulvany, M.J. (1990). Responses to noradrenaline in human subcutaneous resistance arteries are mediated by both α₁- and α₂-adrenoceptors. Br. J. Pharmacol. 99 : 31-34.

Nyborg, N.C.B., and Mikkelsen, E.O. (1985). Characterization of β-adrenoceptor subtype in isolated ring preparations of intramural rat coronary small arteries. J. Cardiovasc. Pharmacol. 7 : 1113-1117.

Ohlstein, A.J., Zabko-Potatovich, B., and Berkowitz, B.A. (1984). Studies on vascular dopamine receptors with the dopamine receptor agonist SKNF 82526. J. Pharmacol. Exp. Ther. 229 : 433.

Okamura, T., Miyazaki, M., and Toda, N. (1986). Responses of isolated dog blood vessels to glucagon.
Eur. J. Pharmacol. 125 : 395-401.

Oriowo, M.A., and Bevan, J.A. (1987). Characterization of histamine H₁ and H₂ receptors in the rabbit isolated ovarian artery and vein. J. Cardiovasc. Pharmacol. 10 : 76-81.

Ottosson, A., Jansen, I., and Edvinsson, L. (1988). Characterization of histamine receptors in isolated human cerebral arteries. Br. J. Pharmacol. 94 : 901-907.

Ozaki, N., Bito, K., Kinoshita, M., and Kawakita, S. (1983). Effects of a cardiotonic agents, TA-064, on isolated canine cerebral, coronary, femoral, mesenteric, and renal arteries. J. Cardiovasc. Pharmacol. 5 : 818-821.

Palmer, R.M.J., Ashton, D.S., and Moncada, S. (1988). Vascular endothelial cells synthesize nitric oxide from L-arginine. Nature 33 : 664-666.

_____. Ferridge, A.G., and Moncada, S. (1987). Nitric oxide release accounts for the biological activity of endothelium derived relaxing factor. Nature 327 : 524-526.

- Peach, M.J., Singer, H.A., and Loeb, A.L. (1985). Mechanisms of endothelium-dependent vascular smooth muscle relaxation. Biochem. Pharmacol. 34 : 1867-1874.
- Petersen, E.N. (1981). Pre- and postsynaptic α -adrenoceptor antagonism by neuroleptics in vivo. Eur. J. Pharmacol. 69 : 399-405.
- Poch, G., and Kukovetz, W.R. (1971). Papaverine-induced inhibition of phosphodiesterase activity in various mammalian tissue. Life Sci. 10 : 133-144.
- Purdy, R.E., Murray, D.L., and Stupecky, G.L. (1987). Receptors for 5-hydroxytryptamine in rabbit blood vessels : activation of alpha-adrenoceptors in rabbit thoracic aorta. J. Pharmacol. Exp. Ther. 240 : 535-541.
- Rapoport, R.M., and Murad, F. (1983). Agonist-induced endothelium-dependent relaxation in rat thoracic aorta may be mediated through cGMP. Cir. Res. 52 : 352-357.
- Reinsberg, J., and Kullmann, R. (1986). Characterization of vascular dopamine receptors in the gastric circulation of the rabbit. J. Cardiovasc. Pharmacol. 8 : 1067-1073.

Rhodin, J.A.G. (1980). Architecture of the vessel wall.

In D.F. Bohr, A.P. Somlyo, H.V. Sparks Jr. (eds.),
Handbook of Physiology : the CVS, pp. 1-32,
Baltimore : American Physiological Society.

Robinson, B.F. and Collier, J.G. (1979). Vascular smooth muscle : correlations between basic properties and responses of human blood vessels. Br. Med. Bull. 35 : 305-312.

Rubanyi, G., and Vanhoutte (1985). Endothelium-removal decreases relaxations of canine coronary arteries caused by β -adrenergic agonists and adenosine.
J. Cardiovasc. Pharmacol. 7 : 139-144.

Sakakibara, Y., Fujiwara, M., and Muramatsu, I. (1982). Pharmacological of the alpha-adrenoceptors of the dog basilar artery. Naunyn-Schmiedeberg's Arch. Pharmacol. 319 : 1-7.

Salom, J.B., Torregrosa, G., Miranda, F.J., Alabadi, J.A., Alvarez, C., and Alborch, E. (1990). Effects of Ca^{2+} entry blockers on $CaCl_2$, KCl and noradrenaline-induced contractions of goat cerebral arteries.
Eur. J. Pharmacol. 177 : 1-8.

Satoh, H., and Inui, J. (1984). Endothelial cell dependent relaxation and contraction induced by histamine in the isolated guinea-pig pulmonary artery. Eur. J. Pharmacol. 97 : 321-324.

- _____. and Sato, S. (1984). Prostaglandin E₂ and I₂ production in isolated dog renal arteries in the absence or presence of vascular endothelial cells. Biochem. Biophys. Res. Commun. 118 : 873-876.
- Savino, E.A. and Varela, A. (1991). Characterization of postsynaptic α -adrenoceptor in the isolated rat tail artery. Arch. int. Pharmacodyn. Ther. 309 : 137-146.
- Schaub, M.C., Kunz, B. (1986). Regulation of contraction in cardiac and smooth muscles. J. Cardiovasc. Pharmacol. 8 : S117-S123.
- Scheid, C.R., Honeyman, T.W., and Fay, F.S. (1979). Mechanism of β -adrenergic relaxation of smooth muscle. Nature 277 : 32-36.
- Schmidt, M., and Imbs, J.L. (1980). Pharmacological characterization of renal vascular dopamine receptors. J. Cardiovasc. Pharmacol. 2 : 595-605.
- _____. Imbs, J.L., and Schwartz, J. (1981). The vascular dopamine receptor : a review. J. Pharmacol. (Paris). 12 : 355-382.
- Schmitz, J.M., Graham, R.M., Sagalowsky, A., and Pettinger, W.A. (1981). Renal alpha-1 and alpha-2 adrenergic receptors : biochemical and pharmacological correlations. J. Pharmacol. Exp. Ther. 219 : 400-406.

Schoeffter, P., Dion, R., and Godfraind, T. (1988). Modulatory role of the vascular endothelium in the contractility of human isolated internal mammary artery. Br. J. Pharmacol. 95 : 531-543.

_____. and Godfraind, T. (1991). Characterization of histamine-induced contraction in rat isolated aorta. Eur. J. Pharmacol. 197 : 193-200.

Seidel, C.L., and LaRochella (1987). Venous and arterial endothelium : different dilation abilities in dog vessels. Cir. Res. 60 : 626-630.

Singer, H.A., and Peach, M.J. (1983). Endothelium dependent relaxation of rabbit aorta: I. Relaxation stimulated by arachidonic acid. J. Pharmacol. Exp. Ther. 226 : 790-795.

Somlyo, A.V., Bond, M., Somlyo, A.P., and Scarpa, A. (1985). Inositol-triphosphate (InsP_3) induced calcium release and contraction in vascular smooth muscle. Proc. Natl. Acad. Sci., U.S.A. 82 : 5231-5235.

_____. and Somlyo, A.V. (1968). Vascular smooth muscle. I. Normal structure, pathology, biochemistry and biophysics. Pharmacol. Rev. 20 : 197-272.

Steinberg, S.F., Jaffe, E.A., and Bilezikian, J.P. (1984). Endothelial cells contain beta-adrenoceptors. Naunyn-Schmiedeberg's Arch. Pharmacol. 325:310-313.

Stitzel, R.E., and Robinson, R.L., (1990). Adrenomimetic drugs. In C.R. Craig and R.E. Stitzel (eds.), Modern Pharmacology, pp. 136-154. 3rd ed. Boston : Little, Brown and Company.

Takenaka, F., and Morishita, H. (1976). Possible mechanisms involved in the coronary vasodilator response to dopamine. Arch. int. Pharmacodyn. Ther. 222 : 81-93.

Tayo, F.M., and Bevan, J.A. (1986). Pharmacological characterization of histamine receptors in the rabbit renal artery. Eur. J. Pharmacol. 121 : 129-133.

Thom, S., Hughes, A., Martin, G., and Sever, P.S. (1987). Endothelium -dependent relaxation in isolated human arteries and veins. Clin. Sci. 73 : 547-552.

Toda, N. (1974). The action of vasodilating drugs on isolated basilar, coronary and mesenteric arteries of the dog. J. Pharmacol. Exp. Ther. 1901 : 139-146.

_____. (1976). Influence of dopamine and noradrenaline on isolated cerebral artery of the dog. Br. J. Pharmacol. 58 : 121-126.

- _____. (1983a). Heterogeneity in the response to dopamine of monkey cerebral and peripheral arteries. Am. J. Physiol. 245 : H930-H936.
- _____. (1983b). Isolated human coronary arteries in response to vasoconstrictor substances. Am. J. Physiol. 245 : H939-H941.
- _____. (1984). Endothelium-dependent relaxation induced by angiotensin II and histamine in isolated arteries of dog. Br. J. Pharmacol. 81 : 301-307.
- _____. (1986). Mechanisms of histamine-induced relaxation in isolated monkey and dog coronary arteries. J. Pharmacol. Exp. Ther. 239 : 529-535.
- _____. (1987). Mechanism of histamine actions in human coronary arteries. Circ. Res. 61 : 280-286.
- _____. (1990). Mechanism underlying responses to histamine of isolated monkey and human cerebral artery. Am. J. Physiol. 258 : H311-H317.
- _____. and Fujita, Y. (1973). Responsiveness of isolated cerebral and peripheral arteries to serotonin, norepinephrine and transmural stimulation. Cir. Res. 38 : 98-104.

_____. and Goldberg, L.I. (1973). Dopamine-induced relaxation of isolated arterial strips. J. Pharm. Pharmacol. 25 : 587-589.

_____. and Goldberg, L.I. (1975). Effects of dopamine on isolated canine coronary artery. Cardiovasc. Res. 9 : 384-389.

_____. and Hatano, Y. (1979). Antagonism by droperidol on dopamine-induced relaxation in isolated dog arteries. Eur. J. Pharmacol. 57 : 231-238.

_____. Okamura, T., Nakajima, M., and Miyazaki, M. (1984). Modification by yohimbine and prazosin of the mechanical response of isolated dog mesenteric, renal and coronary arteries to transmural stimulation and norepinephrine. Eur. J. Pharmacol. 98 : 69-78.

Towart, R. (1982). Effects of nitrendipine (BAY e 5009), nifedipine, verapamil, phentolamine, papaverine and minoxidil on contractions of isolated rabbit aortic smooth muscle. J. Cardiovasc. Pharmacol. 4 : 895-902.

Ueda, S., Yano, S., and Sakanashi, M. (1982). In vitro evidence for dopaminergic receptors in human renal artery. J. Cardiovasc. Pharmacol. 4 : 76-81.

Uma, S., Tuncer, M., and Kayaalp, S.O. (1987). The subtype of 5-HT receptors in the rat pulmonary artery.
Arch. int. Pharmacodyn. Ther. 288 : 248-255.

Van Breeman, C., Cauvin, C., Yamamoto, H., and Zshauer, A. (1987). Vascular smooth muscle calcium channels.
J. Cardiovasc. Pharmacol. 10 : S10-S15.

Vanhoutte, P.M. (1977). Cholinergic inhibition of adrenergic neurotransmission. Fed. Proc. 36 : 2444-2449.

_____. (1981). Pharmacology of the blood vessel wall. J. Cardiovasc. Pharmacol. 3 : 1351-1369.

_____. (1988). The endothelium-modulator of vascular smooth muscle tone. N. Eng. J. Med. 319 : 512-513.

_____. Cohen, R.A., and Van Nueten, J.M. (1984). Serotonin and arterial vessels. J. Cardiovasc. Pharmacol. 6 : S421-S428.

_____. and Miller, V.M. (1989). alpha-2 adrenoceptor and EDRF. Am. J. Med. 87 : 1S-5S.

_____. Rubanyi, G.M., Miller, V.M., and Houston, D.S. (1986). Modulation of vascular smooth muscle contraction by the endothelium. Annu. Rev. Physiol. 48 : 308-320.

Van de Voorde, J., and Leusen, I. (1982). Vascular endothelium and the relaxation effect of histamine on aorta of the rat. Arch. int. Pharmacodyn. Ther. 251 : 329-330.

_____. and Leusen, I. (1983). Role of the endothelium in the vasodilator response of rat thoracic aorta to histamine. Eur. J. Pharmacol. 87 : 113-120.

_____. and Leusen, I. (1984). Effect of histamine on aorta preparations of different species. Arch. int. Pharmacodyn. Ther. 268 : 95-105.

Van Nueten, J.M., Janssen, P.A.J., VanBeek, J., Xhonneux, R., Verbeuren, T.J., and Vanhoutte, P.M. (1981). Vascular effects of ketanserin (R41468), a novel antagonist of 5-HT₂-serotonergic receptors. J. Pharmacol. Exp. Ther. 218 : 217-230.

_____. Leysen, J.E., de Clerck, F., and Vanhoutte, P.M. (1984). Serotonergic receptor subtypes and vascular reactivity. J. Cardiovasc. Pharmacol. 6 : S564-S574.

Van Rossum, J.M. (1963). Cumulative dose-response curves. II. Technique for the making of dose-response curves in isolated organs and the evaluation of drug parameter. Arch. int. Pharmacodyn. Ther. 143 : 229-330.

Warner, T.D. (1990). Simultaneous perfusion of rat isolated superior mesenteric arterial and venous beds : comparison of their vasoconstrictor and vasodilator responses to agonists. Br. J. Pharmacol. 99 : 427-433.

Woodman, O.L., and Lang, W.J. (1980). Dopamine-induced vasodilation in the isolated perfused rat kidney. Arch. int. Pharmacodyn Ther. 243 : 228-235.

Wolff, D.W., Buckalew, V.M., Jr., and Stranhouy, J.W. (1984). Renal α_1 - and α_2 -adrenoceptor mediated vasoconstriction in dogs : comparison of phenylephrine, clonidine, and guanabenz. J. Cardiovasc. Pharmacol. 6 : S793-S798.

Wright, C.E., and Angus, J.A. (1987). Diverse vascular response to serotonin in the conscious rabbit : effects of serotonin antagonists on renal artery spasm. J. Cardiovasc. Pharmacol. 10 : 415-423.

Yamauchi, M., Kobayashi, Y., Hattori, K., Yamada, K., and Nakase, A. (1989). Dopamine-induced relaxation in human pulmonary arteries. Experimentia 45 : 150-152.

Yang, Z.H., and Others (1989). Endothelium-derived relaxing factor and protection against contractions induced by histamine and serotonin in the human internal mammary artery and in the sphenous vein. Circulation 80 : 1041-1048.

Yeh, B.K., McMAY, J.L. and Goldberg, L.I. (1969). Attenuation
of dopamine renal and mesenteric vasodilation by
haloperidol : evidence for specific dopamine receptor.
J. Pharmacol. Exp. Ther. 168 : 303-309.

Yelmosky, J., Katz, R., and Dietrich, E.V. (1964). A study
of some of the pharmacologic actions of droperidol.
Toxicol. Appl. Pharmacol. 6 : 37-47.

ประวัติผู้เขียน

นางสาวนิศากร ไกรสุรย์ เกิดวันที่ 6 กรกฏาคม พ.ศ. 2507
ที่อำเภอสตึก จังหวัดบุรีรัมย์ สำเร็จการศึกษาปริญญาตรีวิทยาศาสตรบัณฑิต สาขา
ชีววิทยา ภาควิชาชีววิทยา คณะวิทยาศาสตร์ มหาวิทยาลัยขอนแก่น ในปีการ
ศึกษา 2528 และเข้าศึกษาต่อในหลักสูตรวิทยาศาสตรบัณฑิต ที่จุฬาลงกรณ์
มหาวิทยาลัย เมื่อ พ.ศ. 2531



ศูนย์วิทยทรัพยากร
อุดหนุนศึกษาและพัฒนา
ศูนย์วิทยาศาสตร์และเทคโนโลยี