

## REFERENCES

- Adams, R.F., Vandemark, F.L., and Schmidt, G.J. More sensitive high-pressure liquid - chromatographic determination of theophylline in serum. Clinical Chemistry 22 (1976) : 1903-1906.
- Aviram, M., Tal, A., Ben-Zvi, Z., and Gorodischer, R. Monitoring theophylline therapy using citric acid- stimulated saliva in infants and children with asthma. Pediatrics 8 (December 1987) : 894-897.
- Bock, J.L., Lam, S. and Karmen, A. Therapeutic drug monitoring using high - speed liquid chromatography and rapid sample preparation: an assay for serum theophylline. Journal of Chromatography. 308 (1984) : 354-358.
- Bonham, A., Hendeles, L., Vaughan, L., and Weinberger, M. The reliability of serum theophylline determinations from clinical laboratories. Am. Rev. Respir Dis. 122 (1980) : 829-832.
- Butrimovitz, G.P., and Raisys, V.A. An improved micromethod for theophylline determination by reversed - phase liquid chromatography. Clinical Chemistry 25 (1979) : 1461-1464.

Chiou, W.L., Chang, K., and Peng, G.W. Precaution in the monitoring of drug levels in saliva: Abnormal salicylate levels after oral dosing. J. Clin. Pharmac. 8 (1976) : 459-463.

Chrzanowski, F.A. Gas chromatographic analysis of theophylline in human serum. Biochem. Med. 11 (1974) : 26-31.

Cohen, A., Johnson, C., and Osvaldo, R.E. A rapidly dissolving theophylline tablet. Curr. Ther. Res. 17 (1975) : 497-505.

Cohen, J.L. Theophylline. In K.Florey(ed.), Analytical profiles of drug substances, pp. 466-493. New York: Academic Press, 1975.

Culig, J., Johnson, A., and Turner, P. Saliva theophylline concentrations after a single oral dose. Br. J. Clin. Pharmac. 13 (1982) : 243-245.

Danhof, M., and Breimer, D.D. Therapeutic drug monitoring in saliva. In M. Gibaldi and L. Prescott (eds.), Clinical Pharmacokinetics, Australia: ADIS Health Science, 1983.

Dusci, L.J., Hackett, L.P., and McDonald, I.A. Gas-liquid chromatographic determination of theophylline in human plasma. Journal of Chromatography 104 (1975) : 147-150.

Ellis, E.F., Koysooko, R. and Levy, G. Pharmacokinetics of theophylline in children with asthma. Pediatrics 58 (1976) : 542-547.

Eney, R.D., and Goldstein, E.O. Compliance of chronic asthmatics with oral administration of theophylline as measured by serum and salivary levels. Pediatrics 57 (April 1976) : 513-517.

Evenson, M.A., and Warren, B.L. Serum theophylline analysis by high-pressure liquid chromatography. Clinical Chemistry 22 (1976) : 851-855.

Franconi, L.C., Hawk, G.L., Sandmann, B.J. and Haney, W.G. Determination of theophylline in plasma ultrafiltrate by reversed phase high pressure liquid chromatography. Analytical Chemistry. 48 (February 1976) : 372-375.

Galant, S.P., Gillman, S.A., Cummins, L.H., Kozak, P.P. and Orcutt, J.J. Reliability of salivary theophylline as a guide to plasma theophylline levels. Am. J. Dis. Child. 131 (1977) : 970-976.

Glynn, J.P., and Bastain, W. Salivary excretion of paracetamol in man. J. Pharm. Pharmacol. 25 (1973) : 420-421.

Habib, M.P., Schifman, R.B., Shon, B.Y., Fastro, J.F., and Campbell, S.C. Evaluation of whole blood theophylline enzyme immunochromatography assay. Chest 92 (July 1987) : 129-131.

Hendeles, L., Massanari, M., and Weinberger, M. Theophylline. In W.E. Evan, J.J. Schentag and W.J. Jusko (eds.), Applied pharmacokinetics : Principles of therapeutic drug monitoring, pp.1105-1181. Washington : Applied Therapeutics, 1986.

Hendeles, L., Weinberger, M., and Johnson, G. Monitoring serum theophylline levels. In M.Gibaldi and L.Presscott (eds.), Clinical pharmacokinetics, Australia : ADIS Health Science, 1983.

Hill, R.E. Retention behaviour of a bonded reversed phase in a high performance liquid chromatographic assay of serum theophylline. Journal of Chromatography 135 (1977) : 419-425.

Jenne, J.W., Wyze, E., Rood, F.S., and MacDonald, F.M. Pharmacokinetics of theophylline : Application to adjustment of the clinical dose of aminophylline: Clin. Pharmacol. Ther. 13 (1972) : 349-360.

Johnson, G.F., Dechtiaruk, W.A., and Solomon, H.M. Gas-chromato-graphic determination of theophylline in human serum and saliva Clinical Chemistry 21 (1975) : 144-147.

Killman, S.A, and Thaysen, J.H. The permeability of the human parotid gland to a series of sulfonamide compounds, paraminohippurate and inulin, Scand. J. Clin. Lab. Invest. 7 (1955) : 86-91.

Koysooko, R., Ellis, E.F., and Levy, G. Relationship between theophylline concentration in plasma and saliva of man. Clin. Pharmacol. Ther. 15 (1974) : 454-460.

Least, C.J., Johnson, G.F., and Solomon, H.M. Gas chromatographic micro-scale procedure for theophylline, with use of nitrogen-sensitive detector. Clinical Chemistry 22 (1976) : 765-768.

Leung, P., Kalisker, A. and Bell, T.D. Variation in theophylline clearance rate with time in chronic childhood asthma. J. Allergy Clin. Immunol. 59 (June 1977) : 440-444.

Levy, G. Ellis, E.F. and Koysooko, R. Indirect plasma-theophylline monitoring in asthmatic children by determination of theophylline concentration in saliva. Pediatrics 53 (1974) : 873-876.

Loomis, K.F. , and Frye, R.M. Evaluation of the Abbott TDx<sup>TM</sup> for the stat measurement of phenobarbital, phenytoin, carbamazepine and theophylline. Am. J. Clin. Pathol. 80 (1983) : 686-691.

Matheson, L.E., Bighley, L., and Hendeles, L. Drug interference with the Schack and Waxler plasma theophylline assay. Am. J. Hosp. Pharm. 34 (1977) : 496-499.

Mathews, S.J., Schneiwiss, F., and Cersosimo, R.J. A clinical manual of adverse drug reactions. USA : Appleton Crofts, 1986.

McDonald, J.M., and Ladenson, J.H. Theophylline toxicity. Clinical Chemistry. 24 (1978) : 1603-1608.

Montri Tuchinda, Suprida Habanananda, Nualonong Srimaruta, Amnuay Thithapandha, and Katharee Chaichanwathanakul. Rational dosage of intravenous aminophylline for Thai asthmatic children. J. Med. Ass. Thailand. 67 (1984) : 170-175.

Montri Tuchinda, Renu Koysooko, Suprida Habanananda, Jongkonnee Wattanapermpool, and Napa Arunyanark. Relationship between theophylline concentration in serum and saliva in Thai asthmatic children. Siriraj Hosp. Gaz. 39 (1987) : 245-252.

Nelson, J.W., Cordry, A.L. Aron, C.G., and Bartell, R.A. Simplified micro-scale procedure for preparing samples for theophylline determination by liquid chromatography. Clinical Chemistry 23 (1977) : 124-126.

Novak, L.P., Ageing, total body potassium, fat-free mass, and cell mass in males and females between ages 18 and 85 years, J.Gerontology. 27(1972) : 438-443.

Ohmori, H. et al. Study of bioavailability and pharmacokinetics of theophylline following administration of two sustained release dosage forms as assessed by salivary data: Part II. Int. J.Clin. Pharmacol. Ther. Toxicol. 24 (1986) : 196-201.

Orcutt, J.J., Kozak, P.P. Gillman, S.A. and Cummins, L.H. Micro-scale method for theophylline in body fluids by reversed phase, high-pressure liquid chromatography. Clinical Chemistry 23 (1977) : 599-601.

Paxton, J.W., and Foote, S. Aberrantly high phenytoin concentrations in saliva:Precautions in monitoring phenytoin concentration in whole saliva. Br. J.Clin. Pharmac. 8 (1979) 508-509.

Piafsky, K.M., and Ogilvie, R.I. Dosage of theophylline in bronchial asthma. N. Eng. J. Med. 292 (June 1975) : 1218-1222.

Posti, J. Saliva-plasma drug concentration ratios during absorption: Theoretical considerations and pharmacokinetic implications. Pharmacutica Acta Helveticae 3 (1982) : 3-12.

Rall, T.W. The methylxanthines. In A.G. Gilman, L.S. Goodman and A.Gilman (eds.), The pharmacological basis of therapeutics. pp. 589-603. New York: MacMillan Publishing, 1985.

Reynold, J.E.F. Martindale: The extra pharmacopoeia. 29 th ed.

London: Pharmaceutical Press, 1989.

Rowe, D.J.F., Watson, I.D., Williams, J. and Berry, D.J. The clinical use and measurement of theophylline. Ann. Clin. Biochem. 25 (1988) :4-26.

Saming Kaojaroen. The importance of pharmacokinetic in patient treatment. Proceedings Symposium on Clinical Pharmacokinetics, pp. 5-18. Mahidol University, 1988.

Schack, J.A. and Waxler, S.H. An ultraviolet spectrophotometric method for the determination of theophylline and theobromine in blood and tissues. J. Pharm. Exp. Ther. 97 (1949) : 283-291.

Schwertner, H.A., Wallace, J.E. and Blum, K. Improved ultraviolet spectrophotometry of serum theophylline. Clinical Chemistry 24 (1978) : 360-361.

Shah, V.P. et. al. Analytical methods validation: bioavailability, bioequivalence and pharmacokinetic studies. Pharmaceutical Research 9 (1992) : 588-592.

Shah, V.P., and Riegelman, R. GLC determination of theophylline in biological fluids. J. Pharm. Sci. 63 (August 1974) : 1283-1285.

Shargel, L., and A.B.C. Yu., Applied Biopharmaceutics and Pharmacokinetics. New York:Appleton-Century-Crofts, 1985.

Silva, J.A.F. Analytical strategies for therapeutic monitoring of drugs in biological fluids. Journal of Chromatography 340 (1985) : 3-30.

Smith, R.V., and Stewart, T.T. Textbook of biopharmaceutic analysis. Philadelphia: Lea-Febiger, 1981.

Soifer, H. Aminophylline Toxicity. J. Pediatrics. 50 (1957) : 657- 669.

Soldin, S.J. and Hill, J.G. A rapid micromethod for measuring theophylline in serum by reverse-phase high-performance liquid chromatography. Clin. Biochem. 10 (1977) : 74-77.

Szepesi, G. HPLC in pharmaceutical analysis; vol I: General consideration. Florida: CRC Press, 1990.

Wan, S.H., Matin, S.B., and Azarnoff, D.L. Kinetics, salivary excretion of amphetamine isomers, and effect of urinary pH. Clin. Pharmacol. Ther. 23 (1978) : 585-590.

Welch, R.M., DeAngelis, R.L., Wingfield, M. and Farmer, T.W. Elimination of antipyrine from saliva as a measure of metabolism in man. Clin. Pharmacol. Ther. 18 (1975) : 249-258.

Wilson, J.D. et al. Harrison's principles of internal medicine. 12 th ed.  
vol. 1, 2. USA:McGraw-Hill, 1991.

Windholz, M., Budavari, S., Blumetti, R.F., and Otterbein, E.S. Merck  
Index. 9 th ed. New Jersey : Merck & Co., 1976.

ศูนย์วิทยาพรพยาบาล  
จุฬาลงกรณ์มหาวิทยาลัย



## **APPENDICES**

ศูนย์วิทยทรัพยากร  
จุฬาลงกรณ์มหาวิทยาลัย

**Appendix A. Physiological characteristics and administered theophylline dose of volunteer subjects**

subject number	sex	age (yr)	height (cm.)	weight (kg.)	dose (mg.)
1	M*	30	175.0	58.0	300
2	M*	30	171.0	61.0	300
3	M*	40	170.0	62.0	350
4	M*	32	165.0	60.0	300
5	M*	31	156.0	50.0	250
6	M*	29	162.0	54.0	300
7	M*	34	159.0	50.0	250
8	M	22	165.0	54.0	300
9	M	22	167.0	57.0	300
10	M	20	180.0	68.0	350
11	M	40	167.0	69.0	350
12	M	36	167.0	50.0	250
13	M	26	178.0	58.0	300
14	M	25	174.0	60.0	300
15	M	26	170.0	60.0	300
16	M	26	170.0	60.0	300
17	M	26	168.0	59.0	300
18	M	28	171.0	61.0	300
$\bar{X}$ (S.D)	-	29 (6)	168.6(6.1)	58.4(5.4)	-

Appendix A. (continue)

subject number	sex	age (yr)	height (cm.)	weight (kg.)	dose (mg.)
19	F	23	164.0	61.0	300
20	F	23	160.0	45.0	250
21	F	27	159.0	54.0	300
22	F	21	162.0	51.0	300
23	F	40	158.0	52.0	300
24	F	24	157.0	48.0	250
25	F	24	160.0	45.0	250
26	F	24	158.0	46.0	250
27	F	34	164.0	58.0	300
28	F	35	163.0	65.0	350
29	F	22	158.0	50.0	250
30	F	25	165.0	67.0	350
31	F	29	160.0	54.0	300
32	F	23	157.0	44.0	250
33	F	28	160.0	57.0	300
34	F	35	152.0	40.0	200
35	F	37	155.0	43.0	250
36	F	30	171.0	61.0	300
$\bar{X}$ (S.D)	-	28 (6)	160.2(4.3)	52.3(7.9)	-

M\* = smoking male , M = non-smoking male , F = female

**Appendix B. Individual data of theophylline concentration in plasma of 36 volunteers**

subject number	sampling time (hour)								
	0	0.5	1.0	1.5	2.0	4.0	6.0	8.0	12.0
1	0	4.882	6.106	7.165	8.046	7.531	6.411	5.842	4.190
2	0	8.099	7.486	8.394	8.247	6.000	4.690	4.429	2.530
3	0	6.418	9.158	7.079	8.862	6.190	5.528	4.737	2.722
4	0	1.297	4.385	6.179	7.120	9.724	7.919	7.444	4.478
5	0	1.606	5.282	7.934	7.080	8.973	6.061	4.670	2.926
6	0	11.20	14.91	14.73	13.73	9.884	6.837	4.516	5.565
7	0	11.16	10.76	10.20	10.05	8.848	8.579	7.602	5.507
8	0	6.982	10.17	11.74	11.72	9.724	8.802	8.002	5.776
9	0	9.001	10.34	10.02	10.11	7.699	6.854	6.061	4.052
10	0	8.758	10.14	10.65	9.896	8.537	7.871	7.222	5.934
11	0	7.222	8.065	9.710	10.55	9.038	8.014	6.824	5.002
12	0	6.035	7.681	8.524	7.979	7.286	6.251	5.796	4.550
13	0	10.19	10.50	10.98	10.34	8.877	8.050	6.732	4.794
14	0	6.098	9.944	9.169	10.89	8.971	7.879	6.864	5.039
15	0	2.160	4.944	7.687	8.868	9.998	9.579	8.018	5.142
16	0	7.615	9.788	9.330	8.881	7.958	5.600	5.040	4.180
17	0	1.457	5.306	6.987	10.85	7.781	6.359	4.866	3.643
18	0	3.693	7.036	7.636	9.436	8.867	7.436	6.351	4.580
19	0	3.192	5.393	7.783	8.781	10.60	9.894	9.510	6.666
20	0	9.155	12.14	10.52	11.09	9.872	9.049	8.266	4.751
21	0	2.054	5.199	9.357	9.740	9.892	8.921	7.524	5.601
22	0	10.45	16.51	18.39	17.21	14.96	12.79	11.20	7.396
23	0	10.03	10.70	11.74	11.61	11.94	10.15	7.999	6.229
24	0	1.659	7.749	11.84	11.09	10.38	9.901	8.550	6.915
25	0	10.73	11.75	11.78	11.81	10.87	9.630	8.249	6.115
26	0	4.095	9.018	11.96	11.31	10.47	9.613	8.344	7.525
27	0	10.17	11.56	12.14	11.69	10.75	8.828	6.976	5.456
28	0	5.108	9.599	10.87	13.16	9.594	8.577	7.522	4.526
29	0	6.196	9.352	10.71	11.03	10.41	8.611	7.211	5.359
30	0	5.331	12.11	13.01	13.11	11.47	9.661	8.777	6.300
31	0	16.43	14.42	13.35	13.09	9.617	8.078	6.540	4.314
32	0	6.675	10.45	11.59	12.64	11.76	11.44	8.899	5.851
33	0	8.935	11.92	11.89	12.13	10.98	9.566	8.601	6.922
34	0	6.048	9.513	13.88	12.39	9.198	9.760	8.260	5.840
35	0	10.51	10.27	9.508	10.18	7.585	6.370	5.431	2.470
36	0	6.958	9.296	9.434	10.67	11.01	9.547	7.803	5.905

**Appendix C. Individual data of theophylline concentration in saliva of 36 volunteers**

subject number	sampling time (hour)								
	0	0.5	1.0	1.5	2.0	4.0	6.0	8.0	12.0
1	0.6178	3.409	4.232	4.432	4.541	5.186	3.702	3.188	2.545
2	0.4844	6.284	6.549	5.682	4.005	3.588	2.943	1.643	0.9995
3	0	5.052	2.548	4.801	3.424	4.982	3.154	1.996	2.074
4	1.505	2.050	5.628	3.276	4.161	7.918	5.803	4.094	2.970
5	0.3496	0.361	2.939	4.452	2.504	4.655	3.120	3.374	1.535
6	5.033	4.235	5.843	6.426	3.390	4.860	2.097	1.751	2.832
7	2.050	4.705	4.826	5.079	4.458	4.999	2.902	5.217	2.010
8	0.4830	5.293	6.421	7.441	5.703	5.259	4.985	4.119	2.990
9	0	6.133	3.329	4.649	6.010	3.931	5.598	4.045	3.146
10	1.213	5.419	5.973	6.547	5.299	4.122	3.058	4.192	3.480
11	0.3193	4.411	5.231	5.820	6.323	5.170	4.142	4.110	2.978
12	0	4.709	5.360	5.690	5.812	6.452	5.218	3.610	3.333
13	1.062	6.026	4.871	3.166	4.318	3.330	4.844	2.045	2.741
14	0	3.806	5.494	5.028	5.550	4.722	3.865	2.938	2.501
15	0.2854	1.694	4.604	5.807	5.473	4.909	4.390	3.232	3.667
16	0.7170	5.990	5.914	5.130	5.178	4.652	3.166	2.632	2.003
17	0.1792	1.196	3.604	3.932	6.068	4.723	3.642	3.091	1.822
18	0	3.629	4.206	5.205	5.448	4.279	3.696	3.621	2.382
19	0.4415	2.828	5.318	6.941	6.430	6.300	6.912	5.576	3.456
20	0.8614	6.218	5.872	5.607	5.238	4.825	4.465	4.860	3.316
21	0.2479	1.902	3.410	5.034	6.451	6.357	4.602	4.781	3.136
22	2.170	6.042	6.960	5.230	7.851	6.496	6.293	4.995	3.295
23	1.499	5.228	5.174	5.860	5.452	4.574	3.524	3.967	2.776
24	0.2169	1.510	7.873	6.706	7.591	8.606	6.817	4.890	3.245
25	1.674	9.150	7.469	7.620	7.739	6.417	5.436	4.717	3.604
26	1.097	5.265	7.322	7.659	10.54	7.480	6.651	5.357	3.494
27	0.7828	6.946	6.582	7.086	6.316	5.463	5.571	4.030	3.162
28	0.4014	4.296	6.855	6.087	7.132	6.829	2.838	4.648	2.893
29	5.923	4.203	4.122	4.922	1.194	5.078	4.776	3.561	2.855
30	1.339	2.803	8.586	5.328	7.806	4.141	4.678	5.902	3.886
31	0.1935	9.050	8.918	7.500	7.845	6.948	5.033	4.380	2.746
32	0.5809	5.131	6.970	6.669	6.294	5.870	4.256	4.339	2.779
33	0.8993	7.524	9.174	6.667	3.839	5.585	4.955	4.168	2.508
34	0.4975	4.902	7.263	11.04	6.674	7.526	5.182	4.337	2.675
35	1.248	6.516	4.043	6.675	3.747	3.398	3.032	3.569	2.116
36	2.702	4.646	5.633	6.174	5.299	5.986	5.312	3.662	3.279

**Appendix D. Individual data of saliva-plasma theophylline concentration ratio of 36 volunteers**

subject number	sampling time (hour)								
	0	0.5	1.0	1.5	2.0	4.0	6.0	8.0	12.0
1	0	0.70	0.69	0.62	0.56	0.69	0.58	0.54	0.61
2	0	0.78	0.87	0.68	0.48	0.60	0.63	0.37	0.40
3	0	0.79	0.28	0.68	0.39	0.80	0.57	0.42	0.76
4	0	1.60	1.30	0.53	0.58	0.81	0.73	0.55	0.66
5	0	0.22	0.56	0.56	0.35	0.52	0.51	0.72	0.52
6	0	0.38	0.39	0.44	0.25	0.49	0.31	0.39	0.51
7	0	0.42	0.45	0.50	0.44	0.56	0.34	0.69	0.36
8	0	0.76	0.63	0.63	0.49	0.54	0.57	0.51	0.52
9	0	0.68	0.32	0.46	0.59	0.51	0.82	0.67	0.78
10	0	0.62	0.59	0.61	0.54	0.48	0.39	0.58	0.59
11	0	0.61	0.65	0.60	0.60	0.57	0.52	0.60	0.60
12	0	0.78	0.70	0.67	0.73	0.88	0.83	0.62	0.73
13	0	0.59	0.46	0.29	0.42	0.38	0.60	0.30	0.57
14	0	0.62	0.55	0.55	0.51	0.53	0.49	0.43	0.50
15	0	0.78	0.93	0.76	0.62	0.49	0.46	0.40	0.71
16	0	0.79	0.60	0.55	0.58	0.58	0.56	0.52	0.48
17	0	0.82	0.68	0.56	0.56	0.61	0.57	0.64	0.50
18	0	0.98	0.60	0.68	0.58	0.48	0.50	0.57	0.52
19	0	0.89	0.99	0.89	0.73	0.59	0.70	0.59	0.52
20	0	0.68	0.48	0.53	0.47	0.49	0.49	0.59	0.70
21	0	0.93	0.66	0.54	0.66	0.64	0.52	0.64	0.56
22	0	0.58	0.42	0.28	0.46	0.43	0.49	0.45	0.44
23	0	0.52	0.48	0.50	0.50	0.38	0.35	0.50	0.44
24	0	0.91	1.00	0.57	0.68	0.83	0.69	0.57	0.47
25	0	0.85	0.64	0.65	0.66	0.59	0.56	0.57	0.59
26	0	1.30	0.81	0.64	0.93	0.71	0.69	0.64	0.46
27	0	0.68	0.57	0.58	0.54	0.51	0.63	0.58	0.58
28	0	0.84	0.71	0.56	0.54	0.71	0.33	0.62	0.64
29	0	0.68	0.44	0.46	0.11	0.49	0.55	0.49	0.53
30	0	0.52	0.71	0.41	0.60	0.36	0.48	0.67	0.62
31	0	0.55	0.62	0.56	0.60	0.72	0.62	0.67	0.64
32	0	0.77	0.67	0.58	0.50	0.50	0.37	0.49	0.48
33	0	0.84	0.77	0.56	0.32	0.51	0.52	0.48	0.36
34	0	0.81	0.76	0.80	0.54	0.82	0.53	0.52	0.46
35	0	0.62	0.39	0.70	0.37	0.45	0.48	0.66	0.86
36	0	0.67	0.61	0.65	0.50	0.54	0.56	0.47	0.56
X	0	0.70	0.61	0.59	0.54	0.58	0.54	0.54	0.56
S.D.	0	0.16	0.16	0.12	0.13	0.13	0.12	0.10	0.12

**VITA**

**Miss Chadarat Ampasavate was born on August 18th, 1968 in Chiang Mai , Thailand , graduated with a Bachelor of Science in Pharmacy (First Class Honors) from Chiang Mai University in 1991.**



ศูนย์วิทยบรพยากร  
จุฬาลงกรณ์มหาวิทยาลัย