



REFERENCES

1. Wagner, J.G. Biopharmaceutics and Relevant Pharmacokinetics. p. 99. Illinois : Drug Intelligence Publications, 1971.
2. American Pharmaceutical Association. The United States Pharmacopeia. 19th rev. Easton, Pa : Mack Publishing Co., 1975.
3. American Pharmaceutical Association. The National Formulary. 14th ed. Easton, Pa : Mack Publishing Co., 1975.
4. Jacob, J.T., and Plein, E.M. "Factors Affecting the Dissolution Rate of Medicaments from Tablets I." In Vitro Dissolution Rate of Commercial Phenobarbital Tablets J. Pharm.Sci. 57 (May 1968) : 798-805.
5. Wagner, J.G., et al. "Equivalence Lack in Digoxin Plasma Levels." JAMA. 224 (April 1973) : 199-204.
6. Pernarowski, M. "Dissolution Methodology." in Dissolution Technology., p. 61-62. Edited by L.J. Leeson and J.T. Carstensen. Washington, D.C. : I,P,T. Academy of Pharmaceutical Science, 1974.
7. Hirschorn, J.O., and Kornblum, S.S. "Dissolution of Poorly Water-Soluble Drugs. II : Exipient Dilution and Force

- of Compression Effects on Tablets of a Quinazalinone Compound." J.Pharm.Sci. 60 (March 1971) : 445-448.
8. Marlowe, E., and Shangraw, R.F. "Dissolution of Sodium Salicylate from Tablet Metrices Prepared by Wet Granulation and Direct Compression." ibid. 56(April 1967) : 498-504.
9. Shah, N., et al. "Influence of Dispersion Method on Dissolution Rate and Bioavailability of Digoxin. Triturations and Compressed Tablets II." ibid. 63 (March 1974) : 339-344.
10. Jacob, J.T., and Plein, E.M. "Factors Affecting Dissolution of Medicaments from tablets II. Effect of Binder Concentration, Tablet Hardness, and Storage Conditions on the Dissolution Rate of Phenobarbital from Tablets". ibid. 57(May 1968) : 802-805.
11. Sakr, A.M., and Elsabbagh, H.M. "Delayed Released in Compressed Nicotinic Acid Tablets". Mfg.Chem & Aerosol News. 43(December 1973) : 41-42, 45.
12. Shubair, M.S., and Dingwall, D. "Effects of Starch Concentration on Dissolution". ibid. 46(October 1976) : 52, 55-56.
13. Gibaldi, M., and Weintraub, H. "Dissolution of Salicylic Acid and Polyvinylpyrrolidone from Compressed Mix-

- ture". J.Pharm.Sci. 57(May 1968) : 832-835
14. Collett, J.H., and Kesteven, G."Some Effects of Polyvinylpyrrolidone on the Solubility and Dissolution Rate of Allopurinol". J.Pharm.Pharmac. 26, Suppl. (1974) : 84P-85P.
15. Florence, A.T., and Rahman, A. "Polyvinylpyrrolidones and Their Influence on Dissolution Rates of Compounds of Varying Aqueous Solubilities". ibid. 27, Suppl. (1975) : 55P.
16. Solvang, S., and Finholt, P. "Effect of Tablet Processing and Formulation Factors on Dissolution Rate of the Active Ingredient in Human Gastric Juice". J.Pharm.Sci. 59(January 1970) : 49-52.
17. Levy, G., and Gumtow, R.H. "Effect of Certain Tablet Formulation Factors on Dissolution Rate of the Active Ingredients III, Tablet Lubricant." J.Pharm.Sci. 52 (December 1963) : 1139-1144.
18. Nogami, H., Nagai, T., and Suzuki, A. "Studies on Powdered Preparations. XVII. Dissolution Rate of Sulfonamides by Rotating Disk Method." Chem.Pharm.Bull. 14(1966) : 329-338.

19. Bisaillon, S., and Tawashi, R. "Influence of Fluid Motion in Dissolution-Rate Determinations." J.Pharm.Sci. 60(December 1971) : 1874-1877.
20. Piccolo, J., and Tawashi, R., "Inhibited Dissolution of Drug Crystal by a Certified Water Soluble Dye." ibid. 59(January 1970) : 56-59.
21. Piccolo, J., and Tawashi, R. "Inhibited Dissolution of Drug Crystals by Certified Water Soluble Dyes III." ibid. 60(December 1971) : 1818-1820.
22. Block, L.H., and Patel, R.N. "Solubility and Dissolution of Triamcinolone Acetonide." ibid. 62(April 1973) : 617-621.
23. Gibaldi, M. "Biopharmaceutics" in Theory and Practice of Industrial Pharmacy., p. 20-21. Edited by L.Lachman H.A. Lieberman, J.L. Kanig. Philadelphia : Lea Febiger, 1976.
24. Florence, A.T., Elworthy, P.H., and Rahman, A. "The Influence of Solution Viscosity on the Rate of Soluble Salts, and the Measurment of an Effective Viscosity" J.Pharm.Pharmac. 25(June 1973) : 779-786.
25. Parrotte, E.L., and Sharma, V.K. "Dissolution Kinetics of Benzoic Acid in High Concentrations of Surface-Active Agents" J.Pharm.Sci. 56(October 1967) : 1341-1343.

26. Singh, P., et al. "Mechanistic Study of the Influence of Micelle Solubilization and Hydrodynamic Factors on the Dissolution Rate of Solid Drugs." ibid. 57 (June 1968) : 959-965.
27. Finholt, P., and Solvang., S. "Dissolution Kinetics of Drugs in Human Gastric Juice-The Role of Surface Tension." ibid. 57(August 1968) : 1322-1326.
28. Wurster, E.D., and Taylor, P.W., Jr. "Dissolution Kinetics of Certain Crystalline Forms of Prednisolone." ibid. 54(May 1965) : 670-676.
29. Tawashi, R. "Aspirin : Dissolution Rates of Two Polymorphic Forms." Science 160(March 1968) :76.
30. Higuchi, W.I., Bernado, P.D., and Mehta, S.C. "Polymorphism and Drug Availability II. Dissolution Rate Behaviors of the Polymorphic Forms of Sulfathiazole and Methyl-prednisolone." J.Pharm.Sci. 56(February 1967) : 200-207.
31. Aguiar, J.A., et al. "Effect of Polymorphism on the Absorption of Chloramphenicol from Chloramphenicol Palmitate." ibid. 56(July 1967) : 847-853.
32. Mullins, J.D., and Macek, T.J. "Some Pharmaceutical Properties of Novobiocin." J.Am.Pharm.Assoc. (Sci.Ed.) 49 (April 1960) : 245-248.

33. Shefter, E., and Higuchi, T. "Dissolution Behavior of Crystalline Solvated and Nonsolvated Forms of Some Pharmaceuticals." J.Pharm.Sci. 52(August 1963) 781-791.
34. Nelson, E. "Comparative Dissolution Rates of Weak Acids and Their Sodium Salts." J.Am. Pharm. Assoc. 47(April 1958) : 297-300.
35. Kornblum, S.S., and Hirschorn, J.O. "Dissolution of Poorly Water-Soluble Drugs I : Some Physical Parameter Related to Method of Micronization and Tablet Manufacture of Quinazolinone Compound." J.Pharm.Sci. 59 (May 1970) : 606-609.
36. Milosovich, G. "Determine of Solubility of a Metastable Polymorph" ibid. 53 (May 1964) : 484-487.
37. Noyes, A.A. and Whitney, W.R. "The Rate of Solution of Solid Substance in Their Own Solutions." J.Am.Chem.Soc. 19 (1897) : 930-934.
38. Nernst, W. and Brunner, E. "Theorie der Reakstionsgeschwindigkeit in Heterogenen Systemen. Reaktionsgeschwindigkeit in Heterogen Systemen." Zeit.Phys.Chem. 47(1904) : 52-55, 56-102.
39. Carstensen, J.T. "Theories of Dissolution-Single Particulate Systems." in Dissolution Technology, p. 3-4 Edited by L.J. Leeson and J.T. Carstensen. Washington, D.C.

- : I.P.T. Academy of Pharmaceutical Science, 1974.
40. Carstensen, J.T. Pharmaceutics of Solid Dosage Form,  
p. 63-86. New York : John Wiley & Sons, 1977.
41. Nelson, E. "Solution Rate of Theophylline Salts and Effects  
from Oral Administration." J.Am.Pharm.Assoc. (Sci.  
Ed.) 46(October 1957) : 607-614.
42. Shami, E.G., Dudzinski, J.R., and Lantz, R.J. "Preformula-  
tion." in The Theory and Practice of Industrial  
Pharmacy, p. 19-20 Edited by L.Lachman, H.A. Lieber-  
man, J.L. Kanig. Philadelphia : Lea & Febiger, 1976.
43. Hersey, J.A. "Method Available for the Determination of in  
Vitro Dissolution Rate." Mfg. Chem. & Aerosol News.  
40(February 1969) : 32-35.
44. Levy, G., and Hayes, B.A, "Physicochemical Basis of the  
Buffered Acetylsalicylic Acid Controversy." New  
Eng. J. Med. 262 (May 1960) : 1053-1058.
45. Levy, G. and Sahli, B.A. "Comparison of the Gastrointesti-  
nal Absorption of Aluminum Acetylsalicylate and  
Acetylsalicylic Acid in Man." J.Pharm.Sci. 51  
(January 1962) : 58-62.
46. Pernarowsky M., Woo, W., and Searl, R.O. "Continuous flow  
Apparatus for the Determination of the Dissolution

Characteristics of Tablets and Capsules." ibid.

57 (August 1968) 1419-1421.

47. Gibaldi, M. and Wientraub, H."Quantitative Correlation of Absorption and In Vitro Dissolution Kinetics of Aspirin from Several Dosage Forms." ibid. 59(May 1970) : 725-726.

48. Wood, J.H., Syarto, J.E., and Letterman, H. "Improved Holder for Intrinsic Dissolution Rate Studies." ibid. 64 (July 1965) : 1068.

49. Prakongpan, S., et al. "Dissolution Rate Studies of Cholesterol Monohydrate in Bile Acid-Lecithin Solution Using the Rotating-Disk Method" ibid. 65(May 1976) : 685-689.

APPENDIX



Table 10. The Absorbance of Standard Solution of Paracetamol at 244 nm.

Concentration, mg %	Absorbance
0.25	0.17
0.50	0.32
0.75	0.48
1.00	0.65
1.50	0.97
2.50	1.54

Table 11. Dissolution Data by the Method of Nogami et al. in 50 ml of 0.1 N HCl at 37° from a Disk of 1.5 cm in Diameter Rotating at 200 rpm.

Minutes	Concentration, mg/ml
15	1.94
30	3.73
45	5.32
60	6.82
75	8.17
90	9.52
105	10.63
120	11.74
135	12.85

Table 12. Dissolution Profiles of Paracetamol Compressed at 2,000, 3,000, and 4,000 lb Using Compression Die Assembly.

Minutes	Amount Dissolved, mg		
	2,000 lb	3,000 lb	4,000 lb
5	4.76	5.45	5.36
10	9.75	10.17	10.49
15	14.57	15.12	15.66
20	19.76	19.81	21.03
25	24.41	24.79	25.45
30	28.76	29.19	29.78

Table 13. Dissolution Profiles of Paracetamol from Paracetamol-Lactose  
Mixtures Compressed at 2,000, 3,000, and 4,000 lb.

Minutes	Amount Dissolved . mg.								
	10 % Lactose			20 % Lactose			30 % Lactose		
	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb
5	4.49	4.30	4.37	4.56	4.56	4.50	4.04	4.30	4.43
10	9.02	8.80	9.10	9.10	8.90	9.10	8.83	8.83	9.09
15	12.92	13.03	13.89	13.10	13.56	13.76	12.99	12.83	13.78
20	17.29	17.36	18.58	17.30	17.84	18.49	17.23	17.50	18.49
25	21.76	21.64	23.02	21.67	22.23	22.83	21.64	21.63	22.84
30	25.91	25.96	27.36	25.99	26.14	26.97	25.96	25.97	27.10

Table 14. Dissolution Profiles of Paracetamol from Paracetamol-Microcrystalline Cellulose Mixtures Compressed at 2,000, 3,000, and 4,000 lb.

Minutes	Amount Dissolved, mg								
	10 % Microcrystalline Cellulose			20 % Microcrystalline Cellulose			30 % Microcrystalline Cellulose		
	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb
5	3.76	3.90	3.64	3.31	3.24	3.17	2.98	2.78	2.78
10	7.56	7.24	7.63	6.24	5.91	5.65	5.58	4.87	4.59
15	11.03	10.44	11.10	8.85	7.78	7.58	7.79	6.39	6.19
20	13.98	13.91	14.40	11.25	9.85	9.39	9.65	7.85	7.39
25	17.50	16.98	17.37	13.38	11.78	11.05	11.39	9.12	8.52
30	20.24	19.90	20.64	15.05	13.85	12.39	12.92	10.13	9.53

Table 15. Dissolution Profiles of Paracetamol from Paracetamol-Magnesium Stearate Mixtures Compressed at 2,000, 3,000, and 4,000 lb.

Minutes	Amount Dissolved, mg					
	0.5 % Magnesium Stearate			1 % Magnesium Stearate		
	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb
5	4.56	4.56	4.56	3.64	3.84	4.03
10	8.57	8.77	8.77	7.17	7.44	7.37
15	12.77	12.57	12.77	10.77	10.84	10.44
20	16.77	16.37	16.57	14.17	14.04	13.90
25	20.77	20.37	20.57	17.04	17.30	16.78
30	24.57	23.97	24.37	19.19	20.18	20.11

Table 15. (Continued) Dissolution Profiles of Paracetamol from Paracetamol-Magnesium Stearate Mixtures Compressed at 2,000, 3,000, and 4,000 lb.

Minutes	Amount Dissolved, mg					
	3 % Magnesium Stearate			5 % Magnesium Stearate		
	2,000 1b	3,000 1b	4,000 1b	2,000 1b	3,000 1b	4,000 1b
5	3.30	3.57	3.51	3.24	3.44	3.31
10	6.18	6.38	6.18	5.91	5.98	5.91
15	8.58	8.85	8.65	7.92	8.18	7.98
20	10.98	10.98	10.78	9.92	10.05	9.98
25	13.05	13.25	13.05	11.65	11.79	11.65
30	15.12	15.12	14.92	13.25	13.19	13.25

- Table 16. Dissolution Profiles of Paracetamol from Paracetamol-Talc Mixtures  
Compressed at 2,000, 3,000, and 4,000 lb.

Minutes	Amount Dissolved, mg					
	1 % Talc			3 % Talc		
	2,000 lb)	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb
5	4.56	4.36	4.56	3.97	3.90	4.17
10	8.96	8.76	8.96	8.10	7.97	8.17
15	13.56	13.16	13.56	12.19	11.97	12.10
20	18.16	17.96	17.96	16.07	15.77	16.10
25	22.37	22.17	22.36	20.10	19.70	19.84
30	26.37	25.97	26.17	23.77	23.40	23.44

Table 16. (Continued) Dissolution Profiles of Paracetamol from Paracetamol  
-Talc Mixtures Compressed at 2,000, 3,000, and 4,000 lb.

Minutes	Amount Dissolved, mg					
	5 % Talc			7 % Talc		
	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb
5	3.70	3.84	3.70	3.64	3.64	3.57
10	7.44	7.57	7.37	7.11	7.04	7.24
15	11.10	11.37	11.04	10.70	10.64	10.63
20	14.83	15.04	14.64	14.30	14.04	14.17
25	18.37	18.77	18.17	17.77	17.57	17.18
30	21.77	22.11	21.57	21.17	20.91	20.97

Table 17. Dissolution Profiles of Paracetamol from Paracetamol-Starch Paste  
 (10 % w/w) Mixtures Compressed at 2,000, 3,000, and 4,000 lb.

Minutes	Amount Dissolved, mg					
	1 % Starch*			2 % Starch*		
	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb
5	3.37	3.17	3.57	3.44	3.50	3.57
10	6.77	6.57	6.77	5.98	6.24	6.31
15	10.37	9.97	10.37	8.23	8.46	8.45
20	13.38	12.98	13.18	10.28	10.45	10.38
25	16.57	16.17	16.37	12.38	12.17	12.05
30	19.97	19.18	20.17	14.12	13.98	13.79

\*quantity of dry mass in starch paste (10 % w/w) mixed with drug.

Table 17. (Continued) Dissolution Profiles of Paracetamol from Paracetamol-Starch Paste (10 % w/w) Mixtures Compressed at 2,000, 3,000 and 4,000 lb.

Minutes	Amount Dissolved, mg					
	3 % Starch*			4 % Starch*		
	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb
5	3.50	3.31	3.57	3.17	3.24	3.31
10	5.71	5.71	5.98	5.45	5.58	5.51
15	7.72	7.78	8.05	7.28	7.38	7.52
20	9.58	9.25	9.79	8.99	8.99	8.05
25	11.19	10.97	11.39	10.39	10.52	10.65
30	12.65	12.65	12.66	11.59	11.86	11.92

\* quantity of dry mass in starch paste (10 % w/w) mixed with durg.

Table 18. Dissolution Profiles of Paracetamol from Paracetamol

-Polyvinylpyrrolidone (in Solution) Mixtures Compressed at  
3,000 and 4,000 lb.

Minutes	Amount Dissolved, ml					
	3 % Polyvinylpyrrolidone		4 % Polyvinylpyrrolidone		5 % Polyvinylpyrrolidone	
	3,000 lb	4,000 lb	3,000 lb	4,000 lb	3,000 lb	4,000 lb
5	4.50	4.36	4.56	4.43	4.36	4.36
10	9.16	9.09	9.23	9.03	8.90	9.03
15	13.89	13.83	14.03	13.76	13.63	13.76
20	18.63	18.56	18.63	18.56	18.29	18.49
25	23.09	23.01	23.01	22.96	22.76	22.76
30	27.76	27.56	27.50	27.23	26.97	26.97

Table 19. Dissolution Profiles of Paracetamol from Paracetamol-Polyvinylpyrrolidone (Dry Adding) Compressed at 3,000 and 4,000 lb.

Minutes	Amount Dissolved, mg					
	3 % Polyvinylpyrrolidone		4 % Polyvinylpyrrolidone		5 % Polyvinylpyrrolidone	
	3,000 lb	4,000 lb	3,000 lb	4,000 lb	3,000 lb	4,000 lb
5	4.36	4.56	4.36	4.36	4.56	4.56
10	9.36	9.12	9.16	9.36	9.36	9.16
15	14.16	13.96	13.96	13.96	13.96	14.16
20	18.76	18.76	18.56	18.76	18.56	18.65
25	23.56	22.96	22.96	23.17	23.21	22.96
30	27.76	27.56	27.36	27.56	27.56	27.56

Table 20. Dissolution Profiles of Paracetamol from Granule-Dry Corn Starch  
 Mixtures Compressed at 2,000, 3,000 and 4,000 lb.

Minutes	Amount Dissolved, mg								
	1 % Starch			3 % Starch			5 % Starch		
	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb	2,000 lb	3,000 lb	4,000 lb
5	2.98	2.98	2.98	2.78	2.98	2.78	2.58	2.58	2.58
10	5.18	5.18	4.78	4.78	5.18	5.18	4.78	4.58	4.78
15	6.98	6.98	6.98	6.78	6.99	6.78	6.78	6.58	6.58
20	8.73	8.59	8.39	8.39	8.79	8.73	8.58	8.19	8.58
25	10.19	10.39	10.39	10.18	10.12	10.12	10.19	9.59	10.19
30	11.59	11.39	11.39	11.59	11.59	11.59	11.58	10.99	11.58

Table 21. Dissolution Profile of Paracetamol from Granule-Lubricants  
 Mixture (Combination of Magnesium Stearate and Talc) Compressed  
 at 2,000 and 3,000 lb.

Minutes	Amount Dissolved, mg							
	0.3% Magnesium Stearate + 3% Talc		0.3% Magnesium Stearate + 5% Talc		1% Magnesium Stearate + 3% Talc		1% Magnesium Stearate + 5% Talc	
	2,000 lb	3,000 lb	2,000 lb	3,000 lb	2,000 lb	3,000 lb	2,000 lb	3,000 lb
5	3.37	3.37	2.98	2.98	2.98	3.17	2.78	2.78
10	5.75	5.56	5.16	5.36	5.16	5.56	5.16	5.16
15	7.54	7.54	6.00	7.14	6.94	7.14	6.94	6.94
20	9.13	9.13	8.53	8.53	8.53	8.73	8.13	8.33
25	10.32	10.52	9.92	9.92	9.92	19.12	9.72	9.72
30	11.71	11.90	11.50	11.11	11.31	11.51	10.91	11.11

## VITA

Mr. Poj Kulvanich was born on July 2, 1951. He got his degree in Bachelor of Science in Pharmacy in 1975 from Faculty of Pharmaceutical Sciences, Chulalongkorn University.

He is now working as an instructor in the Department of Manufacturing Pharmacy, Faculty of Pharmaceutical Sciences, Chulalongkorn University.

