

REFERENCES

- Aalbersberg, W. and Singh, Y. 1991. Dammarane triterpenoids from *Dysoxylum richii*. Phytochemistry 30: 921-926.
- Adesogan, E.K. 1973. Coumarins and other components of *Afraegele paniculata*. Phytochemistry 12: 2310-2312.
- Ahad, A.M., Goto, Y., Kiuchi, F., Tsuda, Y., Kondo, K., and Sato, T. 1991. Nematocidal principles in "Oakmoss Absolute" and nematocidal activity of 2,4-dihydroxybenzoate. Chem. Pharm. Bull. 39: 1043-1046.
- Ahmad, V.U. and Alvi, K.A. 1987. Deacetoxybrachycarpone, a trinortriterpenoid from *Cleome brachycarpa*. Phytochemistry 26: 315-316.
- Assimopoulou, A.N. and Papageorgiou, V.P. 2005. GC-MS analysis of penta- and tetracyclic triterpenes from resins of *Pistacia* species. Part II. *Pistacia terebinthus* var. Chia. Biomed. Chromatogr. 19: 586-605.
- Ayimele, G.A., Tane, P., and Connolly, J.D. 2004. Aulacocarpin A and B, Nerlidol and β -sitosterol glucoside from *Aframomum escapurum*. Biochem. Syst. Ecol. 32: 1205-1207.
- Aziz, N.H., Farag, S.E., Mousa, L.A., and Abo-Zaid, M.A. 1998. Comparative antibacterial and antifungal effects of some phenolic compounds. Microbios. 93: 43-54.
- Bacher, M., Hofer, O., Brader, G., Vajrodaya, S., and Greger, H. 1999. Thapsakins: Possible biogenetic intermediates towards insecticidal cyclopenta[b] benzofurans from *Aglaia edulis*. Phytochemistry 52: 253-263.
- Baetas, A.C.S., Arruda, M.S.P., Muller, A.H., and Arruda, A.C. 1999. Coumarins and alkaloids from the stems of *Metrodorea flavaida*. J. Braz. Chem. Soc. 10: 181-183.
- Batey, I.L., Hellyer, R.O., and Pinhey, J.T. 1971. The structure of squamulosone, a new sesquiterpene ketone from *Phebalium squamulosum*. Aust. J. Chem. 24: 2173-2177.
- Baumann, B., Bohnenstengel, F., Siegmund, D., Wajant, H., Weber, C., Herr, I., Debatin, K.-M., Proksch, P., and Wirth, T. 2002. Rocaglamide derivatives are potent inhibitors of NF- κ B activation in T-cells. J. Biol. Chem. 277: 44791-44800.

- Benosman, A., Richomme, P., Roussakis, C., Sevenet, T., Hadi, A.H.A., and Bruneton, J. 2000. Effects of triterpenes from the stem bark of *Dysoxylum cauliflorum* on a non-small-cell bronchopulmonary carcinoma cell line (NSCLC-N[6]). *Anticancer Res.* 20: 1855-1859.
- Benosman, A.B., Richomme, P., Sevenet, T., Hadi, A.H.A., and Bruneton, J. 1994. Secotirucallane triterpenes from the stem bark of *Aglaia leucophylla*. *Phytochemistry* 37: 1143-1145.
- Bhattacharyya, N.K., Sarmah, G.K., Goswami, B.N., Borthakur, N., and Kataky, J.C.S. 2003. New protolimonoids and limonoids: Part V-Synthesis of new heterocyclic compounds from naturally occurring limonoids. *Indian J. Heterocycl. Chem.* 12: 297-300.
- Bhattacharyya, N.K., Sarmah, G.K., Goswami, B.N., Borthakur, N., and Kataky, J.C.S. 2004. New protolimonoids and limonoids. Part-IV. Transformation of naturally occurring 6α -acetoxyazadirone to new limonoids. *J. Indian Chem. Soc.* 81: 433-435.
- Binutu, O.A. 1997. Phytochemical and antimicrobial studies on *Crescentia cujete*. *Fitoterapia* 68: 184-185.
- Boar, R.B. and Damps, K. 1977. Triterpenoids of *Aglaia odorata*. Configuration of trisubstituted epoxides. *J. Chem. Soc. Perkin Trans. I* 510-512.
- Bohlmann, F., Grenz, M., Jakupovic, J., King, R.M., and Robinson, H. 1983. Four heliangolides and other sesquiterpenes from *Brasilia sickii*. *Phytochemistry* 22: 1213-1218.
- Bohlmann, F., Zdero, C., King, R.M., and Robinson, H. 1984. New aromadendrane derivatives from *Wyethia arizonica*. *Planta Med.* 50: 195-196.
- Bohlmann, F., Singh, P., and Jakupovic, J. 1982. Sesquiterpenes and a dimeric spiroketone from *Cineraria fruticulorum*. *Phytochemistry* 21: 2531-2535.
- Bordoloi, M., Saikia, B., Mathur, R.K., and Goswami, B.N. 1993. A meliacin from *Chisocheton paniculatus*. *Phytochemistry* 34: 583-584.
- Bouic, P.J.D., Etsebeth, S., Liebenberg, R.W., Albrecht, C.F., Pegel, K., and Jaarsveld, P.P.V. 1996. Beta-sitosterol and beta-sitosterol glucoside stimulate human peripheral blood lymphocyte proliferation: Implications for their use as an immunomodulatory vitamin combination. *Int. Immunopharmacol.* 18: 693-700.

- Brader, G., Vajrodaya, S., Greger, H., Bacher, M., Kalchhauser, H., and Hofer, O. 1998. Bisamides, lignans, triterpenes, and insecticidal cyclopenta[b] benzofurans from *Aglaia* species. *J. Nat. Prod.* 61: 1482-1490.
- Bringmann, G., Muhlbacher, J., Messer, K., Dreyer, M., Ebel, R., Nugroho, B.W., Wray, V., and Proksch, P. 2003. Cyclorocaglamide, the first bridged cyclopentatetrahydrobenzofuran, and a related “open chain” rocaglamide derivative from *Aglaia oligophylla*. *J. Nat. Prod.* 66: 80-85.
- Brochini, C.B., and Roque, N.F. 2000. Two new cneorubin related diterpenes from the leaves of *Guarea guidonia* (Meliaceae). *J. Braz. Chem. Soc.* 11: 361-364.
- Cabrera, G.M., Oberti, J., Gros, E.G., and Seldes, A. 1993. Aromadendrane sesquiterpene from *Mandevilla pentlandiana*. *Phytochemistry* 34: 1642-1643.
- Camacho, M.D.R., Phillipson, J.D., Croft, S.L., Kirby, G.C., Warhurst, D.C., and Solis, P.N. 2001. Terpenoids from *Guarea rhophalocarpa*. *Phytochemistry* 56: 203-210.
- Cascon, S.C. and Brown Jr., K.S. 1972. Biogenetically significant triterpenes in a species of Meliaceae: *Cabralea polytricha*. *Tetrahedron* 28: 315-323.
- Cassady, J.M., Ojima, N., Chang, C., and McLaughlin, J.L. 1979. An investigation of the antitumor activity of *Micromelum integerrimum* (Rutaceae). *J. Nat. Prod.* 42: 274-278.
- Castillo, J.B.D., Luis, F.R., and Secundino, M. 1984. Angustifolin, a coumarin from *Ruta angustifolia*. *Phytochemistry* 23: 2095-2096.
- Chadir., Hiort, J., Nugroho, B.W., Bohnenstengel, F.I., Wray, V., Witte, L., Hung, P.D., Kiet, L.C., Sumaryono, W., and Proksch, P. 1999. New insecticidal rocaglamide derivatives from flowers of *Aglaia duperreana* (Meliaceae). *Phytochemistry* 52: 837-842.
- Chadir., Lin, W.H., Ebel, R., Edrada, R., Wray, V., Nimtz, M., Sumaryono, W., and Proksch, P. 2001. Rocaglamide, glycosides, and putrescine bisamides from *Aglaia dasyclada*. *J. Nat. Prod.* 64: 1216-1220.
- Chang, H.M. and But, P.P.H. 1987. *Pharmacology and applications of Chinese materia medica*. World Scientific, Singapore, pp. 867-871.
- Chang, C.-T., Doong, S.-L., Tsai, I.-L., and Chen, I.-S. 1997. Coumarins and anti-HBV constituents from *Zanthoxylum schinifolium*. *Phytochemistry* 45: 1419-1422.

- Chatterjee, A., Nayak, L., Das, B., Patra, A., Dhara, K.P., Mukherjee, K., Banerji, J., and Shoolery, J.N. 1989. Chemistry of phytoconstituents of *Chisocheton paniculatus* Hiern (Meliaceae). *Indian J. Chem., Sect. B: Org.* 28B: 231-136.
- Chumkaew, P., Kato, S., and Chantrapromma, K. 2006. Potent cytotoxic rocaglamide derivatives from the fruits of *Amoora cucullata*. *Chem. Pharm. Bull.* 54: 1344-1346.
- Collins, D.O., Buchanan, G.O., Reynolds, W.F., and Reese, P.B. 2001. Biotransformation of squamulosone by *Curvularia lunata* ATCC 12017. *Phytochemistry* 57: 377-383.
- Collins, L.A. and Franzblau, S.G. 1997. Microplate Alamar Blue assay versus BACTEC 460 system for high-throughput screening of compounds against *Mycobacterium tuberculosis* and *Mycobacterium avium*. *Antimicrob. Agents Chemother.* 41: 1004-1009.
- Connolly, J.D., Labbe, C., Rycroft, D.S., and Taylor, D.A.H. 1979. Tetraneortriterpenoids and related compounds. Part 22. New apotirucallol derivatives and tetraneortriterpenoids from the wood and seeds of *Chisocheton paniculatus* Hiern (Meliaceae). *J. Chem. Soc., Perkin Trans. I* 1972-1999.
- Cui, B., Chai, H., Santisuk, T., Reutrakul, V., Farnsworth, N.R., Cordell, G.A., Pezzuto, J.M., and Kinghorn, A.D. 1997. Novel cytotoxic 1*H*-cyclopenta[*b*]benzofuran lignans from *Aglaiella elliptica*. *Tetrahedron* 53:17625-17632.
- Daniewski, W.M., Añczewski, W., Gumulka, M., Danikiewicz, W., Jacobsson, U., and Norin, T. 1996. Sesquiterpenoid constituents of *Entandrophragma cylindricum*. *Phytochemistry* 43: 811-814.
- De Campos Braga, P.A., Soares, M.S., Da Silva, M.F.D.G.F., Vieira, P.C., Fernandes, J.B., and Pinheiro, A.L. 2006. Dammarane triterpenes from *Cabralea canjerana* (Vell.) Mart. (Meliaceae): Their chemosystematic significance. *Biochem. Syst. Ecol.* 34: 282-290.
- Deepak, M., Dipankar, G., Prashanth, D., Asha, M.K., Amit, A., and Venkataraman, B.V. 2002. Tribulosin and β -sitosterol-D-glucoside, the anthelmintic principles of *Tribulus terrestris*. *Phytomedicine* 9: 753-756.

- De Siqueira, J.M., Muller, L., Carollo, C.A., Garcez, W.S., Boaventura, M.A.D., and Nascimento, E.A. 2003. Aromadendrane sesquiterpenoids from the essential oil of leaves of *Duguetia glabriuscula* – Annonaceae. *J. Chil. Chem. Soc.* 48: 89-97.
- Desjardins, R.E., Canfield, C.J., Haynes, J.D., and Chulay, J.D. 1979. Quantitative assessment of antimalarial activity in vitro by a semiautomated micro dilution technique. *Antimicrob. Agents Chemother.* 16: 710-718.
- De Tommasi, N., Pizza, C., Conti, C., Orsi, N., and Stein, M.L. 1990. Structure and in vitro antiviral activity of sesquiterpene glycosides from *Calendula arvensis*. *J. Nat. Prod.* 53: 830-835.
- Dolejs, L., Motl, O., Soucek, M., Herout, V., and Sorm, F. 1960. Terpenes. CVIII. Epimeric aromadendrenes, stereo-isomerism of ledol, viridiflorol, and globulol. *Collect. Czech. Chem. Commun.* 25: 1483-1491.
- Dreyer, M., Nugroho, B.W., Bohnenstengel, F.I., Ebel, R., Wray, V., Witte, L., Bringmann, G., Muhlbacher, J., Herold, M., Hung, P.D., Kiet, L.C., and Proksch, P. New insecticidal rotaglamide derivatives and related compounds from *Aglaia oligophylla*. *J. Nat. Prod.* 64: 415-420.
- Dumontet, V., Thoison, O., Omobuwajo, O.R., Martin, M.-T., Perromat, G., Chiaroni, A., Riche, C., Pais, M., and Sevenet, T. 1996. New nitrogenous and aromatic derivatives from *Aglaia argentea* and *A. forbesii*. *Tetrahedron* 52: 6931-6942.
- Engelmeier, D., Hadacek, F., Pacher, T., Vajrodaya, S., and Greger, H. 2000. Cyclopenta[*b*]benzofurans from *Aglaia* species with pronounced antifungal activity against rice blast fungus (*Pyricularia grisea*). *J. Agric. Food Chem.* 48:
- Fang, Y., Li, Z., and Watanabe, Y. 2003. Pharmacokinetics of novel anti-asthmatic, scoparone, in the rabbit serum assessed by a simple HPLC method. *J. Ethnopharmacol.* 86: 127-130.
- Feliciano, A.S., Medarde, M., Gordaliza, M., Olmo, E.D., and Corral, J.M.M.D. 1989. Sesquiterpenoids and phenolics of *Pulicaria paludosa*. *Phytochemistry* 28: 2717-2721.
- Fernandez, M.A., Garcia, M.D., and Saenz, M.T. 1996. Antibacterial activity of the phenolic acid fractions of *Scrophularia frutescens* and *Scrophularia sambucifolia*. *J. Ethnopharmacol.* 53: 11-14.

- Friedel, H.D., and Matusch, R. 1987. New aromadendrane derivatives from Tolu Balsum. *Helv. Chim. Acta* 70: 1753-1759.
- Fuzzati, N., Dyatmiko, W., Rahman, A., Achmad, F., and Hostettmann, K. 1996. Triterpenoids, lignans and a benzofuran derivative from the bark of *Aglaia elaeagnoidea*. *Phytochemistry* 42: 1395-1398.
- Goldsby, G. and Burke, B.A. 1987. Sesquiterpene lactones and a sesquiterpene diol from Jamaican *Ambrosia peruviana*. *Phytochemistry* 26: 1059-1063.
- Govindachari, T.R., Suresh, G., and Kumari, G.N. 1994. Triterpenoids from *Dysoxylum malabaricum*. *Phytochemistry* 37: 1127-1129.
- Graham, B.A., Jefferies, P.R., Melrose, G.J.H., Thieberg, K.J.L. and White, D.E. 1960. (+)-Aromadendrene as chiral starting material for the synthesis of fragrances and pheromones. *Aust. J. Chem.* 13: 372-379.
- Greger, H., Pacher, T., Brem, B., Bacher, M., and Hofer, O. 2001. Insecticidal flavaglines and other compounds from Fijian *Aglaia* species. *Phytochemistry* 57: 57-64.
- Gunning, P.J., Jeffs, L.B., Isman, M.B., and Towers, G.H.N. 1994. Two limonoids from *Chisocheton microcarpus*. *Phytochemistry* 36: 1245-1248.
- Gupta, A.S. and Dev, S. 1971. Triterpenoids from the oleoresin of *Dipterocarpus pilosus*: Hollongdione and dipterocarpolic acid. *Tetrahedron* 27: 823-834.
- Haque, M.E., Uddin, M.N., and Jalal, M.S. 1995. Dammarane triterpenoids from the stem bark of *Amoora cucullata* (Meliaceae). *Bangladesh J. Bot.* 24: 165-171.
- Harraz, F.M., Ulubelen, A., Oksuz, S., and Tan, N. 1995. Dammarane triterpenes from *Cleome amblyocarpa*. *Phytochemistry* 39: 175-178.
- Haque, M.E., Uddin, M.N., and Jalal, M.S. 1996. Cabralealactone and isofouquierone from the stem bark of *Amoora cucullata* L. (Meliaceae). *Bangladesh J. Bot.* 25: 177-181.
- Hausott, B., Greger, H., and Marian, B. 2004. Flavaglines: A group of efficient growth inhibitors block cell cycle progression and induce apoptosis in colorectal cancer cells. *Int. J. Cancer* 109: 933-940.
- Herz, W., Bhat, S.V., and Santhanam, P.S. 1970. Coumarins of *Artemisia dracunculoides* and 3',6'-dimethoxy-4',5,7-trihydroxyflavone in *A. arctica*. *Phytochemistry* 9: 891-894.

- Hiort, J., Chaidir., Bohnenstengel, F.I., Nugroho, B.W., Schneider, C., Wray, V., Witte, L., Hung, P.D., Kiet, L.C., and Proksch, P. 1999. New insecticidal rocaglamide derivatives from the roots of *Aglaia duperreana*. *J. Nat. Prod.* 62: 1632-1635.
- Hisham, A., Ajitha, B.M.D., Fujimoto, Y., Hara, N., and Shimada, H. 1996. Complete ¹H and ¹³C NMR spectral assignment of cabraleadiol, dammarane triterpene from *Dysoxylum malabaricum* Bedd. *Magn. Reson. Chem.* 34: 146-150.
- Hoult, J.R.S. and Paya, M. 1996. Pharmacological and biochemical actions of simple coumarins: Natural products with therapeutic potential. *Gen. Pharmacol.* 27: 713-722.
- Hsiao, J.-J. and Chiang, H.-C. 1995. Phenylpropanoids from *Aralia bipinnata*. *Phytochemistry* 39: 825-827.
- Huang, R., Harrison, L.J., and Sim, K.-Y. 1999. A triterpenoid with a novel *abeo*-dammarane skeleton from *Dysoxylum cauliflorum*. *Tetrahedron Lett.* 40: 1607-1610.
- Hwang, B.Y., Su, B.-N., Chai, H., Mi, Q., Kardono, L.B.S., Afriastini, J.J., Riswan, S., Santarsiiero, B.D., Mesecar, A.D., Wild, R., Fairchild, C.R., Vite, G.D., Rose, W.C., Farnsworth, N.R., Cordell, G.A., Pezzuto, J.M., Swanson, S.M., and Kinghorn, A.D. 2004. Silvestrol and episilvestrol, potential anticancer rocaglate derivatives from *Aglaia silvestris*. *J. Org. Chem.* 69: 3350-3358.
- Iinuma, M., Ohyama, M., Tanaka, T., Mizuno, M., and Hong, S.-K. 1993. Five flavonoid compounds from *Echinosophora koreensis*. *Phytochemistry* 33: 1241-1245.
- Inada, A., Shono, K., Murata, H., Inatomi, Y., Darnaedi, D., and Nakanishi, T. 2000. Three putrescine bisamides from the leaves of *Aglaia grandis*. *Phytochemistry* 53: 1091-1095.
- Inada, A., Somekawa, M., Murata, H., Nakanishi, T., Tokuda, H., Nishino, H., Iwashima, A., Darnaedi, D., and Jurata, J. 1993. Phytochemical studies on meliaceous plants. VIII. Structures and inhibitory effects on Epstein-Barr virus activation of triterpenoids from leaves of *Chisocheton macrophyllus* King. *Chem. Pharm. Bull.* 41: 617-619.
- Ishibashi, F., Satasook, C., Isman, M.B., and Towers, G.H.N. 1993. Insecticidal 1H-cyclopentatetrahydro[b]benzofurans from *Aglaia odorata*. *Phytochemistry* 32: 307-310.

- Jakupovic, J., Schuster, A., Bohlmann, F., King, R.M., and Lander N.S. 1988. Sesquiterpene lactones from *Gnephosis* species. *Phytochemistry* 27: 3181-3185.
- Janprassert, J., Satasook, C., Sukumalanand, P., Champagne, D.E., Isman, M.B., Wiriyachitra, P., and Towers, G.H. 1993. Rocaglamide, a natural benzofuran insecticide from *Aglaia odorata*. *Phytochemistry* 32: 67-69.
- Jang, S.I., Kim, Y.J., Lee, W.Y., Kwak, K.C., Baek, S.H., Kwak, G.B., Yun, Y.G., Kwon, T.O., Chung, H.T., and Chai, K.Y. 2005. Scoparone from *Artemisia capillaries* inhibits the release of inflammatory mediators in RAW 264.7 cells upon stimulation cells by interferon-gamma plus LPS. *Arch. Pharm. Res.* 28: 203-208.
- Jimenez, C., Marcos, M., Villaverde, M.C., Riguera, R., Castedo, L., and Stermitz, F. 1989. A chromone from *Zanthoxylum* species. *Phytochemistry* 28: 1992-1993.
- Kang, T.H., Pae, H.O., Jeong, S.J., Yoo, J.C., Choi, B.M., Jun, C.D., Chung, H.T., Miyanoto, T., Higuchi, R., and Kim, Y.C. 1999. Scopoletin: An inducible nitric oxide synthesis inhibitory active constituent from *Artemisia feddei*. *Planta Med.* 65: 400-403.
- Kang, S.Y., Sung, S.H., Park, J.H., and Kim, Y.C. 1998. Hepatoprotective activity of scopoletin, a constituent of *Solanum lyratum*. *Arch. Pharm. Res.* 21: 718-722.
- Kim, S., Kinghorn, A.D., and Swanson, S.M. 2006. Phytochemical investigation of *Aglaia edulis* and mechanistic studies of silvestrol-induced apoptosis. *Anticancer Agents Med. Chem.* 6: 319-345.
- King, M.L., Chiang, C.-C., Ling, H.-C., Fujita, E., Ochiai, M., and McPhail, A.T. 1982. X-ray crystal structure of rocamamide, a novel antileukemic 1*H*-cyclopenta[*b*]benzofuran from *Aglaia elliptifolia*. *J. Chem. Soc., Chem. Commun.* 1150-1151.
- Kingston, D.G., Rao, M.M., Spittler, T.D., Pettersen, R.C., and Cullen, D.L. 1975. Sesquiterpenes from *Flourensia cernua*. *Phytochemistry* 14: 2033-2037.
- Ko, F.-N., Wu, T.-S., Liou, M.-J., Huang, T.-F., and Teng, C.-M. 1992. PAF antagonism in vitro and in vivo by aglafoline from *Aglaia elliptifolia*. *Eur. J. Pharmacol.* 218: 129-135.
- Kojima, H., Sato, N., Hatano, A., and Ogura, H. 1990. Sterol glucosides from *Prunella vulgaris*. *Phytochemistry* 29: 2351-2355.

- Kokpol, U., Venaskulchai, B., Simpson, J., and Weavers, R.T. 1994. Isolation and X-ray structure determination of a novel pyrimidinone from *Aglaia odorata*. J. Chem. Soc., Chem. Commun. 773-774.
- Koul, S.K., Taneja, S.C., Malhotra, S., and Dhar, K.L. 1993. Phenylpropanoids and (-)-ledol from two *Piper* species. Phytochemistry 32: 478-480.
- Krishnappa, S. and Dev, S. 1973. Sesquiterpenes from *Lansium anamalayanum*. Phytochemistry 12: 823-825.
- Krishnaswamy, N.R. and Prasanna, S. 1975. α - and β -Amyrin esters and sitosterol glucoside from *Spilanthes acmella*. Phytochemistry 14: 1666-1667.
- Ku, Y.-L., Chen, C.-H., and Lee, S.-S. 2006. Chemical constituents from *Phoebe minutiflora* II. Nat. Prod. Res. 20: 1199-1206.
- Kumar, V., Bulumulla, H.N.K., Wimalasiri, W.R., and Reisch, J. 1994. Coumarins and an indole alkaloid from *Pamburus missionnis*. Phytochemistry 36: 879-881.
- Lee, S.K., Cui, B., Mehta, R.R., Kinghorn, A.D., and Pezzuto, J.M. 1998. Cytostatic mechanism and antitumor potential of novel 1*H*-cyclopenta[*b*]benzofuran lignans isolated from *Aglaia elliptica*. Chem. Biol. Interact. 115: 215-228.
- Lee, S.-S., Chang, S.-M., and Chen, C.-H. 2001. Chemical constituents from *Alseodaphne andersonii*. J. Nat. Prod. 64: 1548-1551.
- Li, X.-C., Elsohly, H.N., and Clark, A.M. 2000. 7-Caffeoylsedoheptulose from *Nyssa sylvatica*. Phytochemistry 53: 1033-1037.
- Limberger, R.P., Scopel, M., Sobral, M., and Henriques, A.M. 2007. Comparative analysis of volatiles from *Drimys brasiliensis* Miers and *D. angustifolia* Miers (Winteraceae) from Southern Brazil. Biochem. Syst. Ecol. 35: 130-137.
- Liu, H.R., Li, Z., Wang, X.H., Han, X.S., Teng, Z., and Sun, Z. 2002. Effect of scoparone on $[Ca^{2+}]_i$ of the isolated tracheal smooth muscle cells of guinea-pig. J. Chin. Med. Univ. 31: 249-251.
- Liu, X.-L., Zhang, L., Fu, X.-L., Chen, K., and Qian, B.-C. 2001. Effect of scopoletin on PC 3 cell proliferation and apoptosis. Acta Pharmacol. Sin. 22: 929-933.
- Luo, X.-D., Wu, S.-H., Ma, Y.-B., and Wu, D.-G. 2000. Dammarane triterpenoids from *Amoora yunnanensis*. Heterocycles 53: 2795-2802.

- Ma, C.H., Ke, W., Sun, Z.L., Peng, J.Y., Li, Z.X., Zhou, X., Fan, G.R., and Huang, C.G. 2006. Large-scale isolation and purification of scoparone from *Herba artemisiae scopariae* by high-speed counter-current chromatography. *Chromatographia* 64: 83-87.
- Mabberly, D.J. 1979. The species of *Chisocheton* (Meliaceae). *Bull. Br. Mus. Nat. Hist.* 6: 301-386.
- Majinda, R.R.T., Gray, A.I., Waigh, R.D., and Waterman, P.G. 1995. A *sec*-olean-18-ene triterpene acid from *Vahlia capensis*. *Phytochemistry* 38: 461-463.
- Matos, M.F.C., Leite, L.I.S.P., Brustolim, D., De Siqueira, J.M., Carollo, C.A., Hellmann, A.R., Pereira, N.F.G., and Da silva, D.B. 2006. Antineoplastic activity of selected constituents of *Duguetia glabriuscula*. *Fitoterapia* 77: 227-229.
- Miyazawa, M., Uemura, T., and Kameoka, H. 1995. Biotransformation of sesquiterpenoids, (+)-aromadendrane and (-)-alloaromadendrane by *Glomerella cingulata*. *Phytochemistry* 40: 793-796.
- Mizushina, Y., Nakanishi, R., Kuriyama, I., Kamiya, K., Satake, T., Shimazaki, N., Koiwai, O., Uchiyama, Y., Yonezawa, Y., Takemura, M., Sakaguchi, K., and Yoshida, H. 2006. β -Sitosterol-3-O- β -D-glucoside: A eukaryotic DNA polymerase λ inhibitor. *J. Ster. Biochem. Mol. Biol.* 99: 100-107.
- Mohamad, K., Sevenet, T., Dumontet, V., Pais, M., Tri, M.V., Hadi, H., Awang, K., and Martin, M.-T. 1999. Dammarane triterpenes and pregnane steroids from *Aglaia lawii* and *A. tomentosa*. *Phytochemistry* 51: 1031-1037.
- Molleyres, L.-P., Rindlisbacher, A., Winkler, T., and Kumar, V. 1999. Insecticidal natural products: New rocaglamide derivatives from *Aglaia roxburghiana*. *Pest. Sci.* 55: 486-503.
- Moreira, I.C., Lago, J.H.G., Young, M.C.M., and Roque, N.F. 2003. Antifungal aromadendrane sesquiterpenoids from the leaves of *Xylopia brasiliensis*. *J. Braz. Chem. Soc.* 14: 828-831.
- Mulholland, D.A., Nair, J.J., and Taylor, D.A.H. 1994. Dammaranes from *Astrotrichilia asterotricha*. *Phytochemistry* 35: 542-544.
- Mulholland, D.A. and Naidoo, N. 2000. Dammarane triterpenoids from *Dysoxylum muellerii*. *Biochem. Syst. Ecol.* 28: 295-297.
- Mulholland, D.A. and Naidoo, N. 1998. A revision of the structure of ferrugin from *Aglaia ferruginea*. *Phytochemistry* 47: 1163.

- Mulholland, D.A., Naidoo, D., Randrianarivelojosia, M., Cheplogoi, P.K., and Coombes, P.H. 2003. Secondary metabolites from *Cedrelopsis grevei* (Ptaeroxylaceae). *Phytochemistry* 64: 631-635.
- Muhtadi, Hakim, E.H., Juliawaty, L.D., Syah, Y.M., Achmad, S.A., Latip, J., and Ghisalberti, E.L. 2006. Cytotoxic resveratrol oligomers from the tree bark of *Dipterocarpus hasseltii*. *Fitoterapia* 77: 550-555.
- Muellner, A.N., Samuel, R., Chase, M.W., Pannell, C.M., and Greger, H. 2005. *Aglaia* (Meliaceae): An evaluation of taxonomic concepts based on DNA data and secondary metabolites. *Am. J. Bot.* 92: 534-543.
- Nakamura, N., Kojima, S., Lim, Y.A., Meselhy, M.R., Hattori, M., Gupta, M.P., and Correa, M. 1997. Dammarane-type triterpenes from *Cordia spinescens*. *Phytochemistry* 46: 1139-1141.
- Niyonzima, G., Pieters, L., Balde, A.M., Claeys, M., Laekeman, G.M., and Vlietinck, A.J. 1991. Isolation of 6-O-caffeyl catalpol and some other compounds from *Spathodea campanulata*. *Planta Med.* 57: A85-A87.
- Nugroho, B.W., Gussregen, B., Wray, V., Witte, L., Bringmann, G., and Proksch, P. 1997b. Insecticidal rocamamide derivatives from *Aglaia elliptica* and *A. harmsiana*. *Phytochemistry* 45: 1579-1585.
- Nugroho, B.W., Edrada, R.A., Wray, V., Witte, L., Bringmann, G., Gehling, M., and Proksch, P. 1999. An insecticidal rocamamide derivatives and related compounds from *Aglaia odorata* (Meliaceae). *Phytochemistry* 51: 367-376.
- Nugroho, B.W., Edrada, R.A., Gussregen, B., Wray, V., Witte, L., and Proksch, P. 1997a. Insecticidal rocamamide derivatives from *Aglaia duppereana*. *Phytochemistry* 44: 1455-1461.
- Ohse, T., Ohba, S., Yamamoto, T., Koyano, T., and Umezawa, K. 1996. Cyclopentabenzofuran lignan protein synthesis inhibitors from *Aglaia odorata*. *J. Nat. Prod.* 59: 650-652.
- Ojewoleand, J.A.O. and Adesina, S.K. 1983. Cardiovascular and neuromuscular action of scopoletin from fruit of *Tetrapleura tetraptera*. *Planta Med.* 49: 99-102.
- Ojewole, J.A.O. 2002. Hypoglycaemic effect of *Clausena anisata* (Willd) Hook methanolic root extract in rats. *J. Ethnopharmacol.* 81: 231-237.
- Padilla, S.P. and Soliven, F.A. 1933. Chemical analysis for possible sources of oils of forty-five species of oil-bearing seeds. *Philipp. Agric.* 22: 408-415.

- Pannell, C.M. 1992. A taxonomic monograph of the genus *Aglaia* Lour. (Meliaceae). Kew Bulletin additional series XVI. London: Royal Botanic Gardens, Kew, pp. 76-84.
- Patra, A., Mandal, T.K., Majumdar, A., and Mandal, K.K. 1998. Chemical transformations of 6α -acetoxyazadirone and two-dimensional NMR studies on tetranol triterpenoids. J. Indian Chem. Soc. 75: 804-809.
- Pennington, T.D. and Styles, B.T. 1975. A generic monograph of the Meliaceae. Blumea 22: 419-540.
- Pistelli, L., Bertoli, A., Bilia, A.R., and Morelli, I. 1996. Minor constituents from *Bupleurum fruticosum* roots. Phytochemistry 41: 1579-1582.
- Pizza, C. and De Tommasi, N. 1988. Sesquiterpene glycosides based on the alloaromadendrane skeleton from *Calendula arvensis*. Phytochemistry 27: 205-208.
- Plumb, J.A., Milroy, A., and Kaye, S.B. 1989. Effect of the pH dependence of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide-formazan absorption on chemosensitivity determinated by a novel tetrazolium-based assay. Cancer Res. 49: 4435-4440.
- Proksch, P., Edrada, R., Ebel, R., Bohnenstengel, F.I., and Nugroho, B.W. 2001. Chemistry and biological activity of rocaglamide derivatives and related compounds in *Aglaia* species (Meliaceae). Curr. Org. Chem. 5: 923-938.
- Qiu, S.-X., Hung, N.V., Xuan, L.T., Gu, J.-Q., Lobkovsky, E., Khanh, T.C., Soejarto, D.D., Clardy, J., Pezzuto, J.M., Dong, Y., Tri, M.V., Huong, L.M., and Fong, H.H.S. 2001. A pregnane steroid from *Aglaia lawii* and structure confirmation of cabraleadiol monoacetate by x-ray crystallography. Phytochemistry 56: 775-780.
- Rao, M.M., Meshulam, H., Zelnik, R., and Lavie, D. 1975. *Cabralea eichleriana* DC. (Meliaceae)-I Structure and stereochemistry of wood extractives. Tetrahedron 31: 333-339.
- Rashid, M.A., Gray, A.I., Waterman, P.G., and Armstrong, J.A. 1995. Aromadendrane sesquiterpenes from *Phebalium*, *Drummondita*, and *Eriostemon* species. J. Nat. Prod. 58: 618-621.
- Razdan, T.K., Qadri, B., Harkar, S., and Waight, E.S. 1987. Chromones and coumarins from *Skimmia laureola*. Phytochemistry 26: 2063-2069.

- Rivero-Cruz, J.F., Chai, H.-B., Kardono, L.B.S., Setyowati, F.M., Afriatini, J.J., Riswan, S., Farnsworth, N.R., Cordell, G.A., Pezzuto, J.M., Swanson, S.M., and Kinghorn, A.D. 2004. Cytotoxic constituents of the twigs and leaves of *Aglaia rubiginosa*. *J. Nat. Prod.* 67: 343-347.
- Rizk, A.M. and Rimpler, H. 1972. Isolation of daphnoretin and β -sitosterol- β -D-glucoside from *Thymelea hirsuta*. *Phytochemistry* 11: 473-475.
- Robeson, D.J., Ingham, J.L., and Harborne, J.B. 1980. Identification of two chromone phytoalexins in the sweet pea, *Lathyrus odoratus*. *Phytochemistry* 19: 2171-2173.
- Roux, D., Martin, M.-T., Adeline, M.-T., Sevenet, T., Hadi, A.H.A., and Pais, M. 1998. Foveolins A and B, dammarane triterpenes from *Aglaia foveolata*. *Phytochemistry* 49: 1745-1748.
- Saikia, B., Kataky, J.C.S., Mathur, R.K., and Baruah, J.N. 1978. New meliacins from the fruits of *Chisocheton paniculatus* Hiern. *Indian J. Chem. B. Org.* 16B: 1042-1044.
- Sakushima, A., Coskun, M., and Maoka, T. 1995. Hydroxybenzoic acids from *Boreava orientalis*. *Phytochemistry* 40: 257-261.
- Salim, A.A., Pawlus, A.D., Chai, H.-B., Farnsworth, N.R., Kinghorn, A.D., and Carcache-Blanco, E.J. 2007. Ponapensin, a cyclopenta[bc]benzopyran with potent NF- κ B inhibitory activity from *Aglaia ponapensis*. *Bioorg. Med. Chem. Lett.* 17: 109-112.
- Sarker, S.D., Armstrong, J.A., Gray, A.I., and Waterman, P.G. 1994. Sesquiterpenyl coumarins and geranyl benzaldehyde derivatives from the aerial parts of *Eriostemon myoporoides*. *Phytochemistry* 37: 1287-1294.
- Sarmah, G.K., Bhattacharyya, N.K., Goswami, B.N., Borthakur, N., Baruah, P., and Kata, J.C.S. 2003a. Studies of antifungal activity of naturally occurring paniculatin and their derivatives. *Chemistry* 1: 42-46.
- Sarmah, G.K., Bhattacharyya, N.K., Goswami, B.N., Borthakur, N., Baruah, P., and Kata, J.C.S. 2003b. Derivatives of paniculatin and their antifungal activity. *J. Indian Chem. Soc.* 80: 1163-1168.
- Schneider, C., Bohnenstengel, F.I., Nugroho, B.W., Wray, V., Witte, L., Hung, P.D., Kiet, L.C., and Proksch, P. 2000. Insecticidal rocaglamide derivatives from *Aglaia spectabilis* (Meliaceae). *Phytochemistry* 54: 731-736.

- Shaw, C.Y., Chen, C.H., Hsu, C.C., Chen, C.C., and Tsai, Y.C. 2003. Antioxidant properties of scopoletin isolated from *Sinomonium acutum*. *Phytother. Res.* 17: 823-825.
- Shiengthong, D., Verasarn, A., Nanonggai-Suanrath, P., and Warnhoff, E.W. 1965. Constituents of Thai medicinal plants-I Aglaiol. *Tetrahedron* 21: 917-924.
- Shiengthong, D., Kokpol, U., and Karntiang, P. 1974. Triterpenoid constituents of Thai medicinal plant-II Isomeric aglaitriols and aglaiondiol. *Tetrahedron* 30: 2211-2215.
- Shoeb, A., Kapil, R.S., and Popli, S.P. 1973. Coumarins and alkaloids of *Aegle marmelos*. *Phytochemistry* 12: 2071-2072.
- Sibanda, S., Ndengu, B., Multari, G., Pompi, V., and Galeffi, C. 1989. A coumarin glucoside from *Xeromphis obovata*. *Phytochemistry* 28: 1550-1552.
- Singh, Y. and Aalbersberg, W. 1992. Dammarane triterpenoids from *Dysoxylum richii*. *Phytochemistry* 31: 4033-4035.
- Skehan, P., Storeng, R., Scudiero, D., Monks, A., McMahon, J., Vistica, D., Warren, J.T., Bokesch, H., Kenny, S., and Boyd, M.R. 1990. New colorimetric cytotoxicity assay for anticancer-drug screening. *J. Natl. Cancer. Inst.* 82: 1107-1112.
- Smitinand, T. 2001. *Thai plants names revised Edition*. Bangkok: The Forest Herbarium, Royal Forest Department.
- Staerk, D., Skole, B., Jorgensen, F.S., Budnik, B.A., Ekpe, P., and Jaroszewski, J.W. 2004. Isolation of a library of aromadendranes from *Landolphia dulcis* and its characterization using the Volsurf approach. *J. Nat. Prod.* 67: 799-805.
- Steck, W. and Mazurek, M. 1972. Identification of natural coumarins by NMR spectroscopy. *J. Nat. Prod.* 35: 418-438.
- Stefanovic, M., Krstic, L., and Mladenovic, S. 1973. Extractives of *Artemisia scoparia*. *Phytochemistry* 12: 2996-2997.
- Su, B.-N., Chai, H., Mi, Q., Riswan, S., Kardono, L.B.S., Afriastini, J.J., Santarsiero, B.D., Mesecar, A.D., Farnworth, N.R., Cordell, G.A., Swanson, S.M., and Kinghorn A.D. 2006. Activity-guided isolation of cytotoxic constituents from the bark of *Aglaia crassinervia* collected in Indonesia. *Bioorg. Med. Chem.* 14: 960-972.
- Tanaka, R., Matsuda, M., and Matsunaga, S. 1987. 3β -Hydroxyhexanordammaran-20-one from *Euphorbia supina*. *Phytochemistry* 26: 3365-3366.

- Terhune, S.J., Hogg, J.W., and Lawrence, B.M. 1974. β -Spathulene: A new sesquiterpene in *Schinus molle* oil. *Phytochemistry* 13: 865-866.
- Trager, W. and Jensen, J.B. 1976. Human malaria parasites in continuous culture. *Science* 193: 673-675.
- Tsukamoto, H., Hisada, S., Nishibe, S., Roux, D.G., and Rourke, J.P. 1984. Coumarins from *Olea Africana* and *Olea capensis*. *Phytochemistry* 23: 699-700.
- Tzouros, M., Bigler, L., Bienz, S., Hesse, M., Inada, A., Murata, H., Inatomi, Y., Nakanishi, T., and Darnaedi, D. 2004. Two new spermidine alkaloids from *Chisocheton weinlandii*. *Helv. Chim. Acta* 87: 1411-1425.
- Valenciennes, E., Smadja, J., and Conan, J.Y. 1999. Screening for biological activity and chemical composition of *Euodia borbonica* var. *borbonica* (Rutaceae), a medicinal plant in Reunion Island. *J. Ethnopharmacol.* 64: 283-288.
- van der Doelen, G.A., van den Berg, K.J., Boon, J.J., Shibayama, N., Rie, R., Genuit, W.J.L. 1998. Analysis of fresh triterpenoid resins and aged triterpenoid varnishes by high-performance liquid chromatography-atmospheric pressure chemical ionisation (tandem) mass spectrometry. *J. Chromatogr. A* 809: 21-37.
- Vinokurova, E.Y., Shults, E.E., Bagryanskaya, I.Y., Gatilov, Y.V., and Tolstikov, G.A. 1999. (+)-Globulol as a new sesquiterpene alcohol from *Angelica sylvestris* L. *Russ. Chem. Bull.* 48: 600-603.
- Vizzotto, L., Muzzi, R.M., and De Lima, D.P. 2003. Complete NMR assignment of (+)- $10\beta,14$ -dihydroxy-*allo*-aromadendrane. *Magn. Reson. Chem.* 41: 1034-1037.
- Wang, B.-G., Peng, H., Huang, H.-L., Li, X.-M., Eck, G., Gong, X., and Proksch, P. 2004. Rocaglamide, aglain, and other related derivatives from *Aglaia testicularis* (Meliaceae). *Biochem. Syst. Ecol.* 32: 1223-1226.
- Wang, S.-K., Cheng, Y.-J., and Duh, C.-Y. 2001. Cytotoxic constituents from leaves of *Aglaia elliptifolia*. *J. Nat. Prod.* 64: 92-94.
- Waterman, P.G. and Ampofo, S. 1985. Dammarane triterpenes from the stem bark of *Commiphora dalzielii*. *Phytochemistry* 24: 2925-2928.
- Wickramaratne, D.B.M., Kumar, V., and Balasubramaniam, S. 1984. Murragleinin, a coumarin from *Murraya gleinei* leaves. *Phytochemistry* 23: 2964-2966.

- Wu, T.-S., Chan, Y.-Y., and Leu, Y.-L. 2000. The constituents of the root and stem of *Aristolochia heterophylla* Hemsl. Chem. Pharm. Bull. 48: 357-361.
- Wu, T.-S., Li, C.-Y., Leu, Y.-L., and Hu, C.-Q. 1999. Limonoids and alkaloids of the root bark of *Dictamnus angustifolius*. Phytochemistry 50: 509-512.
- Xu, Y.-J., Wu, X.-H., Tan, B.K.H., Lai, Y.-H., Vittal, J.J., Imiyabir, Z., Madani, L., Khozirah, K.S., and Goh, S.H. 2000. Flavonol-cinnamate cycloadducts and diamide derivatives from *Aglaia laxiflora*. J. Nat. Prod. 63: 473-476.
- Yadav, R.D., Kataky, J.C.S., and Mathur, R.K. 1999b. New protolimonoids and limonoids. II. Study of the stereochemistry of new tetrnortriterpenoids isolated from root-wood of *Chisocheton paniculatus* Hiern (Meliaceae). J. Indian Chem. Soc. 76: 575-581.
- Yadav, R.D., Kataky, J.C.S., and Mathur, R.K. 1999c. New protolimonoids and limonoids: Part III. Arunachalin, a new tetrnortriterpenoid from *Chisocheton paniculatus* Hiern (Meliaceae). Indian J. Chem. B. Org. 38B: 243-245.
- Yadav, R.D., Kataky, J.C.S., and Mathur, R.K. 1999a. New protolimonoids: part I- Isolation, structure elucidation of new protolimonoids and limonoid from the root wood of *Chisocheton paniculatus* Hiern (Meliaceae). Indian J. Chem. B. Org. 38B: 1359-1363.
- Ye, Q., Qin, G., and Zhao, W. 2002. Immunomodulatory sesquiterpene glycosides from *Dendrobium nobile*. Phytochemistry 61: 885-890.
- Zhao, M.Y., Zhu, Z., and Li, Z. 2000. Effects of scoparone on isolated guinea-pig tracheal smooth muscle. J. Chin. Med. Univ. 29: 335-337.
- Zhao, W., Ye, Q., Tan, X., Jiang, H., Li, X., Chen, K., and Kinghorn, A.D. 2001. Three new sesquiterpene glycosides from *Dendrobium nobile* with immunomodulatory activity. J. Nat. Prod. 64: 1196-1200.

APPENDICES




Figure 9. IR Spectrum of compound CP-1 (KBr disc)




Figure 10. ESI Mass spectrum of compound CP-1




Figure 11a. ^1H NMR (500 MHz) Spectrum of compound CP-1 (in CDCl_3)




Figure 11b. ^1H NMR (500 MHz) Spectrum of compound CP-1 (in CDCl_3)
(expansion δ 0.78-1.23 ppm)




Figure 11c. ^1H NMR (500 MHz) Spectrum of compound CP-1 (CDCl_3)
(expansion δ 1.24-1.86 ppm)




Figure 12. ^{13}C NMR (125 MHz) Spectrum of compound CP-1 (in CDCl_3)




Figure 13a. DEPT 135 Spectrum of compound CP-1




Figure 13b. DEPT 135 Spectrum of compound CP-1 (expansion δ 49.0-55.5 ppm)




Figure 14. ^1H - ^1H COSY Spectrum of compound CP-1




Figure 15a. HMQC Spectrum of compound CP-1 (δ_{H} 0.7-1.1 ppm, δ_{C} 14-20 ppm)




Figure 15b. HMQC Spectrum of compound CP-1 (δ_{H} 0.8-1.8 ppm, δ_{C} 18-37 ppm)




Figure 15c. HMQC Spectrum of compound CP-1 (δ_{H} 0.8-2.7 ppm, δ_{C} 30-60 ppm)




Figure 16a. HMBC Spectrum of compound CP-1 (δ_H 1.85-2.7 ppm, δ_C 13-29 ppm)




Figure 16b. HMBC Spectrum of compound CP-1 (δ_H 1.85-2.7 ppm, δ_C 30-60 ppm)




Figure 16c. HMBC Spectrum of compound CP-1 (δ_H 1.85-2.7 ppm, δ_C 200-220 ppm)




Figure 16d. HMBC Spectrum of compound CP-1 (δ_H 0.80-1.10 ppm, δ_C 13-29 ppm)




Figure 16e. HMBC Spectrum of compound CP-1 (δ_H 0.80-1.10 ppm, δ_C 30-60 ppm)




Figure 16f. HMBC Spectrum of compound CP-1
(δ_H 0.80-1.10 ppm, δ_C 207-220 ppm)




Figure 16g. HMBC Spectrum of compound CP-1
(δ_{H} 1.10-1.75 ppm, δ_{C} 207-220 ppm)




Figure 16h. HMBC Spectrum of compound CP-1 (δ_{H} 1.10-1.75 ppm, δ_{C} 30-60 ppm)




Figure 16i. HMBC Spectrum of compound CP-1 (δ_H 1.10-1.75 ppm, δ_C 13-30 ppm)




Figure 17. NOESY Spectrum of compound CP-1




Figure 18. IR Spectrum of compound CP-2 (KBr disc)




Figure 19. ESI Mass spectrum of compound CP-2




Figure 20a. ^1H NMR (500 MHz) Spectrum of compound CP-2 (in CDCl_3)




Figure 20b. ^1H NMR (500 MHz) Spectrum of compound CP-2 (in CDCl_3)
(expansion δ 0.84-1.69 ppm)




Figure 20c. ^1H NMR (500 MHz) Spectrum of compound CP-2 (in CDCl_3)
(expansion δ 1.80-5.15 ppm)




Figure 21. ^{13}C NMR (125 MHz) Spectrum of compound CP-2 (in CDCl_3)




Figure 22a. DEPT 135 Spectrum of compound CP-2




Figure 22b. DEPT 135 Spectrum of compound CP-2
(expansion δ 100-135 ppm)




Figure 23. ^1H - ^1H COSY Spectrum of compound CP-2




Figure 24a. HMQC Spectrum of compound CP-2
(δ_{H} 0.8-2.3 ppm, δ_{C} 19-33 ppm)




Figure 24b. HMQC Spectrum of compound CP-2
(δ_{H} 0.7-1.7 ppm, δ_{C} 14-28 ppm)




Figure 24c. HMQC Spectrum of compound CP-2
(δ_{H} 0.7-1.7 ppm, δ_{C} 28-60 ppm)




Figure 24d. HMQC Spectrum of compound CP-2
(δ_H 1.8-2.7 ppm, δ_C 25-58 ppm)




Figure 24e. HMQC Spectrum of compound CP-2
(δ_H 4.1-5.9 ppm, δ_C 106-126 ppm)




Figure 24f. HMQC Spectrum of compound CP-2
(δ_H 0.75-1.15 ppm, δ_C 14-23 ppm)




Figure 25a. HMBC Spectrum of compound CP-2
(δ_H 4.55-5.45 ppm, δ_C 30-60 ppm)




Figure 25b. HMBC Spectrum of compound CP-2
(δ_H 4.55-5.45 ppm, δ_C 11-33 ppm)




Figure 25c. HMBC Spectrum of compound CP-2
(δ_H 1.8-2.9 ppm, δ_C 150-220 ppm)




Figure 25d. HMBC Spectrum of compound CP-2
 $(\delta_H 1.75-2.65 \text{ ppm}, \delta_C 107-136 \text{ ppm})$




Figure 25e. HMBC Spectrum of compound CP-2
 $(\delta_H 1.75-2.65 \text{ ppm}, \delta_C 30-60 \text{ ppm})$




Figure 25f. HMBC Spectrum of compound CP-2
 $(\delta_H$ 1.80-2.55 ppm, δ_C 11-33 ppm)




Figure 25g. HMBC Spectrum of compound CP-2
 $(\delta_H$ 0.6-1.7 ppm, δ_C 150-220 ppm)




Figure 25h. HMBC Spectrum of compound CP-2
(δ_H 0.6-1.7 ppm, δ_C 107-136 ppm)




Figure 25i. HMBC Spectrum of compound CP-2
(δ_H 0.70-1.25 ppm, δ_C 34-59 ppm)




Figure 25j. HMBC Spectrum of compound CP-2
(δ_{H} 0.80-1.75 ppm, δ_{C} 11-33 ppm)




Figure 26. NOESY Spectrum of compound CP-2




Figure 27. IR Spectrum of compound CP-3 (KBr disc)




Figure 28. EI Mass spectrum of compound CP-3




Figure 29a. ^1H NMR (500 MHz) Spectrum of compound CP-3 (in CDCl_3)




Figure 29b. ^1H NMR (500 MHz) Spectrum of compound CP-3 (in CDCl_3) (expansion δ 0.80-1.70 ppm)




Figure 29c. ^1H NMR (500 MHz) Spectrum of compound CP-3 (in CDCl_3)
(expansion δ 1.32-3.35 ppm)




Figure 30. ^{13}C NMR (125 MHz) Spectrum of compound CP-3 (in CDCl_3)




Figure 31. DEPT 135 Spectrum of compound CP-3




Figure 32. ^1H - ^1H COSY Spectrum of compound CP-3




Figure 33a. HMQC Spectrum of compound CP-3
(δ_H 0.80-2.00 ppm, δ_C 16-52 ppm)




Figure 33b. HMQC Spectrum of compound CP-3
(δ_H 3.0-4.0 ppm, δ_C 74-89 ppm)




Figure 34a. HMBC Spectrum of compound CP-3
(δ_H 1.00-1.30 ppm, δ_C 30-39 ppm)




Figure 34b. HMBC Spectrum of compound CP-3
(δ_H 1.14-1.34 ppm, δ_C 15-23 ppm)




Figure 34c. HMBC Spectrum of compound CP-3
(δ_{H} 1.14-1.34 ppm, δ_{C} 24-30 ppm)




Figure 34d. HMBC Spectrum of compound CP-3
(δ_{H} 1.34-1.68 ppm, δ_{C} 15-23 ppm)




Figure 34e. HMBC Spectrum of compound CP-3
(δ_{H} 1.35-2.00 ppm, δ_{C} 30-53 ppm)




Figure 34f. HMBC Spectrum of compound CP-3
(δ_{H} 1.35-2.00 ppm, δ_{C} 68-90 ppm)




Figure 34g. HMBC Spectrum of compound CP-3
(δ_H 1.06-1.18 ppm, δ_C 85-88 ppm)




Figure 34h. HMBC Spectrum of compound CP-3
(δ_H 1.58-1.90 ppm, δ_C 84.5-88.5 ppm)




Figure 34i. HMBC Spectrum of compound CP-3
(δ_{H} 0.75-1.20 ppm, δ_{C} 15-29 ppm)




Figure 34j. HMBC Spectrum of compound CP-3
(δ_{H} 0.75-1.20 ppm, δ_{C} 31-52 ppm)




Figure 34k. HMBC Spectrum of compound CP-3
(δ_H 3.35-3.75 ppm, δ_C 31-52 ppm)




Figure 34l. HMBC Spectrum of compound CP-3
(δ_H 3.35-3.75 ppm, δ_C 21-30 ppm)




Figure 34m. HMBC Spectrum of compound CP-3
 $(\delta_H \text{ 3.35-3.75 ppm}, \delta_C \text{ 69-89 ppm})$




Figure 35. NOESY Spectrum of compound CP-3




Figure 36. IR Spectrum of compound CP-4 (KBr disc)




Figure 37. ESI-TOF Mass spectrum of compound CP-4




Figure 38a. ^1H NMR (500 MHz) Spectrum of compound CP-4 (in CDCl_3)




Figure 38b. ^1H NMR (500 MHz) Spectrum of compound CP-4 (in CDCl_3)
(expansion δ 0.0–1.1 ppm)




Figure 38c. ^1H NMR (500 MHz) Spectrum of compound CP-4 (in CDCl_3)
(expansion δ 1.0–1.7 ppm)




Figure 38d. ^1H NMR (500 MHz) Spectrum of compound CP-4 (in CDCl_3)
(expansion δ 1.70-2.04 ppm)




Figure 39. ^{13}C NMR (125 MHz) Spectrum of compound CP-4 (in CDCl_3)




Figure 40a. DEPT 135 Spectrum of compound CP-4




Figure 40b. DEPT 135 Spectrum of compound CP-4 (expansion δ 16-30 ppm)




Figure 41. ¹H-¹H COSY Spectrum of compound CP-4




Figure 42. HMQC Spectrum of compound CP-4




Figure 43a. HMBC Spectrum of compound CP-4




Figure 43b. HMBC Spectrum of compound CP-4 (δ_{H} 1.4-3.7 ppm, δ_{C} 40-95 ppm)




Figure 43c. HMBC Spectrum of compound CP-4 (δ_{H} 2.7-4.1 ppm, δ_{C} 14-43 ppm)




Figure 43d. HMBC Spectrum of compound CP-4 (δ_{H} 0.0-1.2 ppm, δ_{C} 25-33 ppm)




Figure 43e. HMBC Spectrum of compound CP-4 (δ_{H} 0.75-1.20 ppm, δ_{C} 11-43 ppm)




Figure 43f. HMBC Spectrum of compound CP-4 (δ_H 1.2-2.5 ppm, δ_C 15-45 ppm)




Figure 44. NOESY Spectrum of compound CP-4




Figure 45. IR Spectrum of compound CP-5




Figure 46. ESI-TOF Mass spectrum of compound CP-5




Figure 47a. ^1H NMR (500 MHz) Spectrum of compound CP-5 (in CDCl_3)




Figure 47b. ^1H NMR (500 MHz) Spectrum of compound CP-5 (in CDCl_3)
(expansion δ 0.6-2.2 ppm)




Figure 48. ^{13}C NMR (125 MHz) Spectrum of compound CP-5 (in CDCl_3)




Figure 49a. DEPT 135 Spectrum of compound CP-5




Figure 49b. DEPT 135 Spectrum of compound CP-5
(expansion δ 15-45 ppm)




Figure 50. ¹H-¹H COSY Spectrum of compound CP-5




Figure 51. HMQC Spectrum of compound CP-5




Figure 52. HMBC Spectrum of compound CP-5




Figure 53a. NOESY Spectrum of compound CP-5




Figure 53b. NOESY Spectrum of compound CP-5 (δ 0.5-2.4 ppm)




Figure 54. IR Spectrum of compound CP-6 (KBr disc)




Figure 55. EI Mass spectrum of compound CP-6




Figure 56a. ^1H NMR (500 MHz) Spectrum of compound CP-6 (in CDCl_3)




Figure 56b. ^1H NMR (500 MHz) Spectrum of compound CP-6 (in CDCl_3)
 (expansion δ 0.8-2.8 ppm)




Figure 57a. ^{13}C NMR (125 MHz) Spectrum of compound CP-6 (in CDCl_3)




Figure 57b. ^{13}C NMR (125 MHz) Spectrum of compound CP-6 (in CDCl_3)
(expansion δ 15-51 ppm)





Figure 58. DEPT 135 Spectrum of compound CP-6

Figure 59. ^1H - ^1H COSY Spectrum of compound CP-6




Figure 60a. HMQC Spectrum of compound CP-6




Figure 60b. HMQC Spectrum of compound CP-6
(δ_{H} 4.0-6.0 ppm, δ_{C} 108-120 ppm)




Figure 61a. HMBC Spectrum of compound CP-6




Figure 61b. HMBC Spectrum of compound CP-6

(δ_H 4.0-6.0 ppm, δ_C 15-60 ppm)




Figure 61c. HMBC Spectrum of compound CP-6

(δ_H 1.2-3.0 ppm, δ_C 80-190 ppm)




Figure 62. NOESY Spectrum of compound CP-6




Figure 63. IR Spectrum of compound CP-7




Figure 64. IR Spectrum of compound CP-7 (KBr disc)




Figure 65. ¹H NMR (500 MHz) Spectrum of compound CP-7 (in CDCl₃)




Figure 66. ¹³C NMR (125 MHz) Spectrum of compound CP-7 (in CDCl₃)




Figure 67a. DEPT 135 Spectrum of compound CP-7




Figure 67b. DEPT 135 Spectrum of compound CP-7
(expansion δ 15–32 ppm)




Figure 67c. DEPT 135 Spectrum of compound CP-7
(expansion δ 33-56 ppm)




Figure 68. IR Spectrum of compound CP-8 (KBr disc)




Figure 69. EI Mass spectrum of compound CP-8




Figure 70a. ^1H NMR (500 MHz) Spectrum of compound CP-8 (in CDCl_3)




Figure 70b. ^1H NMR (500 MHz) Spectrum of compound CP-8 (in CDCl_3)
(expansion δ 0.80-1.52 ppm)




Figure 70c. ^1H NMR (500 MHz) Spectrum of compound CP-8 (in CDCl_3)
(expansion δ 1.68-3.40 ppm)




Figure 71. ^{13}C NMR (125 MHz) Spectrum of compound CP-8 (in CDCl_3)




Figure 72a. DEPT 135 Spectrum of compound CP-8
(expansion δ 14-32 ppm)




Figure 72b. DEPT 135 Spectrum of compound CP-8
(expansion δ 33-51 ppm)




Figure 73. ^1H - ^1H COSY Spectrum of compound CP-8




Figure 74a. HMQC Spectrum of compound CP-8




Figure 74b. HMQC Spectrum of compound CP-8

(δ_{H} 0.75-1.65 ppm, δ_{C} 30-54 ppm)




Figure 74c. HMQC Spectrum of compound CP-8

(δ_H 1.0-2.8 ppm, δ_C 13-30ppm)




Figure 74d. HMQC Spectrum of compound CP-8
(δ_H 1.4-2.8 ppm, δ_C 20-30 ppm)




Figure 74e. HMQC Spectrum of compound CP-8
(δ_{H} 1.0-2.8 ppm, δ_{C} 30- 54ppm)




Figure 74f. HMQC Spectrum of compound CP-8
(δ_{H} 2.4-3.5 ppm, δ_{C} 62- 98ppm)




Figure 75a. HMBC Spectrum of compound CP-8




Figure 75b. HMBC Spectrum of compound CP-8
(δ_H 0.80-1.04 ppm, δ_C 13-30ppm)




Figure 75c. HMBC Spectrum of compound CP-8
(δ_H 1.09-1.65 ppm, δ_C 13-30ppm)




Figure 75d. HMBC Spectrum of compound CP-8
(δ_H 0.80-1.55 ppm, δ_C 74-91ppm)




Figure 75e. HMBC Spectrum of compound CP-8
(δ_{H} 1.6-3.5 ppm, δ_{C} 30-54 ppm)




Figure 75f. HMBC Spectrum of compound CP-8
(δ_{H} 1.6-3.5 ppm, δ_{C} 70-180 ppm)




Figure 76. NOESY Spectrum of compound CP-8




Figure 77. UV Spectrum of compound CP-9 (MeOH)




Figure 78. IR Spectrum of compound CP-9 (KBr disc)




Figure 79. EI- Mass spectrum of compound CP-9




Figure 80. ¹H NMR (500 MHz) Spectrum of compound CP-9 (in CDCl₃)




Figure 81. ¹³C NMR (125 MHz) Spectrum of compound CP-9 (in CDCl₃)




Figure 82. DEPT 135 Spectrum of compound CP-9




Figure 83. ^1H - ^1H COSY Spectrum of compound CP-9




Figure 84. HMQC Spectrum of compound CP-9




Figure 85. HMBC Spectrum of compound CP-9




Figure 86. UV Spectrum of compound CP-10 (MeOH)




Figure 87. IR Spectrum of compound CP-10 (KBr disc)




Figure 88. ESI-TOF Mass spectrum of compound CP-10




Figure 89. ¹H NMR (500 MHz) Spectrum of compound CP-10 (in DMSO-d₆)




Figure 90. ^{13}C NMR (125 MHz) Spectrum of compound CP-10
(in $\text{DMSO}-d_6$)




Figure 91. DEPT 135 Spectrum of compound CP-10




Figure 92. ^1H - ^1H COSY Spectrum of compound CP-10




Figure 93. HMQC Spectrum of compound CP-10




Figure 94. HMBC Spectrum of compound CP-10




Figure 95. UV Spectrum of compound CP-11 (MeOH)




Figure 96. IR Spectrum of compound CP-11 (KBr disc)




Figure 97. EI Mass spectrum of compound CP-11




Figure 98a. ^1H NMR (500 MHz) Spectrum of compound CP-11 (in $\text{DMSO}-d_6$)




Figure 98b. ^1H NMR (500 MHz) Spectrum of compound CP-11 (in $\text{DMSO}-d_6$) (expansion δ 6.5-7.7 ppm)




Figure 99. ^{13}C NMR (125 MHz) Spectrum of compound CP-11 (in $\text{DMSO}-d_6$)




Figure 100. DEPT 135 Spectrum of compound CP-11




Figure 101. ^1H - ^1H COSY Spectrum of compound CP-11




Figure 102. HMQC Spectrum of compound CP-11




Figure 103. HMBC Spectrum of compound CP-11




Figure 104. ^1H NMR (500 MHz) Spectrum of compound CP-12 (in $\text{DMSO}-d_6$)




Figure 105a. ^{13}C NMR (125 MHz) Spectrum of compound CP-12 (in $\text{DMSO}-d_6$)




Figure 105b. ^{13}C NMR (125 MHz) Spectrum of compound CP-12 (in $\text{DMSO}-d_6$)
(expanded δ 10-50 ppm)




Figure 105c. ^{13}C NMR (125 MHz) Spectrum of compound CP-12 (in $\text{DMSO}-d_6$) (expansion δ 45-80 ppm)




Figure 106. DEPT 135 Spectrum of compound CP-12




Figure 107. IR Spectrum of compound CP-13 (KBr disc)




Figure 108. ESI Mass spectrum of compound CP-13




Figure 109. ¹H NMR (500 MHz) Spectrum of compound CP-13 (in CDCl_3)




Figure 110. ¹³C NMR (125 MHz) Spectrum of compound CP-13 (in CDCl_3)




Figure 111. DEPT 135 Spectrum of compound CP-13




Figure 112. ¹H-¹H COSY Spectrum of compound CP-13




Figure 113. HMQC Spectrum of compound CP-13




Figure 114a. HMBC Spectrum of compound CP-13




Figure 114b. HMBC Spectrum of compound CP-13 (δ_H 0.1-2.0 ppm, δ_C 50-85 ppm)




Figure 114c. HMBC Spectrum of compound CP-13 (δ_H 3.0-3.8 ppm, δ_C 9-34 ppm)





Figure 114d. HMBC Spectrum of compound CP-13 (δ_H 3.0-3.8 ppm, δ_C 66-82 ppm)






Figure 116. UV Spectrum of compound CP-14 (MeOH)




Figure 117. IR Spectrum of compound CP-14 (KBr disc)




Figure 118. ESI-TOF Mass spectrum of compound CP-14




Figure 119. ^1H NMR (500 MHz) Spectrum of compound CP-14 (in CDCl_3)




Figure 120. ^{13}C NMR (125 MHz) Spectrum of compound CP-14 (in CDCl_3)




Figure 121. DEPT 135 Spectrum of compound CP-14




Figure 122. ^1H - ^1H COSY Spectrum of compound CP-14




Figure 123. HMQC Spectrum of compound CP-14




Figure 124. HMBC Spectrum of compound CP-14




Figure 125. UV Spectrum of compound CP-15 (MeOH)




Figure 126. IR Spectrum of compound CP-15 (KBr disc)




Figure 127. ESI- Mass spectrum (Negative mode) of compound CP-15




Figure 128a. ^1H NMR (500 MHz) Spectrum of compound CP-15 (in CDCl_3)




Figure 128b. ^1H NMR (500 MHz) Spectrum of compound CP-15 (in CDCl_3)
(expansion δ 1.0-4.8 ppm)




Figure 128c. ^1H NMR (500 MHz) Spectrum of compound CP-15 (in CDCl_3) (expansion δ 6.0-12.0 ppm)




Figure 129. ^{13}C NMR (125 MHz) Spectrum of compound CP-15 (in CDCl_3)




Figure 130. DEPT 135 Spectrum of compound CP-15




Figure 131. ^1H - ^1H COSY Spectrum of compound CP-15




Figure 132. HMQC Spectrum of compound CP-15




Figure 133. HMBC Spectrum of compound CP-15




Figure 134. UV Spectrum of compound AE-4 (MeOH)




Figure 135. IR Spectrum of compound AE-4 (KBr disc)




Figure 136. EI Mass spectrum of compound AE-4




Figure 137a. ¹H NMR (500 MHz) Spectrum of compound AE-4 (in CDCl₃)




Figure 137b. ^1H NMR (500 MHz) Spectrum of compound AE-4 (in CDCl_3)
(expansion δ 6.0-7.4 ppm)




Figure 138. ^{13}C NMR (125 MHz) Spectrum of compound AE-4 (in CDCl_3)




Figure 139. DEPT 135 Spectrum of compound AE-4




Figure 140. ^1H - ^1H COSY Spectrum of compound AE-4




Figure 141. HMQC Spectrum of compound AE-4




Figure 142. HMBC Spectrum of compound AE-4




Figure 143. IR Spectrum of compound AE-5 (KBr disc)




Figure 144. ESI-TOF mass spectrum of compound AE-5




Figure 145a. ^1H NMR (500 MHz) Spectrum of compound AE-5 (in CDCl_3)




Figure 145b. ^1H NMR (500 MHz) Spectrum of compound AE-5 (in CDCl_3)
(expansion δ 0.8-2.4 ppm)




Figure 145c. ^1H NMR (500 MHz) Spectrum of compound AE-5 (in CDCl_3)
(expansion δ 3.6-5.0 ppm)




Figure 146. ^{13}C NMR (125 MHz) Spectrum of compound AE-5 (in CDCl_3)




Figure 147a. DEPT 135 Spectrum of compound AE-5
(expansion δ 14-38 ppm)




Figure 147b. DEPT 135 Spectrum of compound AE-5
(expansion δ 38-90 ppm)




Figure 147c. DEPT 135 Spectrum of compound AE-5




Figure 148. ^1H - ^1H COSY Spectrum of compound AE-5




Figure 149a. HMQC Spectrum of compound AE-5




Figure 149b. HMQC Spectrum of compound AE-5
(δ_{H} 3.5-5.0 ppm, δ_{C} 55-115 ppm)




Figure 150a. HMBC Spectrum of compound AE-5




Figure 150b. HMBC Spectrum of compound AE-5
(δ_H 0.6-2.6 ppm, δ_C 10-55 ppm)




Figure 150c. HMBC Spectrum of compound AE-5
 $(\delta_H \text{ } 0.8\text{-}2.0 \text{ ppm}, \delta_C \text{ } 59\text{-}90 \text{ ppm})$




Figure 150d. HMBC Spectrum of compound AE-5
 $(\delta_H \text{ } 1.4\text{-}2.5 \text{ ppm}, \delta_C \text{ } 110\text{-}180 \text{ ppm})$




Figure 150e. HMBC Spectrum of compound AE-5
 $(\delta_H$ 3.6-5.0 ppm, δ_C 10-65 ppm)




Figure 151a. NOESY Spectrum of compound AE-5




Figure 151b. NOESY Spectrum of compound AE-5
(expansion δ 0.8-2.4 ppm)




Figure 152. IR Spectrum of compound AE-7 (KBr disc)




Figure 153. ESI Mass spectrum (Negative mode) of compound AE-7




Figure 154a. ^1H NMR (500 MHz) Spectrum of compound AE-7 (in CDCl_3)




Figure 154b. ^1H NMR (500 MHz) Spectrum of compound AE-7 (in CDCl_3)
(expansion δ 0.8-2.8 ppm)




Figure 155. ^{13}C NMR (125 MHz) Spectrum of compound AE-7 (in CDCl_3)




Figure 156a. DEPT 135 Spectrum of compound AE-7
(expansion δ 15-37 ppm)




Figure 156b. DEPT 135 Spectrum of compound AE-7
(expansion δ 39-52 ppm)




Figure 156c. DEPT 135 Spectrum of compound AE-7
(expansion δ 75-115 ppm)




Figure 157. ^1H - ^1H COSY Spectrum of compound AE-7




Figure 158a. HMQC Spectrum of compound AE-7




Figure 158b. HMQC Spectrum of compound AE-7
(δ_{H} 1.0–2.6 ppm, δ_{C} 20–55 ppm)





Figure 158c. HMQC Spectrum of compound AE-7
(δ_{H} 4.4-5.1 ppm, δ_{C} 106-120 ppm)






Figure 159b. HMBC Spectrum of compound AE-7
(δ_{H} 0.7-1.9 ppm, δ_{C} 65-100 ppm)




Figure 159c. HMBC Spectrum of compound AE-7
(δ_{H} 0.7-2.8 ppm, δ_{C} 100-185 ppm)




Figure 159d. HMBC Spectrum of compound AE-7
(δ_{H} 4.4-5.1 ppm, δ_{C} 10-60 ppm)




Figure 160. UV Spectrum of compound AE-10 (MeOH)




Figure 161. IR Spectrum of compound AE-10 (KBr disc)




Figure 162. ESI-TOF Mass spectrum of compound AE-10




Figure 163a. ^1H NMR (500 MHz) Spectrum of compound AE-10 (in CDCl_3)




Figure 163b. ^1H NMR (500 MHz) Spectrum of compound AE-10 (in CDCl_3)
(expansion δ 0.8-4.5 ppm)




Figure 163c. ^1H NMR (500 MHz) Spectrum of compound AE-10 (in CDCl_3)
(expansion δ 5.0-7.5 ppm)




Figure 164a. ^{13}C NMR (125 MHz) Spectrum of compound AE-10 (in CDCl_3)





Figure 164b. ^{13}C NMR (125 MHz) Spectrum of compound AE-10 (in CDCl_3) (expansion δ 5-80 ppm)






Figure 165. DEPT 135 Spectrum of compound AE-10




Figure 166a. ^1H - ^1H COSY Spectrum of compound AE-10 (expansion δ 3.2-5.4 ppm)




Figure 166b. ^1H - ^1H COSY Spectrum of compound AE-10 (expansion δ 5.5-8.5 ppm)




Figure 167. HMQC Spectrum of compound AE-10




Figure 168a. HMBC Spectrum of compound AE-10




Figure 168b. HMBC Spectrum of compound AE-10 (δ_{H} 3.2-5.4 ppm, δ_{C} 40-85 ppm)




Figure 169. NOESY Spectrum of compound AE-10

VITA

Miss Jarinporn Phongmaykin was born on July 11, 1977 in Bangkok, Thailand. She received her Bachelor Degree of Pharmacy (1st class honor) from the Faculty of Pharmaceutical Sciences, Huachiew University in 1999. She was granted a Royal Golden Jubilee Ph.D. Scholarship from the Thailand Research Fund (TRF) in the year 2001.

Poster Presentations

1. Phongmaykin, J., Suttisri, R., Saifah, E., Kumamoto, T., Ishikawa, T. Chemical Constituents of Fruits of *Chisocheton tomentosus*. RGJ-Ph.D. Congress VII. April 20-22, 2006. Jomtien Palm Beach Resort, Pattaya, Thailand.
2. Phongmaykin, J., Suttisri, R., Saifah, E., Kumamoto, T., Ishikawa, T. Sesquiterpenoids and triterpenoids from *Chisocheton penduliflorus*. Proceeding of the 23rd Annual Research Conference in Pharmaceutical Sciences & JSPS 1st Medical Chemistry Seminar of Asia/Africa Science Platform Program. December 14-15, 2006, Bangkok, Thailand.