



## Chapter 4

### $T_c$ equations

In this chapter I will show how to obtain  $T_c$  from Eq.(3.55).

Since the structure of the matrix in equation (3.55), which determines  $T_c$ , depends on the nature of different conducting layers in the unit cell and their surrounding.

The transition temperature  $T_c$  for a three-layer type is given by the solution of the determinant,

$$\begin{vmatrix}
 1 - N_d V_d F + K(1 - V_{er}) & -[V_d V_{er} N_d F + K(1 - V_{er})] & 0 \\
 \frac{V_d}{V_c} & \frac{V_c}{V_d} & \\
 -[V_c V_{er} N_c F + K(1 - V_{er})] & 1 - N_c V_c F + K(1 - V_{er}) & -[V_c V_{er} N_c F + K(1 - V_{er})] \\
 \frac{V_d}{V_c} & \frac{V_d}{V_c} & \frac{V_d}{V_c} \\
 0 & -[V_d V_{er} N_d F + K(1 - V_{er})] & 1 - N_d V_d F + K(1 - V_{er}) \\
 & \frac{V_c}{V_d} & \frac{V_d}{V_c}
 \end{vmatrix} = 0$$

### One superconducting layer per unit cell

We first consider the one layer per unit cell system,



Fig. 4.1 One superconducting layers per unit cell

here  $V_d = 0$ ,  $K = K' = 0$  and hence

$$1 - N_c V_c F = 0$$

where

$$F = 2 \pi T_c \sum_{n=0} \frac{1}{\omega_n}$$

$$= \ln(1.13 \omega_D / T_c) \quad (\text{see appendix})$$

then

$$T_c = 1.13 \omega \exp[-1/N_c V_c] \quad (4.1)$$

here  $\omega$  is the cutoff frequency.

#### Two superconducting layers per unit cell




Fig. 4.2 Two superconducting layers per unit cell

Let us now calculate the critical temperature for the sample where there are two superconducting layers per unit cell, our  $T_c$  is determined by the condition

$$\begin{vmatrix} 1 - N_d V_d F + K'(1 - V_{er}) & -[V_d V_{er} N_d F + K'(1 - V_{er})] \\ \frac{V_d}{V_c} & \frac{V_c}{V_c} \\ -[V_c V_{er} N_c F + K(1 - V_{er})] & 1 - N_c V_c F + K(1 - V_{er}) \\ \frac{V_d}{V_c} & \frac{V_c}{V_c} \end{vmatrix} = 0. \quad (4.2)$$

Solving Eq.(4.2), we obtain

$$T_c = 1.13 \exp \left[ - \frac{N_c V_c + N_d V_d + N_d K \left( \frac{V_d - V_{er}}{V_c} \right) + N_c K \left( \frac{V_c - V_{er}}{V_d} \right)}{2N_c N_d (V_c V_d - V_{er}^2)} \right. \\ \left. + \left[ \frac{[N_c V_c (1+K) + N_d V_d (1+K) - V_{er}^2 (N_d K + N_c K)]^2}{- 4N_c N_d (V_c V_d - V_{er}^2) \left( 1 + K \left( 1 - \frac{V_{er}}{V_c} \right) + K \left( 1 - \frac{V_{er}}{V_d} \right) \right)} \right]^{1/2} / 2N_c N_d (V_c V_d - V_{er}^2) \right] \quad (4.3)$$

In solving Eq.(4.2) for F, we obtain two solutions for  $T_c$ . The solution which gives the highest  $T_c$  is acceptable, while the other with the lower  $T_c$  value is rejected because the system is already superconducting once the system reaches the high  $T_c$  value.

Consider limit  $t$  tends to zero,  $N_c = N_d$ , and  $V_c = V_d$ .

From Eq.(4.3) we obtain

$$T_c = 1.13 \omega \exp(-1/N_c (V_c + V_{er})) \quad (4.4)$$

which is precisely the result of Ihm and Yu (46).

From equation (4.3), it is concluded that  $T_c$  is always raised by the presence of  $V_{er}$ . We also see clearly that the effect of small but finite  $t$  is detrimental to high- $T_c$  superconductivity in consistent with ref.(55). Recently Cava et al.(56) synthesized  $La_{1-x}Sr_{0.4}CaCuO_2O_e$ , the layer-stacking sequence in this compound is  $LaO-CuO_2-Ca-CuO_2$ . This system is a bi-layer superconductor. The upper and lower  $CuO_2$  layers of this system are equivalent.

Equation (4.3) also gives

$$T_c = 1.13\omega \exp(-1/N_c(V_c + V_{er})) \quad (4.5)$$

for this system, assuming that  $t \neq 0$ . Eqs. (4.4) and (4.5) are identical this is because of the fact that in a system of equivalent double layers, whether there is a direct electron hopping process or not, does not change the physical situation at all.

For two inequivalent double layer, Eq. (4.3) yields

$$T_c = 1.13\omega \exp \left[ - \frac{N_c V_c + N_d V_d + N_d K(V_d - V_{er})^2 + N_c K(V_c - V_{er})^2}{2N_c N_d (V_c V_d - V_{er}^2)} \right] \quad (4.6)$$

where  $t=0$ .

#### Three identical layers per unit cell




Fig. 4.3 Three identical layers per unit cell

For three identical layers, the interactions  $V_c = V_d$ , if  $t=0$  and  $N_c = N_d$ , the 3X3 determinant of equation (3.55) gives

$$T_c = 1.13\omega \exp(-1/N_c(V_c + \sqrt{2} V_{er})) \quad (4.7)$$

We recover the formula of Ihm and Yu (46).

We next consider, for example the compound  $Tl_2Ba_2Ca_2Cu_3O_{10}$ , here the TlO-TlO-CuO<sub>2</sub> layers bounding the middle CuO<sub>2</sub> sheets are mediating the interaction and maintaining its 2 dimensional property. These layers have the density of states, at the Fermi level, of the same magnitude with the carrier from the middle CuO<sub>2</sub> layer. It is plausible that the excitations from the TlO-TlO layers have a strong influence on the interactions in the bounding CuO<sub>2</sub> layer, the two outer CuO<sub>2</sub> layers are therefore different from the middle CuO<sub>2</sub> layer. We have for this system

$$\begin{aligned}
 T_c = 1.13\omega \exp \left[ - \left[ \frac{N_c V_c (1+K(1-V_{er})) + N_d V_d (1+K(1-V_{er}))}{V_d V_c} \right. \right. \\
 + \frac{2KV_{er} [(1-V_{er}) \frac{V_d N_d}{V_c} + (1-V_{er}) \frac{V_c V_{er}}{V_d}]}{2N_c N_d (V_c V_d - 2V_{er}^2)} \\
 - \left[ \left( \frac{N_c V_c (1+K(1-V_{er})) + N_d V_d (1+K(1-V_{er}))}{V_d V_c} + 2KV_{er} \left( \frac{(1-V_{er}) V_d N_d}{V_d V_c} \right. \right. \right. \\
 \left. \left. \left. + \frac{(1-V_{er}) V_c V_{er}}{V_d} \right) \right)^2 - 4N_c N_d [V_c V_d - 2V_{er}^2] \left[ \frac{1+K(1-V_{er})}{V_c} + \frac{K(1-V_{er})}{V_d} \right] \right. \\
 \left. \left. - \frac{KK(1-V_{er})(1-V_{er})}{V_c V_d} \right]^{1/2} / 2N_c N_d (V_c V_d - 2V_{er}^2) \right] \quad (4.8)
 \end{aligned}$$

For the bounding layers of insulating nature  $V_d=0=V_{er}$ , non-zero hopping becomes important and we have a system of a superconducting plane interposed between two insulating sheets.




Fig. 4.4 A superconducting layer between two insulating layers.

Equation (4.8) yields

$$T_c = 1.13\omega \exp\left\{-1/N_c V_c - \frac{z^2 t^2 N_d}{4T_{co}}\right\}. \quad (4.9)$$

The critical temperature decreases with the increased direct hopping between layers.

In order to further understand the physics of high- $T_c$  superconductor it is important to study the influence of the decoupling of the  $\text{CuO}_2$  bilayer by the insulating layer.




Fig. 4.5 An insulating layer between two superconducting layers.

Here  $V_c = 0 = V_{er}$ , and Eq.(4.8) yields

$$T_c = 1.13\omega \exp\left\{-1/N_d V_d - \frac{z^2 t^2 N_c}{4T_{co}}\right\}. \quad (4.10)$$

This last equation indicates that the coupling between the  $\text{CuO}_2$  layers across the insulating layer is reduced.