

REFERENCES

- Alderton LE, Spector LG, Blair CK, et al. Child and maternal household chemical exposure and the risk of acute leukemia in children with Down's syndrome: a report from the Children's Oncology Group. Am J Epidemiol. 2006; 164 (3): 212-221.
- Andrew Williams J, David H, Phillip, et al. Mammary Expression of Xenobiotic Metabolizing Enzymes and Their Potential Role in Breast Cancer¹. CANCER RESEARCH. 2000; 60: 4667-4677.
- Balta G, Yuksek N, Ozyurek E, et al. Characterization of *MTHFR*, *GSTM1*, *GSTT1*, *GSTP1*, and *CYP1A1* genotypes in childhood acute leukemia. Am J Hematol. 2003 ;73 (3):154-160.
- Bell DA, Stephens EA, Castranio T, et al. Polyadenylation polymorphism in the acetyltransferase 1 gene (*NAT1*) increases risk of colorectal cancer. Cancer Res. 1995; 55 (16): 3537-42.
- Bennett JM, Catovsky D, Daniel MT, et al., Proposal for the classification of the acute leukemia (FAB Cooperative Group) Br journal hematol. 1976; 33:451.
- Bolt H. M, R Their, et al. Relevance of the deletion polymorphisms of the glutathione-S-transferase *GSTT1* and *GSTM1* in pharmacology and toxicology. Current Drug Metabolism. 2006; 7: 613-628.
- Bolufer P, Collado M, Barragán E, et al. The potential effect of gender in combination with common genetic polymorphisms of drug-metabolizing enzymes on the risk of developing acute leukemia. Haematologica. 2007; 92 (3): 308-314.
- Cascorbi I, Brockmøller J, Roots I, et al. A C4887A polymorphism in exon 7 of human *CYP1A1*: Population frequency, mutation linkages, and impact on lung cancer susceptibility. Cancer Res. 1996; 56: 4965.
- Cascorbi I, Brockmøller J, Mrozikiewicz PM, et al. Homozygous rapid arylamine N-acetyltransferase (*NAT2*) genotype as a susceptibility factor for lung cancer. Cancer Res. 1996; 56 (17): 3961-6.

- Cave H, van der Werff ten Bosch J, Suciú S, et al. Clinical significance of minimal residual disease in childhood acute lymphoblastic leukemia. European Organization for Research and Treatment of Cancer-Childhood Leukemia Cooperative Group. N Engl J Med. 1998; 339 (9): 591-8.
- Chen CL, Liu O, Pui CH, Rivera GK, et al. Higher frequency of glutathione S-transferase deletions in black children with acute lymphoblastic leukemia. Blood. 1997; 89 (5): 1701-7.
- Cheok MH, Evans WE, et al. Acute lymphoblastic leukaemia: a model for the pharmacogenomics of cancer therapy. Nat Rev Cancer. 2006; 6 (2): 117-29.
- Chin-Shih F, Ruey Wu, Shieh J, et al. Pesticide exposure on southwestern Taiwanese with *MnSOD* and *NQO1* polymorphisms is associated with increased risk of Parkinson's disease. Clinica Chimica Acta. 2007; 378: 136-141.
- Coustan-Smith E, Behm FG, Sanchez J, et al. Immunological detection of minimal residual disease in children with acute lymphoblastic leukaemia. Lancet. 1998; 351 (9102): 550-4.
- Daniel W, Nebert, Amy L, et al. *NQO1* polymorphism, exposure to benzene, and predisposition to disease: Huge review. Genetic in medicine. 2002; 4: 62-70.
- DeAnn J. Liska. The Detoxification Enzyme Systems. Alternative Medicine Review. 1998; 3: 187-196.
- Edward S, Andrew L, Mel F, et al. Childhood acute lymphoblastic leukemia. Leukemia 7th ed. 2002; 601-620.
- Eguchi-Ishimae M, Eguchi M, Ishii E, et al. The association of a distinctive allele of NAD(P)H:quinone oxidoreductase with pediatric acute lymphoblastic leukemias with MLL fusion genes in Japan. Haematologica. 2005; 90 (11): 1511-1515.
- Errico A, Taioli E, Chen X, Vineis P, et al. Genetic metabolic polymorphisms and the risk of cancer: A review of the literature. Biomarkers. 1996; 1: 149-173.

- Eyada TK, El Ghonemy EG, El Ghoroury EA, et al. Study of genetic polymorphism of xenobiotic enzymes in acute leukemia. Blood Coagul Fibrinolysis. 2007; 18 (5): 489-495.
- Francesco D, Teresa M, Guidi F, et al. Polymorphism of *CYP1A1* and glutathione S-transferase and susceptibility to adult acute myeloid leukemia. Hematologica. 2004; 89: 664-670.
- Gaynon PS. Childhood acute lymphoblastic leukaemia and relapse. Br J Haematol. 2005; 131 (5): 579-587.
- GHyun-Jung Cho, Soo-Youn Lee, Chang-Seok Ki, et al. *GSTM1*, *GSTT1* and *GSTP1* Polymorphisms in the Korean Population. J Korean Med Sci. 2005; 20: 1089-1092.
- Giordano P, Santoro N, Del Vecchio GC, et al. T-immunophenotype is associated with an increased prevalence of thrombosis in children with acute lymphoblastic leukemia. A retrospective study. Haematologica. 2003; 88 (9): 1079-1080.
- Greaves M. Etiology of acute leukemia. Lancet. 1997; 349 (9048): 344-349.
- Greaves M. Molecular Genetics, Natural History and the Demise of Childhood Leukaemia. European Journal of Cancer. 1999; 35 (2): 173-185.
- Guengerich FP, Shimada T. Oxidation of toxic and carcinogenic chemicals by human cytochrome P-450 enzymes. Chem Res Toxicol. 1991; 4(4): 391-407.
- Helfrich, LA, Weigmann, DL, Hipkins, P, et al. Pesticides and aquatic animals: A guide to reducing impacts on aquatic systems. Virginia Cooperative Extension. 2007; 4: 443-456.
- Joseph T, Kusumakumary P, Chacko T, et al. Genetic polymorphism of *CYP1A1*, *CYP2D6*, *GSTM1* and *GSTT1* and susceptibility to acute lymphoblastic leukemia in Indian children. Pediatric Blood Cancer. 2004; 43 (5): 560-567.
- Kadlubar FF, Hammons GJ, et al. Role of cytochrome P-450 in metabolism of chemical carcinogens. In: Guengerich FP, ed. Mammalian Cytochromes P-450. Boca Raton, FL: CRC Press. 1987; 81-130.

- Katoh T, Nagata N, et al. Glutathione-S-transferase M1 and T1 genetic polymorphism and susceptibility to gastric and colorectal adenocarcinoma. Carcinogenesis. 1996; 9: 1855-1865.
- Kaufman D, Issaranisil S, Anderson T, et al. Use of Household Pesticides and the Risk of Aplastic Anaemia in Thailand. International Journal of Epidemiology. 1997; 26: 3: 643-650.
- Khuhaprema T, Srivatanakul P, Sriprang H, et al. Childhood cancer. Cancer in Thailand. 2007; 4: 73-78.
- Kishi R, Katakura Y, et al. Association of parents' occupational exposure to cancer in childhood. A case-control study of acute lymphoblastic leukemia. Sangyo Igaku. 1993; 35(6):515-529.
- Kori B, Jane A, Charles F, et al. Cancer Risk and Parental Pesticide Application in Children of Agricultural Health Study Participants. Toxicology and Applied Pharmacology. 2000; 354: 1400-1413.
- Krajinovic M, Labuda D, Richer C, Karimi S, Sinnett D, et al. Susceptibility to childhood acute lymphoblastic leukemia: influence of *CYP1A1*, *CYP2D6*, *GSTM1*, and *GSTT1* genetic polymorphisms. Blood. 1999; 93 (5): 1496-1501.
- Look AT. Oncogenic transcription factors in the human acute leukemia. Science. 1997; 278: 1059-1064.
- Ma X, Buffler PA, Selvin S, Matthay KK, et al. Daycare attendance and risk of childhood acute lymphoblastic leukaemia. Br J Cancer. 2002; 86 (9): 1419-1424.
- Maja Krajinovic, Damian Labuda, Chantal Richer, et al. Susceptibility to Childhood Acute Lymphoblastic Leukemia: Influence of *CYP1A1*, *CYP2D6*, *GSTM1*, and *GSTT1* Genetic Polymorphisms. Blood. 1999; 93 (5): 1496-1501.
- Margolin JF, Steuber CP, Poplack DG, et al. Acute lymphoblastic leukemia. In: Principles and Practice of Pediatric Oncology 15th ed. 2006: 538-590.
- Martin T. Smith, Cliona M. Mchale, Joseph L. Wiemels, et al. Molecular biomarkers for the study of childhood leukemia. Toxicology and Applied Pharmacology. 2005; 206: 237-245.
- McNally RJ, Parker L, et al. Environmental factors and childhood acute leukemias and lymphomas. Leuk Lymphoma. 2006; 47 (4): 589-598.

- Meinert R, Schuz J, kaletsch U, kaatsch P, michaelis J, et al. Leukemia and Non-Hodgkin's Lymphoma in Childhood and Exposure to Pesticides: Results of a Register-based Case-Control Study in Germany. American Journal of Epidemiology. 2000; 151 (7): 639-646.
- Mimemori E, Mariko E, Eiichi I, et al. The associated of a distinctive allele of NAD(P)H:quinone oxidoreductase with pediatric acute lymphoblastic leukemias with MLL fusion genes in Japan. Hematologica. 2005; 90: 1511-1515.
- Mututes E, Morilla R, Farahat N, et al: Definition of biphenotypic leukemia. Hematologica. 1997; 82-64.
- Nylund SJ, Ruutu T, Saarinen U, et al. Detection of minimal residual disease using fluorescence DNA in situ hybridization: A follow-up study in leukemia and lymphoma patients. Leukemia. 1994; 8: 587-592.
- Pakakasama S, Mukda E, Sasamakul W, et al. Polymorphisms of drug-metabolizing enzymes and risk of childhood acute lymphoblastic leukemia. Am J Hematol. 2005; 79 (3): 202-205.
- Parl R, Vorachek WR, Xu SJ, et al. Role of glutathione-S-transferase and carcinogenesis. Cancer Lett. 2000; 221: 123-129.
- Pearson WR, Vorachek WR, Xu SJ, et al. Role of the deletion polymorphisms of the glutathione-S-transferase *GSTT1* and *GSTM1*. Am. J. Hum. Genet. 1993; 53 (1): 220-223.
- Pui CH, Evans WE, et al. Treatment of acute lymphoblastic leukemia. N Engl J Med. 2006; 354 (2): 166-178.
- Pui CH, Relling MV, Downing JR, et al. Acute lymphoblastic leukemia. N Engl J Med. 2004; 350 (15): 1535-1548.
- Rebbeck TR, et al. Molecular epidemiology of the human glutathione S-transferase genotypes *GSTM1* and *GSTT1* in cancer susceptibility. Cancer Epidemiol Biomarkers Prev. 1997; 6 (9): 733-743.
- Reynolds P, Von Behren J, Gunier R, et al. Agricultural pesticides and lymphoproliferative childhood cancer in California. Scand J Work Environ Health. 2005; 31 Suppl 1: 46-54.

- Rivard C, Labuda D, Krajinovic M, et al. Risk of childhood leukemia associated with exposure to pesticides and with gene polymorphisms. Epidermiology. 1999; 10 (5): 481-487.
- Ross ME, Zhou X, Song G, et al. Classification of pediatric acute lymphoblastic leukemia by gene expression profiling. Blood. 2003; 15;102(8): 2951-2959.
- Salah H, Sheweita A, Drug-Metabolizing Enzymes: Mechanisms and Functions. Current Drug Metabolism. 2000; 1: 107-132.
- Sirma S, Agaoglu L, Yildiz I, et al. NAD(P)H:quinone oxidoreductase 1 null genotype is not associated with pediatric de novo acute leukemia. Pediatr Blood Cancer. 2004; 43 (5): 568-570.
- Smith M, Ries L, Gurney J, Ross J, et al. Cancer Incidence and Survival among Children and Adolescents: United States SEER Program. National Cancer Institute. 1999; 17-34.
- Stacy A, Geisler, Andrew F, Olshan et al. *GSTM1*, *GSTT1*, and the risk of squamous cell carcinoma of the head and neck: A Mini-HuGe Review. American Journal of Epidemiology. 2001; 154: 95-105.
- Stefano Landi, et al. Mammalian class theta GST and differential susceptibility to carcinogen: a review. Mutation Research. 2000; 463: 247-283.
- Uckun FM, Kersey JH, Vallera DA, et al. Autologous bone marrow transplantation in high-risk remission T-lineage acute lymphoblastic leukemia using immunotoxin plus 4-hydroperoxycyclophosphamide for marrow purging. Blood. 1990; 76: 1723-1727.
- Vervoordeldonk SF, Merle PA, Behrendt H, et al. Tripical immunofluorescence staining for prediction of relapse in childhood precursor B cell lymphoblastic leukemia. Br J Hematol. 1996; 92: 922-927.
- Vineis P, Malats N, Lang M, et al. Metabolic polymorphisms and susceptibility to cancer. Lyon, France: IARC. 1999; 32: 23-29.
- Wang Z, Tang L, Sun G, et al. Etiological study of esophageal squamous cell carcinoma in an endemic region: a population-based case control study in Huaian, China. BMC Cancer. 2006; 15 (6): 287-292.

- Weigmann, DL, Hipkins, P, Stinson P, et al. Pesticides and aquatic animals: A guide to reducing impacts on aquatic systems. Virginia Cooperative Extension. 2007; 10-14.
- Wiemels J, Cazzaniga G, Daniotti M, et al. Prenatal Origin of Acute Lymphoblastic Leukaemia in Children. Lancet. 1999; 354: 1499-1503.
- Wiemels J, Leinard B, Wang Y, Segal M, et al. Site-specific translocation and evidence of post-natal origin of the t(1;19) *E2A-PBXq* translocation in childhood acute lymphoblastic leukemia. Proceedings of the National Academy of Sciences. 2002; 203-205.
- Wiemels J, Xiao Z, Buffler PA, Maia AT, Ma X, et al. In utero origin of t(8;21) *AML1-ETO* translocations in childhood acute myeloid leukemia. Blood. 2002; 99 (10): 3801-3815.
- Wiemels J, Pagnamenta A, Taylor GM, Eden OB, et al. A Lack of a Functional NAD(P)H: Quinone Oxidoreductase Allele Is Selectively Associated with Pediatric Leukemias That Have MLL Fusions. Cancer Research. 1999; 59: 4095-4099.
- William L, Deepa B, Cheryl L, Dong-Joon M, Marry R, et al. Pediatric Acute Lymphoblastic Leukemia. American Society of Hematology. 2003; 102-131.
- Xiao Z, Greaves MF, Buffler P, Smith MT, Segal MR, et al. Molecular characterization of genomic *AML1-ETO* fusions in childhood leukemia. Leukemia. 2001; 15 (12): 1906-1913.
- Xu S, Wang Y, Roe B, Pearson W, et al. *GST* and differential susceptibility to carcinogen. J. Biol. Chem. 1998; 273: 3517-3572.
- Zahm S H, Devesa S, et al. Childhood cancer: overview of incidence trends and environmental carcinogens. Environmental Health Perspective (supplement). 1995; 103 (Supplement6): 177-224.
- Zahm S H, Ward M H, et al. Pesticides and childhood cancer. Environ. Health. Perspect. 1998; 106 (suppl. 3): 893-908.

Zhong S, Ma X, Buffler PA, Smith MT, et al. Whole genome amplification increases the efficiency and validity of buccal cell genotyping in pediatric populations. Cancer Epidemiol Biomarkers Prev. 2001; 10 (6): 697-700.

APPENDICES

APPENDIX A
QUESTIONNAIRE

มกราคม 2550

ID code



การวิจัยปฏิสัมพันธ์ระหว่างพันธุศาสตร์ของยีน *GSTM1*, *GSTT1*, *CYP1A1*, *NQO1* และ *XRCC1* และการสัมผัสสารป้องกันกำจัดศัตรูพืชและควันบุหรี่ กับความเสี่ยงในการเกิดโรคมะเร็งเม็ดเลือดขาวชนิด acute lymphoblastic leukemia ในเด็กไทย

กรอกใบยินยอมด้วยความสมัครใจให้ทำการวิจัยในมนุษย์

ได้รับการเก็บเซลล์จากตัวอย่างเลือด

วันที่สัมภาษณ์ / /

สถานที่สัมภาษณ์

1 บ้าน 2 โรงพยาบาล 3 อื่นๆ _____

ส่วนที่ 1: ข้อมูลเบื้องต้น

1. ท่านคือมารดาผู้ให้กำเนิดของเด็กใช่หรือไม่?

1 ใช่

2 ไม่ใช่ → หยุดการสัมภาษณ์

3 ไม่ทราบ

2. วันเดือนปีเกิดของท่าน (มารดา) คือ / /

3. สถานะภาพการสมรสของมารดาคือ

1 แต่งงาน

4 หย่าร้าง

2 อยู่ด้วยกัน

6 ไม่เคยแต่งงาน

3 แยกกันอยู่

9 ไม่ทราบ

มกราคม 2550

ส่วนที่ 2: ข้อมูลการใช้สารป้องกันกำจัดศัตรูพืชในการเกษตร และสารกำจัดแมลงภายในบ้าน

คำถามเกี่ยวกับการใช้สารป้องกันกำจัดศัตรูพืช และสารกำจัดแมลง กรณียบอกชนิดหรือชื่อสารที่ท่านใช้และช่วงเวลาที่ใช้สารเคมีดังกล่าวตามช่วงระยะเวลาที่ได้กำหนดไว้ตามตาราง

1. ท่านเคยใช้สารกำจัดวัชพืช, สารกำจัดแมลง, สารป้องกันกำจัดโรคพืชหรือไม่ สารชื่ออะไร และเมื่อไร?

1 เคย 2 ไม่เคย 3 ไม่ทราบ

1) ชนิดของสารป้องกันกำจัดศัตรูพืช	2) ใครเป็นผู้ใช้สารเคมี	3) ช่วงเวลาที่ใช้สารเคมี					4) ใช้กี่ครั้ง/ช่วง					
		Trimester					อายุ					
		3 mo	1 st trim	2 nd trim	3 rd trim	BF	0-1	1-2	2-3	12 mo	≥ 5	≤ 5
1= สเปรย์ฉีดกันขุงและแมลง 2= ยาจุดกันขุง 3= สารทากันขุง 4= สารฆ่าปลวก 5= อื่นๆ	1 = มารดา 2 = บิดา 3 = เด็ก 7 = อื่นๆ : ระบุ _____	1 = 3 เดือนก่อนมารดาตั้งครรภ์ 2 = 3 เดือนแรกที่มารดาตั้งครรภ์ 3 = 3-6 เดือน 4 = 6-9 เดือน 5 = ช่วงมารดาให้นมบุตร 6 = เด็กแรกเกิดจนอายุ 1 ปี 7 = เด็กอายุ 1-2 ปี 8 = เด็กอายุ 2-3 ปี 9 = ช่วงก่อนสัมผัสภยันต์ 1 ปี					2 = ≥ 5 ครั้ง 3 = ≤ 5 ครั้ง					
		1	2	3	4	5	6	7	8	9	2	3
		1	2	3	4	5	6	7	8	9	2	3
		1	2	3	4	5	6	7	8	9	2	3
		1	2	3	4	5	6	7	8	9	2	3
		1	2	3	4	5	6	7	8	9	2	3
		1	2	3	4	5	6	7	8	9	2	3
		1	2	3	4	5	6	7	8	9	2	3
		1	2	3	4	5	6	7	8	9	2	3
		1	2	3	4	5	6	7	8	9	2	3
		3 mo	1 st trim	2 nd trim	3 rd trim	BF	0-1	1-2	2-3	12 mo	≥ 5	≤ 5

ส่วนที่ 3: ข้อมูลอาชีพของมารดา

1. ท่านเคยทำอาชีพดังต่อไปนี้หรือไม่?

1) ทำสวนผักหรือผลไม้	<input type="checkbox"/> 1 ใช่ → ช่วงระยะเวลาที่ทำงานปี พ.ศ. _____ ถึง _____ <input type="checkbox"/> 2 ไม่ใช่
2) ทำนา	<input type="checkbox"/> 1 ใช่ → ช่วงระยะเวลาที่ทำงานปี พ.ศ. _____ ถึง _____ <input type="checkbox"/> 2 ไม่ใช่
3) ทำไร่	<input type="checkbox"/> 1 ใช่ → ช่วงระยะเวลาที่ทำงานปี พ.ศ. _____ ถึง _____ <input type="checkbox"/> 2 ไม่ใช่
4) รับผิดชอบคนหรือดูแลสวน	<input type="checkbox"/> 1 ใช่ → ช่วงระยะเวลาที่ทำงานปี พ.ศ. _____ ถึง _____ <input type="checkbox"/> 2 ไม่ใช่
5) อาชีพที่เกี่ยวข้องกับการเกษตร อื่นๆ	<input type="checkbox"/> 1 ใช่ → ช่วงระยะเวลาที่ทำงานปี พ.ศ. _____ ถึง _____ <input type="checkbox"/> 2 ไม่ใช่

APPENDIX B

BUFFERS AND REAGENTS

1. 10% SDS solution

Sodium dodecyl sulfate	10	g
Distilled water to	100	ml

Mix the solution and store at room temperature

2. 0.5 M EDTA (pH 8.0)

Disodium ethylenedimine tetraacetate.2H ₂ O	186.6	g
--	-------	---

Dissolve in distilled water and adjust pH to 8.0 with NaOH

Distilled water to	1,000	ml
--------------------	-------	----

Sterilize the solution by autoclaving and store at room temperature

3. 50X Tris acetate buffer (50X TAE buffer)

Tris base	242	g
Glacial acetic	57.1	ml
0.5 M EDTA pH 8.0	100	ml
Distilled water to	1,000	ml

4. 6X loading dye

Bromphenol blue	0.25	g
Xylene cyanol	0.25	g
Glycerol	50	ml
1 M Tris (pH 8.0)	1	ml
Distilled water to	100	ml

Mix and stored at 4°C

5. 2% agarose gel (w/v)

Agarose	1.6	g
1X TAE buffer	80	ml

Dissolve by heating in microwave oven and mix occasionally until no granules of agarose are visible

6. Ethyidium bromide

Ethyidium bromide	10	mg
Distilled water	1	ml

Mix the solution and store at 4°C

7. Denature solution

NaOH 0.5 M	10	g
NaCl 1.5 M	43.8	g
Distilled water to	500	ml

Mix the solution and store at room temperature

8. Neutralizing solution

NaCl 1.5 M	43.8	g
Tris base 1M pH 7.2	60.5	g
Distilled water to	500	ml

Mix the solution and store at room temperature

APPENDIX C

CHEMICAL AGENTS AND INSTRUMENTS

Materials

1. Pipette tip: 10 μ l, 200 μ l, 1000 μ l (Axygen Scientific, USA and Euro Lab[®] Labortechnik KG, German)
2. Microcentrifuge tube: 0.2 ml, 0.5 ml, 4.5 ml (Axygen Scientific, USA)
3. Polypropylene conical tube: 15 ml, 50 ml (Corning, USA)
4. Beaker: 50 ml, 100 ml, 200 ml, 500 ml, 1,000 ml (Pyrex)
5. Flask: 250 ml, 500 ml, 1,000 ml (Pyrex)
6. Reagent bottle: 100 ml, 250 ml, 500 ml, 1,000 ml (Duran, USA)
7. Cylinder: 25 ml, 50 ml, 100 ml, 250 ml, 500 ml, 1,000 ml (Witeg, German)
8. Glass pipette: 5 ml, 10 ml (Witeg, German)
9. Pipette rack (Eppendorf, German)
10. Thermometer (Precision, German)
11. PARAFILM (American National Can, USA)
12. Plastic wrap
13. Aluminum foil
14. Stirring-magnetic bar
15. Combs (Bio-RAD)
16. Electrophoresis chamber set (Bio-RAD)
17. Timer (Canon, China)

Equipment

1. Automatic adjustable micropipette: P2 (0.1-2.5 μ l), P10 (0.5-10 μ l), P20 (2-20 μ l), P100 (10-100 μ l), P200 (10-200 μ l), P1000 (0.1-1 ml) (Eppendorf, German)
2. Pipette boy (Tecnomara, Switzerland)
3. Vertex (Scientific Industry, USA)

4. pH meter (Ecomet, UK)
5. Stirring hot plate (Corning, USA)
6. Microcentrifuge (Eppendorf, German)
7. Thermal centrifuge (Heraeus, German)
8. DNA Thermal cycler 480 (Applied Biosystems, USA)
9. Heat block (Boekel Scientific, UK)
10. Shaker (Armed)
11. Incubator (Mettler, German)
12. Spectrophotometers (Bio-RAD, USA)
13. Gel-doc (Bio-RAD, USA)
14. Refrigerator 4 °C (Hitachi, Sanyo, Japan)
15. Deep freeze -20 °C, -80 °C (Sanyo, Japan)
16. Water Bath

Reagents

1. General reagents
 - 1.1 Hydrochloric acid (Merck)
 - 1.2 Sodium chloride (Merck)
 - 1.3 Tris base (USB)

2. Reagents for PCR analysis
 - 2.1 10X PCR buffer (500 mM KCl, 200 m tris-HCl pH 8.4)
 - 2.2 Magnesium Chloride
 - 2.3 Deoxynucleotide triphosphates (dNTPs)
 - 2.4 Oligonucleotide primers (BSU)
 - 2.5 Taq DNA polymerase (Fermentus)

3. Reagents for electrophoresis

3.1 Agarose, molecular grade (Promega)

3.2 Ethidium bromide (Sigma, USA)

3.3 6X loading dye

3.4 50X TAE

3.5 100 base pair DNA ladder (Fermentas)

BIOGRAPHY

Personal data

Full name Miss Pattramon Aungbamnet
Institutional Affiliation Hematology Laboratory Department of Pediatric
Chulalongkorn University, Bangkok, Thailand
Birth date and Place 10/11/1975; Bangkok, Thailand

Education and Professional Training

1993-1996	Chulalongkorn University Bangkok, Thailand	B.Sc. (Medical Technology)
2006-2008	Chulalongkorn University Bangkok, Thailand	Visiting Graduate. (Medical Science)