REFERENCES

- 1. Kurer FL, Welch DB. Gynecological laparoscopy: Clinical experiences of two anesthetic techniques. Br.J.Anaesth.1984;56:1207-1211.
- 2. Hodgson C, Mc Clelland RMA, Nuwton JR. Some effects of the peritoneal insufflation of carbon dioxide at laparoscopy.

 Anaesthesia. 1970;25:382-390.
- 3. Desmond J, Gordon RA. Ventilation in patients anesthetised for laparoscopy. Can. Anaesth. Soc. J. 1970;17:378-387.
- 4. Eger EI. The pharmacology of isoflurane. Br.J.Anaesth.1984;56:71s-98s.
- 5. Pandit UA, Steude GM, Leach AB. Induction and recovery characteristics of isoflurane and halothane anaesthesia for short outpatient operations in children. Anaesthesia.1985;40:1226-1230.
- 6. Fisher DM, Robinson S, Brett CM, Perin G, Gregory GA. Comparison of enflurane, halothane, and isoflurane for diagnostic and therapeutic procedures in children with malignancies. Anesthesiology 1985; 63:647-650
- 7. Carter JA, Dye AM, Cooper GM. Recovery from day-case anaesthesia.
 Anaesthesia 1985;40:545-548.
- 8. Kingston HGG. Halothane and isoflurane anesthesia in pediatric outpatients. Anesth. Analg. 1986;65:181-184.
- 9. Cattermole RW, Verghese C, Blair IJ, Jones CJH, Flynn PJ, Sebel PS.

 Isoflurane and halothane for outpatient dental anaesthesia in children. Br.J.Anaesth.1986; 58:385-389.
- 10. McAteer PM, Carter JA, Cooper GM, Prys Robert C. Comparison of isoflurane and halothane in outpatient paediatric dental anaesthesia. Br.J.Anaesth.1986;58: 390-393.
- 11. Ghaly RG, Flynn RJ, Moore J. Isoflurane as an alternative to halothane for Caesarean section. Anaesthesia. 1988; 43:5-7.


- 12. Wren WS, McShane AJ, McCarthy JG, Lamont BJ, Casey WF, Hannon VM. Isoflurane in paediatric anaesthesia. Anaesthesia. 1985;40: 315-323.
- 13. Milligan KR, Howe JP, Dundee JW. Halothane and isoflurane in outpatient anaesthesia. Anaesthesia. 1988;43:2-4.
- 14. Steinberg H. Selective effects of an anesthetic drug on cognitive behavior. Quarterly Journal of Experimental Psychology.1954;6: 170-180.
- 15. Chung. F, Cheng DCH, Seyone C. A randomized comparison of midazolam and diazepam injectable emulsion in cataract surgery. Can. J. Anaesth.1990;37:528-533.
- 16. Ghouri AF, Bodner M, White PF. Recovery profile after desflurane nitrous oxide versus isoflurane - nitrous oxide in outpatients. Anesthesiology.1991;74:419-424.
- 17. Korttila K. Recovery period and discharge. In: White PF. ed.
 Outpatient Anaesthesia. New York: Churchill Livingstone, Inc.
 1990;369-395.
- 18. Van SA. Special consideration for gynaecology and obstetrics.
 In: Kallar SK, Whitwam JG ed. Outpatient Anesthesia. Bussum:
 Medicom Europe 1988;85-88.
- 19. Boerner TF, Bartkowski RR, Goldberg ME, Torjman M, Odorisio L,
 DeLuca P. Cognitive recovery after prolong alfentanyl infusion:
 Midazolam and diazepam premedication compared. Anesth. Anal.
 1990;70:S1-S450.
- 20. Tsai M, Lee C, Kwan WF, Chen BJ. Recovery of cognitive functions after anaesthesia with desflurane or isoflurane and nitrous oxide.

 Br.J.Anaesth. 1992;69:255-258.

- 21. Milligan KR, O' Toole DP, Howe JP, Cooper JC, Dundee JW. A comparison of incremental propofol with propofol -isoflurane for outpatient gynecological surgery. Br. J. Anaesth. 1987;59: (in press).
- 22. Henegan C, McAuliffe R, Thomas D, Radford P. Morbidity after outpatient anaesthesia. Anaesthesia. 1981;36:4-8
- 23. Fragen RJ, Shanks CA. Neuromuscular recovery after laparoscopy.

 Anesth. Analg.1984;63:51-54.
- 24. Sosis M, Goldberg ME, Marr AT. Atracurium and vecuronium do not affect extraocular muscle function after outpatient surgery.

 Anesthesiology. 1988;68:465-468.
- 25. MacKenzie N, Grant IS. Comparison of the new emulsion formulation of propofol with methohexitone and thiopentone for induction of anaesthesia in day cases. Br. J. Anaesth.1985;57:725-731.
- 26. Carpenter RL, Eger EI, Johnson BH. Does the duration of anesthetic administration affect the pharmacokinetics or metabolism of inhaled anesthetics in humans ? Aneath Analg. 1987;66:1-5.



Appendix I

Preoperative Classification of Patients

according to the American Society of Anesthesiologists

Class I = A normal healthy patient other than surgical

pathology without systemic disease.

Class II = A patient with mild systemic disease - no functional limitations

Class III = A patient with moderate systemic disease

Class IV = A patient with severe systemic disease

Class V = Moribund

E = Emergency

Appendix II

Postoperative questionnaire

An attempt is being made to assess the effect of an anesthesia on patients in Siriraj Hospital. Your reply will be treated confidentially. Please answer this questionnaire.

| Instruction | : please check the correct answer. |
|-------------|---|
| 1. | Did you have any of the following after anesthesia ? |
| (can choose | more than 1 answer) |
| | [] nausea |
| | [] vomiting |
| | [] dizziness |
| | [] headache |
| | [] sore throat |
| | [] muscle pain |
| | [] pain at the injection site |
| | [] others, specify |
| ٤. | Did you have awareness during the operation ? |
| | [] yes [] no |
| 3. | Over all, how did you rate the anesthesia you had received? |
| | [] very good |
| | [] good |
| | [] satisfactory |
| | [] poor |
| | [] very poor |

| 4. | Should it | be required, | would you | like to ha | ve a similar |
|------------|-----------|--------------|-----------|------------|---------------------|
| anesthesia | again ? | | | | |
| | [] yes, | because | | | • • • • • • • • • • |
| | [] no, | because | | | |

แบบสอบถาม (หลังดมฮาสลบ)

No.

| กร | ภูณาใส่เครื่องหมาย / หน้าคำตอบที่ท่านเลือก หรือเติมคำตอบลงในช่องว่าง |
|----|--|
| 1. | ผลข้างเคียงจากการคมยาสลบในครั้งนี้ |
| | ท่านมีอาการต่าง ๆ เหล่านี้หรือไม่ (ตอบได้มากกว่า 1 คำตอบ) |
| | ย ว คลื่นไส้ |
| | ธ ว อาเจียน |
| | E] รู้สึกมินงง, สับสน |
| | [] ปวดศีรษะ |
| | ย 1 เจ็บคอ |
| | [] ปวดเมื่อยกล้ามเนื้อ |
| | E] เจ็บบริเวณที่ฉีดยา |
| | [] เจ็บแผลผ่าตัดมากจนต้องการยาฉีดแก้ปวด |
| | [] อื่น ๆ โปรดระบุ |
| 2. | ขณะผ่าตัดท่านรู้สึกตัวหรือไม่ |
| | E] รู้สึก E] ไม่รู้สึก ท่านรู้สึกต่อการดมยาสลบที่ท่านได้รับครั้งนี้อย่างไร |
| 3. | ท่านรู้สึกต่อการดมยาสลบที่ท่านได้รับครั้งนี้อย่างไร |
| | [] พอใจมากที่สุด |
| | [] พอใจมาก |
| | [] wala |
| | E] Himala |
| | [] ไม่พอใจมากที่สุด |

Figure I The Ball Bearing Test



Ball bearing test. The subject is asked to transfer ball bearings from the glass dish to the tube as quickly as possible using a pair of forceps. The number transferred in 40 seconds is scored.

Figure II The Digit Symbol Substitution Test.

| DIGIT SYMBOL SUBSTITUTION TEST | | | | | - | TIME | | | | DATE | | | | ACTUAL TIME | | | | | | | | | | |
|--------------------------------|------|------|-------|-----|-----|------|------|------|------|------|---|--------|---|---------------|---|-----|----|-------|----|---|---|------|-----|---|
| | DIG. | LI K | J1111 | JOL | Joi | | .101 | .101 | . 11 | | | | | | | MO. | DA | AY Y | Æ. | v | | | AM. | |
| OIG] | | C | 1 | 2 | - | 3 | | | 5 | | 6 | 7 U | | 8 X | - | 9 | | | | | S | SCOF | E | |
| | | | | | | | | | | | | | | | | | | | | | | | | |
| 6 | 7 | 6 | 9 | 4 | 2 | 5 | 4 | 1 | 2 | 6 | 9 | 6 | 9 | 3 | 7 | 5 | 8 | 3 | 2 | 8 | 3 | 9 | 4 | 5 |
| 2 | 7 | 4 | 3 | 5 | 8 | 7 | 1 | 6 | 5 | 9 | 6 | 2 | 3 | 8 | 7 | 9 | 6 | 4 | 3 | 7 | 9 | 1 | 6 | 9 |
| | | | | | | | | | | | | | | | | | | | | | | | | |
| 4 | 7 | 4 | 1 | 5 | 4 | 3 | 7 | 3 | 9 | 3 | 8 | 1 | 3 | 1 | 6 | 3 | 4 | 2 | 6 | 2 | 6 | 8 | 6 | 8 |
| | | | | | | | | | | | | | | | | | | Asset | | | | | | |

The subject's task is to place the appropriate symbol below each number. The score is the number of correct responses completed in, for example, 90 seconds. In order to eliminate the training effect, different symbols are used when the test is repeated.

VITAE

Mr. Supoj Samattachai was born on 8 October, 1960 in Bangkok, Thailand. He graduated Medical Doctor from Siriraj Hospital, Mahidol University, in 1985. In 1989, he got Thai Board of Anesthesiology. He has been enrolled in the Master Degree of Science in Health Development at Faculty of Medicine, Chulalongkorn University since 1994. The present position is Anesthesiologist at Bhumibol Adulyadej Hospital, RTAF.

