

## REFERENCES

- สาธารณสุข, กระทรวง. ประกาศกระทรวงสาธารณสุข ฉบับที่ 97 (พ.ศ. 2539) เรื่องระบุชื่อและจัดแบ่งประเภทวัตถุออกฤทธิ์ ตามความในพระราชบัญญัติวัตถุออกฤทธิ์ต่อจิตและประสาท พ.ศ. 2518. กรุงเทพมหานคร: กระทรวงสาธารณสุข, 2539. (อัดสำเนา)
- Abad, A., and Montoya, A. 1994. Production of monoclonal antibodies for carbaryl from a hapten preserving the carbamate group. J. Agric. Food. Chem. 42: 1818-1823.
- Abad, A., Moreno, M.J., and Montoya, A. 1998. Hapten synthesis and production of monoclonal antibodies to the N-methylcarbamate pesticide methiocarb. J. Agric. Food. Chem. 46: 2417-2426.
- Abuknesha, R.A., and Maragkou, A. 2004. A high sensitive and specific enzyme immunoassay for detection of  $\beta$ -amanitin in biological fluids. Anal. Clin. Chem. 379: 853-860.
- Andrews, K.M. 1995. Ephedra's role as a precursor in the clandestine manufacture of methamphetamine. J. Forensic. Sci. 40(4): 551-560.
- Aoki, K., Hirose, Y and Kuroiwa, Y. 1990. Immunoassay for methamphetamine with a new antibody. J. Forensic. Sci. 44: 245-255.
- Aoki, A., and Kuroiwa, Y. 1985. A screening method for urinary methamphetamine latex agglutination inhibition reaction test. Forensic. Sci. Int. 27: 49-56.
- Armbruster, D.A., and Krolak, J.M. 1992. Screening for drugs of abuse with the Roche ONTRAK assays. J. Anal. Toxicol. 16(3):172-5.
- Badia, R., dela Torre, R., Corcione, S., and Segura, J. 1998. Analytical approach of European Union laboratories to drug of abuse analysis. Drug. Monitor. Toxicol. 44(4): 790-799.
- Barrett, J.T., ed. Textbook of Immunology. 5<sup>th</sup> Ed, An introduction to Immunochemistry and immunobiology. pp. 48-52. Washington: The C.V. Mosby, 1974.
- Batteiger, B., Newhall, V., and Jones, R.B. 1982. The use of Tween 20 as blocking agent in the immunological detection of proteins transferred to nitrocellulose membranes. J. Immunol. Methods 55: 297-307.

- Bruun, L., Koch, C., Jakobsen, M.H., Pedersen, B., Christiansen, M., and Jens, A. 2001. Characterization of monoclonal antibodies raised against different structures belonging to the s-triazine group of herbicides. Anal. Chim. Acta. 436: 87-101.
- Bureau of Justice Assistance. 1992. Drug Testing in the Criminal Justice System. Office of Justice Programs. Bureau of Justice Statistics. U.S. Department of Justice.
- Caraway, K.L., and Koshland, Jr., D.E. 1972. Carbodiimide modification of proteins. Methods Enzymol. 25: 616-623.
- Catty, D., and Raykundalia, C. Production and quality control of polyclonal antibodies. In D. Catty (ed.), Antibodies. volume I: A practical approach, pp.19-80. Oxford: IRL press, 1988.
- Chappey, O., Debray, M., Niel, E., and Scherrmann, J.M. 1994. Association constants of monoclonal antibodies for haptens: heterogeneity of frequency distribution and possible relationship with haptens molecular weight. J. Immunol. Methods 172: 219-225.
- Cheng, L.T., Kim, S.Y., Chung, A., and Castro, A. 1973. Amphetamine: New radioimmunoassay. FEBS LETTERS 36(3): 339-342.
- Chester, N., Molttram, D.R., Reilly, T., and Powell, M. 2003. Elimination of ephedrine in urine following multiple dosing: the consequences for athletes, in relation to doping control. British J. Clin. Pharmacol. 57(1): 62-67.
- Choi, M.J., Gorovits, B.M., Choi, J., Song, K.S.N., and Park, J. 1994. A visual membrane immunoassay for the detection of methamphetamine using an enzyme-labeled tracer derived from methamphetamine and amphetamine. Biol. Pharm. Bull. 17(7): 875-880.
- Ciabattoni, G. Production of antisera by conventional techniques. In C. Patrono; and B.A. Pedskar (eds.), Radioimmunoassay in basic and clinical pharmacology. pp.23-69. Berlin: Springer Verlag, 1987.
- Cleeland, R., Christensen, J., Usatequi-Gomez, M., Heveran, J., Davis, R., and Grunberg, E. 1976. Detection of drugs of abuse by radioimmunoassay: A summary data and some new information. Clin. Chem. 22(6): 712-725.

- Cody, J.T. 1990. Detection of D,L-amphetamine , D,L-methamphetamine and illicit amphetamine analogs using diagnostic product corporation's amphetamine and methamphetamine radioimmunoassay. J. Anal. Toxicol. 14: 321-324.
- Colbert, D.L., Gallacher, G., and Mainwaring-Burton, R.W. 1985. Single-reagent polarization fluoroimmunoassay for amphetamine in urine. Clin. Chem. 31(7): 1193-1195.
- Cremer, N.E., Cossen, C.K., Hanson, C.V., and Shell, G.R. 1982. Evaluation and reporting of enzyme immunoassay determinations of antibody to herpes simplex virus in sera and cerebrospinal fluid. J. Clin. Microbiol. 15(5): 815-823.
- Dallakian, P., Budzkiewicz, H., and Brzezinka, H. 1996. Detection and quantitation of amphetamine and methamphetamine : Electron impact and chemical ionization with ammonia-comparative investigation on shimadzu QP 5000 GC-MS system. J. Anal. Toxicol. 20(3): 255-261.
- Danilova, N.D. 1994. ELISA screening of monoclonal antibodies to hapten: influence of the chemical structure of hapten-protein conjugates. J. Immunol. Methods. 173: 111-117.
- Danilova, N.D., and Vasilov, R.G. 1991. Production and characterization of anti-theophylline monoclonal antibodies suitable for immunoassay. Immunology Letters 28: 79-84.
- de la Torre, R., et al. 1996. Cross-reactivity of stimulants found in sports drug testing by two fluorescence polarization immunoassays. J. Anal. Toxicol. 20(3): 165-170.
- De Mey, J.R. The preparation of immunoglobulin gold conjugates (IGS Reagents) and their use as markers for light and electron microscopic immunochemistry. In A.C. Cuello (ed.), Immunochemistry. IBRO, 1983.
- DePace, A., Verebey, K., and elSohly, M. 1990. Capillary gas-liquid chromatography separation of phenethylamines in amphetamine-positive urine samples. J. Forensic. Sci. 35(6): 1431-1435.

- Deschamps, R.J.A., Hall, J.C., and McDermott, M.R. 1990. Polyclonal and monoclonal enzyme immunoassay for picloram detection in water, soil, plants, and urine. J. Agric. Food. Chem. 38: 1881-1886.
- D'Nicuola, J., Jones, R., Levine, B., and Smith, M. 1992. Evaluation of six commercial amphetamine and methamphetamine immunoassays for cross-reactivity to phenylpropanolamine and ephedrine in urine. J. Anal. Toxicol. 16: 211-213.
- Dumestre-Toulet, V., and Kintz, P. 1999. Ephedrine abuse for doping purposes as demonstrated by hair analysis. J. Anal. Toxicol. 24(5): 381-382.
- Ekins, R.P. 1974. Basic principles and theory. Br. Med. Bull. 30: 3-11.
- Eremin, S.A., Gallacher, H., Smith, D.S., and Landon, J. 1987. Single-reagent polarization fluoroimmunoassay of methamphetamine in urine. Clin. Chem. 33(10): 1903-1906.
- Erlanger, B.F., Borex, F., Beiser, S.M., and Lieberman, S. 1957. Steroid-protein conjugates: I Preparation and characterization of conjugates of bovine serum albumin with testosterone and with cortisone. J. Biol. Chem. 228: 713-727.
- Galve, R., Sanchez-Baeza, F., Camps, F., and Marco, M.P. 2002. Indirect competitive immunoassay for trichlorophenol determination rational evaluation of the competitor heterology effect. Anal. Chim. Acta. 452: 191-206.
- Geoghegan, W.D. 1988. Immunoassays at the microscopic level: Solid-phase colloidal gold methods. J. Clin. Immunoassay 11(1): 11-23.
- Geoghegan, W.D., and Ackerman, G.A. 1977. Adsorption of horseradish peroxidase, ovomucoid and antiimmunoglobulin to colloidal gold for the indirect detection of concanavalin A, wheat germ agglutinin and goat antihuman immunoglobulin G on cell surfaces at the electron microscope level : a new method, theory and application. J. Histochem. Cytochem. 5(11): 1187-1200.
- Geoghegan, W.D., Ammbegaonker, S., and Calvanico, N.J. 1980. Passive gold agglutination. An alternative to passive hemagglutination. J. Immunol. Methods 34: 11-21.
- Goding, J. W. Monoclonal Antibodies : Principle and Practice. 3<sup>rd</sup>ed.  
London: Academic press, 1996.

- Goodman, S.L., Hodges, G.M., and Livingston, D.C. 1980. A review of the colloidal gold marker system. Scan. Elect. Microsc. 2: 133-144.
- Goodrow, M.H., and Hammock, B.D. 1998. Hapten design for compound-selective antibodies: ELISAS for environmentally deleterious small molecules. Anal. Chim. Acta. 376: 83-91.
- Goodrow, M.H., Harrison, R.O., and Hammock, B.D. 1990. Hapten synthesis, antibody development, and competitive inhibition enzyme immunoassay for s-triazine herbicides. J. Agric. Food. Chem. 38: 990-996.
- Gribnau, T.C.J., Leuvering, J.H.W., and Van Hell, H. 1986. Particle-labelled immunoassay: A review. J. Chromato. 376: 175-189.
- Grinstead, G.F. 1989. Ranitidine and high concentrations of phenylpropanolamine cross-react in the EMIT Monoclonal Amphetamine/Methamphetamine assay. Clin. Chem. 35: 1998-1999.
- Guidance for Prescription Use, Drugs of Abuse Assays: Premarket Notifications Devices for Drugs of Abuse Assays Using Various Methodologies, August 31, 1995. U.S. Department of Health and Human Services, Food and Drug Administration, Center for Devices and Radiological Health, Clinical Chemistry and Toxicology Branch. Division of Clinical Laboratory Devices, Office of Device Evaluation.
- Halbert, S.P., Bastomsky, C.H., and Anken, M. 1983. A rapid standardized enzyme immunoassay for autoantibodies to thyroglobulin. Clin. Chim. Acta. 127: 69-76.
- Handley, D.A., and Chien, S. 1983. Colloidal gold: a pluripotent receptor probe (41697). Proc. So. Exp. Biol. Med. 174: 1-11.
- Holthues, H., Pfeifer-Fukumura, U., Sound, I., and Baumann, W. 2005. Evaluation of the concept of heterology in a monoclonal antibody-based ELISA utilizing direct hapten linkage to polystyrene microtiter plates. J. Immunol. Methods 304: 68-77.
- Horisberger, M. 1977. Colloidal gold: A cytochemical marker for light and fluorescent microscopy and for transmission and scanning electron microscopy. Scan. Elect. Microsc. 2: 9-21.

- Horisberger, M. 1981. Colloidal gold: A cytochemical marker for light and fluorescent microscopy and for transmission and scanning electron microscopy  
Scanning. Electron. Microsc. 2: 9-31.
- Horisberger, M., and Rosset, J. 1977. Colloidal gold, a useful marker for transmission scanning electron microscope. J. Histochem. Cytochem. 25(4): 295-305.
- Hosoda, H., Kawamura N., and Nambara, T. 1981. Effect of bridge heterologous combination on sensitivity in enzyme immunoassay for cortisol.  
Chem. Pharm. Bull. 29(7): 1969-1974.
- Hosoda, H., Kobayashi, N., Kawamoto, H., and Nambara, T. 1983. The specificity of enzyme immunoassays for Plasma 11-Deoxycortisol. Chem. Pharm. Bull. 31(10): 3595-3600.
- Hosoda, H., Kobayashi, N., Ishii, N., and Nambara, T. 1985. The bridge length effect on sensitivity in steroid enzyme immunoassays. Chem. Pharm. Bull. 33(2): 902-904.
- Hosoda, H., Kobayashi, N., Ishii, N., and Nambara, T. 1986. Bridging phenomena in steroid immunoassays. The effect of bridge length effect on sensitivity in steroid enzyme immunoassays. Chem. Pharm. Bull. 34(5): 2105-2111.
- Hosoda, H., Kobayashi, N., and Nambara, T. 1983. Effect of bridge heterologous combination on sensitivity in enzyme immunoassay for 11-deoxycortisol.  
Chem. Pharm. Bull. 31(3): 953-958.
- Hosoda, H., Yoshida, H., Sakai, Y., Miyairi, S., and Nambara, T. 1980. Sensitivity and specificity in enzyme immunoassay of testosterone. Chem. Pharm. Bull. 28(10): 3035-3040.
- Immunochromatographic, lateral flow or strip tests development[Onlins]. 2000.  
Available from <http://www.pall.com/sls-4154.asp>[2004]
- Jenkins, P. 1999. Amphetamine drugs on the pacific rim. Available from  
<http://www.personal.psu.edu>[2006]
- Jeon, S.I., Yang, X., and Andrade, J.D. 2004. Modeling of homogeneous cloned enzyme donor immunoassay. Anal. Biochem. 333(1): 136-147.
- Johnstone, A., and Thrope, R. Immunochemistry in Practice. Oxford: Blackwell, 1982.

- Kelly, K.L. 1990. Ranitidine cross-reactivity in the EMIT<sup>®</sup> d.a.u.<sup>™</sup> monoclonal amphetamine / methamphetamine assay. Clin. Chem. 36(7): 1391-1392.
- Kemeny, D.M. 1992. Titration of antibodies. J. Immunol. Methods. 150: 57-76.
- Khosravi, M.J., and Papanastasiou-Diamandi, A. 1993. Hapten-heterologous conjugates evaluated for application to free thyroxine immunoassays. Clin. Chem. 39(2): 256-262.
- Kim, Y.J., Cho, Y.A., Lee, H-S., Lee, Y.T., Gee, S.J., and Hammock, B.D. 2003. Synthesis of haptens for immunoassay of organophosphorus pesticides and effect of heterology in hapten spacer arm length on immunoassay sensitivity. Anal. Chim. Acta. 475: 85-86.
- Killy, K.L. 1990. Ranitidine cross-reactivity in the EMIT<sup>®</sup> d.a.u. monoclonal amphetamine/methamphetamine assay. Clin. Chem. 36(7): 1391-1392.
- Kitjaroenham, A., Khusmith, S., Laothanvorn, J., Na-Bangchang, K., and Thammapalerd, N. 1998. Monoclonal antibodies to quinine. Southeast Asian J. Trop. Med. Public Health 29(2): 337-340.
- Kolb-Bachofen, V. 1989. Carbohydrate receptor binding using colloidal gold. Methods in Enzymology 179: 111-121.
- Kunsmann, G.W., et al. 1990. Application of the Syva EMIT and Abbott TD<sub>x</sub> Amphetamine immunoassays to the detection of 3,4-methylenedioxymethamphetamine (MADA) and 3,4-methylenedioxyethylamphetamine (MDEA) in Urine. J. Anal. Toxicol. 14: 321-324
- Landsteiner, K. ed. The specificity of serological reactions. Massachusetts: Harvard University Press, 1945.
- Lekskulchai, V., and Kaewpongsri, S. 2004. Reagents kits for urine methamphetamine: rapid test kits v.s. Abbott A/M II reagent. Thai J. Health Res. 18(1): 62-73.
- Lekskulchai, V., and Mokkhaveva, C. 2001. Evaluation of Roche Abuscreen ONLINE amphetamine immunoassay for screening of new amphetamine analogues. J. Anal. Toxicol. 25(6): 471-475.

- Leuvering, J.H.H.W., Hail, P.J.H.M., Van der Waart, M., and Schuurs., A.H.W.M. 1981. A sol particle agglutination for human chorionic gonadotrophin. J. Immunol. Methods. 45: 183-194.
- Leuvering, J.H.H.W., Hail, P.J.H.M., White, D.D., and Schuurs., A.H.W.M. 1988. A homogeneous sol particle immunoassay for total oestrogen in urine and serum samples. J. Immunol. Methods. 62: 163-174.
- Leuvering, J.H.H.W., Thal, P.J.H.M., and Schuurs., A.H.W.M. 1983. Optimization of a sandwich sol particle immunoassay for human chorionic gonadotrophin. J. Immunol. Methods. 62: 175-184.
- Liddell J. E., and Cryer, A. A practical guide to monoclonal antibodies. Newyork: John Wiley and Sons, 1991.
- Lokhnauth, J.K., and Nicholas, H.S. 2005. Solid phase micro-extraction coupled with ion mobility spectrometry for the analysis of ephedrine in urine. J. Sep. Sci. 28(7): 612–618.
- Loomans, E.E.M.G., Van. Wiltenburg, J., Koets, M., and Van. Amerongen, A. 2003. Neamin as immunogen for the development of a generic ELISA detecting gentamicin, kanamycin and neomycin in milk. J. Agric. Food. Chem. 51: 587-593.
- Loor, R., Lingenfelter, C., Wason, P.P., Tang, K., and Davoudzadeh, D. 2002. Multiplex assay of amphetamine, methamphetamine, and ecstasy drug using CEDIA<sup>®</sup> technology. J. Anal. Toxicol. 26(5): 267-273.
- Lucocq, J.M. and Baschong, W. 1986. Preparation of protein colloidal gold complexes in the presence of commonly used buffer. Eur. J. Cell Biol. 42: 332-337.
- Malik, V. S., and Lillehoj, E. P., eds. Antibody techniques. USA: Academic press, 1994.
- Malvano, R., Boniolo, A., DAVIS, M., and Zannino, M. 1982. ELISA for antibody measurement: aspect related to data expression. J. Immunol. Methods. 48(1): 51-60.
- Manclus, J.J., Primo, J., and Montoya, A. 1996. Development of enzyme-linked immunosorbent assays for the insecticide chlorpyrifos. 1. Monoclonal antibody production and immunoassay design. J. Agric. Food. Chem. 44: 4052-4062.





- Martin, J.M.C., Paques, M., Van der Velden-de Groot, T.A.M., and Beuvery, E.C. 1990. Characterization of antibody labeled colloidal gold particles and their application in a sol particle assay (SPIA). J. Immunoassay 11(1): 31-47.
- Matapatara, W. Selective immuno-detection of amphetamine and/or methamphetamine by principle of heterologous combinations. Master's Thesis, Department of Pharmaceutical Chemistry, Graduate School, Chulalongkorn University, 1999.
- Matapatara, W., and Thongnopnua, P. 2003. Effect of heterology on the detection of amphetamine, methamphetamine and ephedrine by enzyme-immunoassay. Thai J. Pharm. Sci. 27 (Suppl.).
- Matapatara, W., and Thongnopnua, P. Effect of heterology on the detection of amphetamine, methamphetamine and ephedrine by enzyme-immunoassay. The 20<sup>th</sup> Annual Research Meeting in Pharmaceutical Sciences. December 1, 2003, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok, 2003.
- Matapatara, W., Thongnopnua, P., and Lipipun, V. Comparison of Specificity and Sensitivity of Antibodies from Cell Culture Supernatant to Their Polyclonal Antibodies for Amphetamine, Methamphetamine and Ephedrine Detection. The 5<sup>th</sup> National Symposium on Graduate Research. pp.300. October 10-11, 2005. Bangkok, 2005a.
- Matapatara, W., Thongnopnua, P., and Lipipun, V. The Influence of Hybridoma Cell Medium on Specificity and Sensitivity of Antibody for Amphetamine, Methamphetamine and Ephedrine Detection. The 2<sup>nd</sup> AASP (Asian Associate of Schools of Pharmacy) Symposium/2<sup>nd</sup> APEM Conference 2005. pp.73. November 14-17, 2005. Bangkok, 2005b.
- Mathoera, R.B., Kok, D.J., Verduin, C.M., and Nijman, R.J.M. 2002. Pathological and therapeutic significance of cellular invasion by *Proteus mirabilis* in an enterocystoplasty infection stone model. Infect. Immun. 70(12): 7022-7032.
- Miagkova, M.A., Lushnikova, M.V., and Polevaia, O.A. 1990. Solid-phase immunoenzyme method for the quantitative determination of ephedrine. Sud. Med. Ekspert. 33(3): 30-32.

- Midha, K.K., Hubbard, J.K., Cooper, J.K., and Mackonka, C. 1983. Stereospecific radioimmunoassays for l-ephedrine and d-ephedrine in human plasma. J. Pharm. Sci. 72(7): 736-739.
- Miki, A., Katagi, M., Shima, N., and Tsuchihashi, H. 2004. Application of ORAL screen saliva drug test for the screening of methamphetamine, MDMA, and MDEA incorporated in hair. J. Anal. Toxicol. 28(2): 132-134.
- Mongkolsirichaikul, D., Tarnchompoo, B., and Ratanabanangkoon, K. 1993. Development of latex agglutination inhibition reaction test for amphetamine in urine. J. Immunol. Methods. 157: 189-195.
- Nakane, P.K., and Kawaoi, A. 1974. Peroxidase-labeled antibody a new method of conjugation. J. Histochem. Cytochem. 22(12): 1084-1091.
- Needleman, S.B., and Romberg, R.W. 1990. Limits of linearity and detection for some drugs of abuse. J. Anal. Toxicol. 14: 34-38.
- Nishikawa, T., Saito, M., and Kubo, H. 1984. Effect of heterologous combination on competitive nephelometric immunoassay II. Bridge-heterologous combination for ethosuximide immunoassay. Chem. Pharm. Bull. 32(12): 4958-4962 .
- Olsen, K.M., Gulliksen, M., and Christophersen, A.S. 1992. Metabolites of chlorpromazine and brompheniramine may cause false-positive urine amphetamine results with monoclonal EMIT<sup>®</sup> d.a.u. immunoassay. Clin. Chem. 38(4): 611-612.
- One Step AMP Test Kit.[Online] (n.d.).  
Available from: [www.passyourdrugtest.com](http://www.passyourdrugtest.com)[2005. Dec 17].
- Paek, S-H., Jang, M-R., Mok, R-S., Kim, S-C., and Kim, H-B. 1998. Immunochromatographic membrane strip assay system for a single-class plasma lipoprotein cholesterol, exemplified by high-density lipoprotein cholesterol measurement. Biotechnology Bioengineering 62(2): 145-154.
- Paek, S-H., Lee, S-H., Cho, J\_H., and Kim, Y-S. 2000. Development of rapid one-step immunochromatographic assay. Methods 22: 53-60.
- Paxton, J.W., Rowell, F.J., Ratcliffe, J.G. 1976. Production and characterization of antisera to diphenylidantoin suitable for radioimmunoassay. J. Immunol. Methods. 10: 317-327.

- Pidetcha, E., Congpnong, P., Putripraw, T., Rekakanakul, R., Suwanton, L., and Tantraronroj, S. 1995. Screening for urinary amphetamine in truck drivers and drug addicts. J. Med. Assoc. Thai. 78(10): 554-558.
- Piran, U., Riordan, W.J., and Silbert, D.R. 1990. Effect of hapten heterology on thyroid hormone immunoassays. J. Immunol. Methods. 133: 207-214.
- Piran, U., Silbert, S.D. and Barlow, E.H. 1993. Role of antibody valency in hapten-heterologous immunoassays. Clin. Chem. 39(5): 879-883.
- Pitchayawasin, S. Effect of heterologous combinations between enzyme labeled theophylline and immunogen on the sensitivity and specificity of enzyme immunoassay for theophylline. Master' s Thesis, Department of Pharmaceutical Chemistry, Graduate School, Chulalongkorn University, 1997.
- Poklis, A., Hall, K.V., and Still, J. 1991. Ranitidine interference with the monoclonal EMIT d.a.u. amphetamine/methamphetamine immunoassay. J. Anal. Toxicol. 15: 101-103.
- Poklis, A., and Moore, K.A. 1995. Stereoselectivity of the TDx ADx/FLx Amphetamine/Methamphetamine II amphetamine/methamphetamine immunoassay--response of urine specimens following nasal inhaler use. J. Toxicol. Clin. Toxicol. 33(1): 35-41.
- Pontecorvo, G. 1975. Production of mammalian somatic cells hybrids by means of polyethylene glycol treatment. Somatic Cell Genet. 1: 397-400.
- Porstmann, T., and Kiessig, S.T. 1992. Enzyme immunoassay techniques: An overview. J. Immunol. Methods. 150: 5-21.
- QuickPac II OneStep Amphetamine Test[Online] (n.d).  
Available from: [www.syntron.net](http://www.syntron.net)[2005. Dec 23]
- QuickScreen One-Step Rapid Amphetamine Test[Online] (n.d).  
Available from: [www.craigmedical.com](http://www.craigmedical.com)[2005. Dec 23]
- Richardson, M.D., Turner, A., Warnock, D.W., and Llewellyn, P.A. 1983. Computer-assited rapid enzyme-linked immunosorbent assay (ELISA) in the serological diagnosis of aspergillosis. J. Immunol. Method. 56: 201-207.

- Robinson, N. 2002. Assay development: immunogold conjugation for IVD applications [Online]. IVD Technology.  
Available from: [http://www.IVD applications \(IVDT archive, Mar 02\).htm\[2005\]](http://www.IVD applications (IVDT archive, Mar 02).htm[2005]).
- Rose, B.G., Kamps-Holtzapfle, C., and Stanker, L.H. 1995. Competitive indirect ELISA for ceftiofur sodium and the effect of different immunizing and coating antigen conjugates. Bioconjugate Chem. 6: 529-535.
- Roth, J. The colloidal gold marker system for light and electron microscopic cytochemistry. In G.R. Bullock; and P. Petrusz (eds.), Techniques in Immunocytochemistry. vol.2, pp. 217-278, New York: Academic Press, 1983.
- Ruangyuttikarn, K., and Moody, D.E. 1988. Comparison of three commercial amphetamine immunoassays for detection of methamphetamine , methylenedioxyethylamphetamine. J. Anal. Toxicol. 12: 229-233.
- Sato, H., Mochizuki, H., Tomita, Y., and Kanamori, T. 1996. Enhancement of the sensitivity of a chemiluminescent immunoassay for estradiol based on hapten heterology. Clin. Biochem. 29(6): 509-513.
- Schreier, M., Kohler, G., and Hengartner, H. 1980. Hybridoma techniques. New York: EMBO, 1980.
- Sengupta, J., Dhar, T.K., and Ali, E. 1992. Sandwich immunoassay of small molecules II effects of heterology and development of a highly sensitive and specific enzyme immunoassay for testosterone. J. Immunol. Method. 147: 181-188.
- Shyu, R-H., Shyu, H-F., Liu, H-W., and Tang, S-S. 2002. Colloidal gold-based immunochromatographic assay for detection of ricin. Toxicon 40: 255-258.
- Sirikatitham, A. Comparative immunodetection of ephedrine between enzyme-labeled and metal particle-labeled techniques. Master's Thesis, Department of Pharmaceutical Chemistry, Graduate School, Chulalongkorn University, 1999.
- Standerfer, J.C., and Saunders, G.C. 1978. Enzyme immunoassay for gentamicin. Clin. Chem. 24: 1903-1907.
- Steimer, K.S., and Boettiger, D. 1977. Complementation rescue of Rous sarcoma virus from transformed mammalian cells by polyethylene glycol-mediated cell fusion. J. Virol. 23(1): 133-141.

- Stout, P.R., Klette, K.L., and Horn, C.K. 2004. Evaluation of ephedrine, pseudoephedrine and phenylpropanolamine concentrations in human urine samples and a comparison of the specificity of DRI amphetamines and Abuscreen online (KIMS) amphetamines screening immunoassays. J. Forensic Sci. 49(1):160-164.
- Stout, P.R., Klette, K.L., and Wiegand, R. 2003. Comparison and Evaluation of DRI(r) Methamphetamine, DRI Ecstasy, Abuscreen(r) ONLINE Amphetamine, and a modified Abuscreen ONLINE amphetamine screening immunoassays for the detection of amphetamine (AMP), methamphetamine (MTH), 3,4-methylenedioxyamphetamine (MDA), and 3,4-methylenedioxymethamphetamine (MDMA) in human urine. J. Anal. Toxicol. 27(5): 265-269.
- Suttijitpasal, P., and Ratanabanangkorn, K. 1992. Immunoassays of amphetamines : immunogen structure & antibody specificity. Asian Pacific J. of Allergy and Immunology 10: 159-164.
- Tattam, B., Starmer, G., and Mascord, D. A comparative study of blood & saliva with analysis by GC/MS/MS for the analysis of ephedrine in potential road users. [online]. (n.d.). Available from: <http://www.latrobe.edu.au/anzsms/Conferences/ANZSMS17/ANZSMS17Abs/WeP/WeP-12.pdf> [2005. Dec 20].
- The National Institutes of Health Chemical Genomics Center. 2005. Immunoassay methods[Online]. Eli Lilly. Available from: <http://www.NIH Chemical Genomics Center -- Assay Guidance -- Assay Guidance Manual - Version 4 1 -- Immunoassay Methods.htm>[2005].
- Thongnopnua, P., Hongratanaworakit, T., and Lipipun, V. The effect of side chain heterologous combination of enzyme-labeled theophylline and immunogenic on theophylline enzyme immunoassay sensitivity. The 12<sup>th</sup> Pharmaceutical Research Meeting of Faculty of Pharmaceutical Sciences. Chulalongkorn University, Dec. 6, 1995. Thailand.
- Thongnopnua, P., Pitchayawasin, S., and Vangveravong, S. The sensitivity and specificity enhancement of solid-phase theophylline enzyme immunoassay by heterologous combinations. AAPS Annual Meeting & Exposition. Nov 14-23, 1998. San Francisco, USA.

- Thongnopnua, P., and Matapatara, W. Heterologous combination for amphetamine and methamphetamine immunodetection. PAC/FAPA Meeting. Austraria. 2000.
- Thongnopnua, P., and Sirikatitham. Ephedrine analytical method development through immunoassay. E-proceeding of 20<sup>th</sup> FAPA Congress 2004. Thailand. Nov 30 – Dec 3, 2004.
- Tijssen, P. Practice and theory of enzyme immunoassay. New York: Elsevier-North Holland, 1985.
- Trouble shooting protein binding in nitrocellulose membrane. 1999. [online].  
Available from: <http://www.devicelink.com/ivdt/archive/99/03/009.html>[2004]
- Tuomola, M., Harpio, R., and Mikola, H. 2000. Production and characterization of monoclonal antibodies against a very small hapten, 3-methylidole.  
J. Immunol. Methods 240: 111-124.
- Turner, G.J., Colbert, D.L., and Chowdry, B.Z. 1991. A broad spectrum immunoassay using fluorescence polarization for the detection of amphetamines in urine.  
Ann. Clin. Biochem. 28: 588-594.
- Urine testing[Online]. Available from: <http://www.erowid.org>[2006].
- Van Weemen, B.K., and Schuurs, A.H.W.M. 1972. Immunoassay using hapten-enzyme conjugates. FEBS LETTERS 24(1): 77-81.
- Van Weemen, B.K., and Schuurs, A.H.W.M. 1975. The influence of heterologous combinations of antiserum and enzyme-labeled estrogen on the characteristic of estrogen enzyme-immunoassays. Immunochemistry 12: 667-670.
- Verstraete, A.G., and Heyden, F.V. 2005. Comparison of the sensitivity and specificity of six immunoassays for the detection of amphetamines in urine. J. Anal. Toxicol. 29(5): 359-364.
- Walker, J.M., eds. The protein protocols handbook. New Jersey: Human Press, 1996.
- Wang, B.L., Scopsi, L., and Larsson, L.I. 1985. Simplified purification and testing of colloidal gold probes. Histochemistry 83: 109-115.
- Ward, C., McNally, A.J., Rusyniak, D., and Salamone, S.J. 1994. 125I radioimmunoassay for the dual detection of amphetamine and methamphetamine.  
J. Forensic. Sci. 39(6): 1486-1496.
- Wild, D., ed. The immunoassay handbook. Basingstock: The Macmillan Press, 1994.

- Wilson, M.B., and Nakane, P.K. Recent developments in the periodate method of conjugation horseradish peroxidase (HRPO) to antibodies. In W. Knapp; R. Holubar; and G. Wick (eds.), Immunofluorescence and related staining techniques. pp. 215-224. New York: Elsevier-North Holland Biochemical Press, 1978.
- Wortberg, M., Goodrow, M.H., Gee, S.J., and Hammock B.C. 1996. Immunoassays for simazine and triazine with low cross-reactivity for promazine. J. Agric. Chem. 44: 2210-2221.
- Wortberg, M., Kreissig, S.B., Jones, G., Rocke, D.M., and Hammock, B.D. 1995. An immunoarray for the simultaneous determination of multiple triazine herbicides. Anal. Chim. Acta. 304: 339-352.
- Wu, Y., Mitchell, J., Cook, C., and Main, L. 2002. Evaluation of progesterone-ovalbumin conjugates with different length linkers in enzyme-linked immunosorbent assay and plasmon resonance-based immunoassay. Steroid 67: 565-572.
- Yalow, R.S., and Berson, S.A. Immunoassay of plasma insulin. In D. Grick (ed.), Methods of biochemical analysis. vol. 12, pp. 69-96. New York: Interscience, 1962.
- Yamamoto, I., Matsuura, E., Horiba, M., Akima, K., Nomura, K., and Aizawa, T. 1985. Enzyme immunoassay for mabuterol, a selective b2-adrenergic stimulant in the trachea. J. Immunoassay. 6: 261-273.
- Zhang, J., Xie, J., Chen, X., and Hua, Z. 2003. Sensitive determination of ephedrine and pseudoephedrine by capillary electrophoresis with laser-induced fluorescence detection. Analyst 128(4): 369-372.

## VITA

Miss Waliluk Matapatara was born on August 15, 1964 in Phitsanuloke, Thailand. She has received her Bachelor of Science in Pharmacy in 1987 from the Faculty of Pharmaceutical Sciences, Chiangmai University, Chiangmai, Thailand. Her Master of Sciences in Pharmacy in 1999 from the Faculty of Pharmaceutical Sciences, Chulalongkorn University. Since her graduation, she has become a member of the Bureau of Drug and Narcotic, Department of Medical Sciences, Public Health Administration, Thailand.

### Poster Presentations

1. Matapatara, W., and Thongnopnua, P. "Effect of Heterology on the Detection of Amphetamine, Methamphetamine and Ephedrine by Enzyme-immunoassay" The 20<sup>th</sup> Annual Research Meeting in Pharmaceutical Sciences, December 1, 2003, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok.
2. Matapatara, W., Thongnopnua, P., and Lipipun, V. "Comparison of Specificity and Sensitivity of Antibodies from Cell Culture Supernatant to Their Polyclonal Antibodies for Amphetamine, Methamphetamine and Ephedrine Detection" The 5<sup>th</sup> National Symposium on Graduate Research, October 10-11, 2005, Kasetsart University, Bangkok.
3. Matapatara, W., Thongnopnua, P., and Lipipun, V. "The Influence of Hybridoma Cell Medium on Specificity and Sensitivity of Antibody for Amphetamine, Methamphetamine and Ephedrine Detection" The 2<sup>nd</sup> AASP (Asian Associate of Schools of Pharmacy) Symposium/2<sup>nd</sup> APEM Conference 2005, November 14-17, 2005, Montein Riverside Bangkok Hotel, Bangkok.

