

REFERENCES

- Abidi S.L. Chromatographic analysis of plant sterols in foods and vegetable oils J Chromatogr A. 935(2001): 173-201(29).
- Adair, C.R. Production and utilization of rice. Ed Houston, D.F., In Rice: Chemistry and Technology. pp. 1-13. American Association of Cereal Chemists, Inc., St. Paul, MN. 1972.
- Akihisa, T., Yasukawa, K., Kimura, Y., Takase, S., Yamanouchi, S. and Tamura, T. Triterpene alcohols from camellia and sasanqua oils and their anti-inflammatory effects. Chem Pharm Bull (Tokyo). 45(1997): 2016-23.
- Akihisa, T., Yasukawa, K., Yamaura, M., Ukiya, M., Kimura, Y., Shimizu, N. and Arai, K. Triterpene alcohol and sterol ferulates from rice bran and their anti-inflammatory effects. J. Agri. Food Chem. 48(2000): 2313-2319.
- Ansari, G.A., Walker, R.D., Smart, V.B. and Smith, L.L. Further investigations of mutagenic cholesterol preparations. Food Chem Toxicol. 20(1982): 35-41.
- Armstrong, M.J. and Carey, M.C. Thermodynamic and molecular determinants of sterol solubilities in bile salt micelles. J Lipid Res 28(1987): 1144-55.
- Awad, A. B., Chen, Y. C. and Fink, C. S. and Hennessey, T. β -sitosterol inhibits HT-29 human colon cancer cell growth and alters membrane lipids. Anticancer Res. 16(1996): 2797-2804.
- Awad, A. B., Down, A. C. and Fink, C. S. Inhibition of growth and stimulation of apoptosis by β sitosterol treatment of MDA-MB- 231 human breast cancer cells in culture. Int. J. Mol. Med. 5(2000b), 541-546.
- Awad, A. B., Downie, A., Fink, C. S. and Kim, U. Dietary phytosterol inhibits the growth and metastasis of MDA-MB-231 human breast cancer cells grown in SCID mice. Anticancer Res. 20(2000c): 821-824.
- Awad, A. B. and Fink, C. S. Phytosterols as anticancer dietary components: evidence and mechanism of action. J. Nutr. 130(2000): 2127-2130.
- Awad, A. B., Gan, Y. and Fink, C. S. Effect of β -sitosterol, a plant sterol, on growth, protein phosphatase 2A and phospholipids D in LNCaP cells. Nutr. Cancer 36 (2000a): 74-78.

- Banthorpe, D.V. Classification of terpenoids and general procedures for their characterization. Eds. Dey, P.M. and Harborne, J.B., In: Methods in Plant Biochemistry Vol. 7. Terpenoids. Eds. Charlwood, B.V. and Banthorpe, D.V., pp. 1-42. Academic Press Inc., San Diego, CA. 1991.
- Becker, M., Staab, D. and Von Bergmann, K. Treatment of severe familial hypercholesterolemia in childhood with sitosterol and sitostanol. J. Pediatr. 122(1993): 292-6.
- Blois, M.S. Antioxidant determination by the use of a stable free radical. Nature 181(1958): 1199-200.
- Booth, A. N., Emerson, O., Jones, F. T. and De Eds, F. Urinary metabolites of caffeic and chlorogenic acids. J. Biol. Chem. 229(1957): 51-59.
- Borgstrom, B. Quantitative aspects of the intestinal absorption and metabolism of cholesterol and beta-sitosterol in the rat. J. Lipid Res. 9(1968): 473-481.
- Bramley, P.M. Isoprenoid metabolism. Eds. Dey, P.M. and Harborne, J.B., In: Plant Biochemistry. pp. 471-426. Academic Press Inc., San Diego, CA. 1997.
- Brand-Williams, W., Cuvelier, M.E. and Berset, C. Use of a free radical method to evaluate antioxidant activity. Lebensm-Wiss U-Technol 28(1995): 25-30.
- Bucci, R., Magri, A. D., Magri, A. L. and Marini, F. Comparison of three spectrophotometric methods for the determination of γ -oryzanol in rice bran oil. Anal. Bioanal. Chem. 375(2003): 1254-1259.
- Carr, T.P., Cornelisona, R.M., Illston, B.J., Stuefer-Powell, C.L., and Gallaher, D.D. Plant sterols alter bile acid metabolism and reduce cholesterol absorption in hamsters fed a beef-based diet. Nutr. Res. 22(2002): 745-754.
- Chen, M.-H. and Bergman, C. J. A rapid procedure for analysing rice bran tocopherol, tocotrienol and γ -oryzanol contents. J. Food Comp. Anal. 18(2005): 319-331.
- Child, P. and Kuksis, A. Uptake of 7-dehydro derivatives of cholesterol, campesterol, and β -sitosterol by rat erythrocytes, jejunal villus cells, and brush border membranes. J. Lipid Res. 24(1983): 552-565.
- Cicero, A.F. and Gaddi, A. Rice bran oil and gamma-oryzanol in the treatment of hyperlipoproteinæmias and other conditions. Phytother. Res. 15(2001): 277-289.

- Clouse, S.D. Plant development: A role for sterols in embryogenesis. Curr. Bio. 10(2000): R601-R604.
- Das, P.K., Chaudhuri, A., Kaimal, T. N. B. and Bhalerao, U. T. Isolation of γ -oryzanol through calcium ion induces precipitation of anionic micellar aggregates. J. Agric. Food Chem. 46(1998): 3073-3080.
- De, B.K. and Bhattacharyya, D.K. Deacidification of high-acid rice bran oil by reesterification with monoglyceride. J Oil Technol Assoc India 32(2000): 10-11.
- Diack, M. and Saska, M. Separation of vitamin E and γ -oryzanol from rice bran oil by normal-phase chromatography. J. Am.Oil Chem. Soc. 71(1994): 1211-1217.
- Dreshner, E. E., Cohen, B. J. and Raicht, R. F. The kinetics of the protective effect of β -sitosterol against MNU-induced colonic neoplasia. J. Cancer Res. Clin. Oncol. 103(1982): 49-54.
- Einhellig, F. A. Mechanisms and modes of action of allelochemicals. Eds A. R. Putnam and C. S. Tang, In The Science of Allelopathy. pp. 171-188. John Wiley & Sons, New York. 1986.
- Endo, T., Misu, O. and Inaba, Y. Process for the isolation of oryzanols from rice bran oil. Yukagaku 18(1969): 63-67.
- Endo, T., Ueno, K. and Iwaba, Y. A new method for measuring oryzanol concentration. Yukagaku 17(1968): 344-348.
- Evershed, R. P., Spooner, N., Prescott, M. C. and Goad, L. J. Isolation and characterization of intact steryl ferulates from seeds. J. Chromatogr. 440(1988): 23-35.
- Fang, N., Yu, S. and Badger, T. M. Characterization of triterpene alcohol and sterol ferulates in rice bran using LC-MS/MS. J. Agri. Food Chem. 51(2003):3260-3267.
- Fieser, L. F. and Fieser, M. Sterols. (Chapt. 11) In Steroids pp. 341-363. Van Nostrand Reinhold: New York. 1959.
- Folch, J., Lees, M. and Sloane Stanley, G.H. A simple method for the isolation and purification of total lipides from animal tissues. J Biol Chem. 226 (1957): 497-509.
- Fry, A.C., Bonner, E., Lewis, D.L., Johnson, R.L., Stone, M.H. and Kraemer, W.J. The effects of gamma-oryzanol supplementation during resistance exercise training. Int J Sport Nutr. 7(1997): 318-29.

- Fujiwara, S., Noumi, K., Sugimoto, I. and Awata, N. Mass fragmentographic determination of ferulic acid in plasma after oral administration of gamma-oryzanol. *Chem Pharm Bull (Tokyo)*. 30(1982): 973-9.
- Fujiwara, S., Sakurai, S., Noumi, K., Sugimoto, I. and Awata, N. Metabolism of gamma-oryzanol in rabbit (author's transl). *Yakugaku Zasshi*. 100(1980): 1011-8.
- Fujiwara, S., Sakurai, S., Sugimoto, I. and Awata, N. Absorption and metabolism of gamma-oryzanol in rats. *Chem Pharm Bull (Tokyo)*. 31(1983): 645-52.
- Gerhardt, A.L. and Gallo, N.B. Full-fat rice bran and oat bran similarly reduce hypercholesterolemia in humans. *J Nutr*. 128(1998): 865-9.
- Gong, Y. and Yao, H. Analysis of γ -oryzanol by LC/MS. *Huaxue Shijie* 42(2001): 123 - 125.
- Gonzalez, F.J. The molecular biology of cytochrome P450s. *Pharmacol Rev* 41(1989): 91-2.
- Graf, E. Antioxidant potential of ferulic acid. *Free Radic Biol Med*. 13(1992): 435-48.
- Gremaud, G., Dalan, E.C.P., Baumgartner, M., Ballabeni, P., Decarli, B., Leser, M., Berger, A., and Fay, L.B. Effects of nonesterified stanols in a liquid emulsion on cholesterol absorption and synthesis in hypercholesterolemic men. *Eur J Nutr* 41(2002): 54-60.
- Grundy, S.M. Absorption and metabolism of dietary cholesterol. *Annu Rev Nutr*. 3(1983): 71-96.
- Grundy, S.M., Ahrens, E.H. and Davignon, J. The interaction of cholesterol absorption and cholesterol synthesis in man. *J Lipid Res*. 10(1969): 304-15.
- Grunewald, K.K. and Bailey, R.S. Commercially marketed supplements for bodybuilding athletes. *Sports Med*. 15(1993): 90-103.
- Guenoerich, F. P. Cytochrome P450: advances and prospects. *FASEB Journal* 6(1992): 667-668.
- Gylling, H. and Miettinen, T.A. Serum cholesterol in gastrointestinal diseases. [Article in Finnish] *Duodecim*. 111(1995): 2123-32.
- Hakala, P., Lampi, A. M., Ollilainen, V., Werner, U., Murkovic, M., Wahala, K., Karkola, S. and Phronen, V. Steryl phenolic acid esters in cereals and their milling fractions. *J Agric. Food Chem*. 50(2002), 5300-5307.

- Hargrove, K.L. Processing and Utilization of Rice Bran in the United States. Eds W.E. Marshall and J.I. Wadsworth, In Rice Science and Technology. pp.381-404. Marcel Dekker, New York, 1994.
- Hartmann, M. Plant sterols and the membrane environment. Trends Plant Sci. 3(1998):170-175.
- Heinemann, T., Kullak-Ublick, G.-K., Pietruck, B. and von Bergmann, K. Mechanisms of action of plant sterols on inhibition of cholesterol absorption. Eur J Clin Pharmacol 40(1991): S50-63.
- Heinemann, T., Axtmann, G. and Von Bergmann, K. Comparison of intestinal absorption of cholesterol with different plant sterols in man. Eur. J. Clin. Invest. 23(1993): 827-831.
- Hendriks, H. F., Weststrate, J. A., Vliet, T. V., and Meijer, G. W. Spreads enriched with three different levels of vegetable oil sterols and the degree of cholesterol lowering in normocholesterolaemic and mildly hypercholesterolaemic subjects. Eur J Clin Nutr 53(1999): 319-27.
- Heuman, D.M. Quantitative estimation of the hydrophilic-hydrophobic balance of mixed bile salt solutions. J Lipid Res. 30(1989): 719-30.
- Hiramitsu, T. and Armstrong, D. Preventive effect of antioxidants on lipid peroxidation in the retina. Ophthalmic Res. 23(1991): 196-203.
- Hirose, M., Ozaki, K., Takaba, K., Fukushima, S., Shirai, T. and Ito, N. Modifying effects of the naturally occurring antioxidants in γ -oryzanol, phytic acid, tannic acid and n-tritriacontane-16, 18-dione in a rat wide-spectrum organ carcinogenesis model. Carcinogenesis 12(1991): 1917-1921.
- Horst, A. and Andreas, K. Hair cosmetic cleaning composition with active compound combinations consisting of gamma oryzanol and calcium salts with a low metal ion concentration. EP1475073. 2004.
- Hu, W., Wells, J. H., Shin, T., and Godber, J. S. Comparison of isopropanol and hexane for extraction of vitamin E and oryzanols from stabilized rice bran. J. Am. Oil Chem. Soc. 73(1996): 1653-1656.
- Huang, D., Ou, B., Hampsch-Woodill, M., Flanagan, J.A. and Deemer, E.K. Development and validation of oxygen radical absorbance capacity assay for

- lipophilic antioxidants using randomly methylated β -cyclodextrin as the solubility enhancer. *J Agric Food Chem.* 50(2002): 1815–1821.
- Huke, R. E. and Huke, E. H. *Rice: Then and Now*. International Rice Research Institute, Manila, The Philippines. 1990.
- Ikeda, I. and Sugano, M. Some aspects of mechanism of inhibition of cholesterol absorption by β -sitosterol. *Biochim Biophys Acta* 732(1983): 651–8.
- Ikeda, I., Sugano, M., Scallen, T.J., Vahouny, G.V. and Gallo, L.L. Transfer of cholesterol and sitosterol from rat intestinal brush border membranes to phospholipid liposomes: effects of SCP. *Agri. Biol. Chem.* 54(1990): 2646-2654.
- Ikeda, I., Tanaka, K., Sugano, M., Vahouny, G.V. and Gallo, L.L. Discrimination between cholesterol and sitosterol for absorption in rats. *J Lipid Res* 29(1988b):1583–91.
- Ikeda, I., Tanaka, K., Sugano, M., Vahouny, G.V. and Gallo, L.L. Inhibition of cholesterol absorption in rats by plant sterols. *J Lipid Res* 29(1988a):1573–82.
- Imai, Y., Ogawa, T., Tsurumi, C., Kitagawa, M. and Tanaka, H. Food additive and use thereof. US patent number 5,514,398. 1996.
- Imaida, K., Hirose, M., Yamaguchi, S., Takahashi, S. and Ito, N. Effects of naturally occurring antioxidants on combined 1,2-dimethylhydrazine- and 1-methyl-1-nitrosourea-initiated carcinogenesis in F344 male rats. *Cancer Lett.* 55(1990): 53-59.
- Iqbal, S., Bhanger, M.I., Anwar, F. Antioxidant properties and components of some commercially available varieties of rice bran in Pakistan. *Food Chem.* 93(2005): 265-272.
- Iwatsuki, K., Akihisa, T., Tokuda, H., Ukiya, M., Higashihara, H., Mukainaka, T., Iizuka, M., Hayashi, Y., Kimura, Y. and Nishino, H. Sterol ferulates, sterols, and 5-alk(en)ylresorcinols from wheat, rye, and corn bran oils and their inhibitory effects on Epstein-Barr virus activation. *J Agric Food Chem.* 51(2003): 6683-8.
- Jariwalla, R.J. Rice-bran products: phytonutrients with potential applications in preventive and clinical medicine. *Drugs Exp Clin Res.* 27(2001): 17-26.
- Johnson, L.A. and Lusas, E W. Comparison of alternative solvents for oils extraction. *J Am Oil Chem Soc* 60(1983): 229–42.

- Juliano, B.O. and Bechtel, D.B. The rice grain and its gross composition. Ed. Juliano, B.O., In: Rice: Chemistry and Technology. pp. 17-57. American Association of Cereal Chemists, Inc., St. Paul, MN. 1985.
- Kahlon, T. S., Chow, F. I., Chiu, M. M., Hudson, C. A. and Sayre, R. N. Cholesterol-lowering by rice bran and rice bran oil unsaponifiable matter in hamsters. Cereal Chem. 73(1996): 69-74.
- Kaimal, T.N.B. γ -Oryzanol from rice bran oil. Oil Technologists' Association of India (Southern Zone) Newsletter. 3(1999): 6-9.
- Kamimura, M., Takahashi, S. and Sato, S. Influence of γ -oryzanol on the skin capillary circulation. Bitamin. 30(1964): 341-345.
- Kamm, W., Dionisi, F., Fay, L.-B., Hischenhuber, C., Schmarr, H.-G. and Engel, K.-H. Analysis of steryl esters in cocoa butter by on-line liquid chromatography-gas chromatography. J. Chromatogr. A. 918(2001): 341-349.
- Kaneko, R. and Tsuchiya, T. New compounds in rice bran and germ oils. J. Chem. Soc. Jpn. 57(1954): 526-529.
- Kao, C., Luh, B.S. Rice oil. Ed. Luh, B.S., In: Rice: Utilization. Vol. 2, pp. 295-362. Van Nostrand Reinhold, New York, NY. 1980.
- Kerckhoffs, D.A., Brouns, F., Hornstra, G. and Mensink, R.P. Effects on the human serum lipoprotein profile of beta-glucan, soy protein and isoflavones, plant sterols and stanols, garlic and tocotrienols. J Nutr. 132(2002): 2494-2505.
- Kikuzaki, H., Hisamoto, M., Hirose, K., Akiyama, K. and Taniguchi, H. Antioxidant properties of ferulic acid and its related compounds. J Agric Food Chem. 50(2002): 2161-8.
- Kim, J.-S., Godber, J.S., King, J.M. and Prinyawiwatkul, W. Inhibition of cholesterol autoxidation by the nonsaponifiable fraction in rice bran in an aqueous model system. J. Am.Oil Chem. Soc. 78(2001): 685-689.
- Klick, S. and Herrmann, K. Glucosides and glucose esters of hydroxybenzoic acids in plants. Phytochem. 27(1988): 2177-2180.
- Kochhar, S.P. Influence of processing on sterols of edible vegetable oils. Prog Lipid Res. 22(1983): 161-188.

- Krishna, A.G.G., Sakina K., Shiela, P.M., Sarmandal, C.V., Indira, T.N. and Mishra, A. Effect of refining of crude oil on retention of oryzanol in refined rice bran oil. *J. Am. Oil Chem. Soc.* 78(2001): 127–131.
- Kudchodkar, B.J., Horlick, L. and Sodhi, H.S. Effects of plant sterol on cholesterol metabolism in man. *Atherosclerosis*. 23(1976): 239–248.
- Kudo, S. and Akiyama, H. High-performance liquid chromatographic procedure for the determination of a new antithrombotic and vasodilating agent in human plasma. *Otsuka Pharmaceut. In-House Rep.*, 27(1982): 419-422.
- Kumar, N. and Singhal, O. P. Cholesterol oxides and atherosclerosis: a review. *J. Sci. Food Agric.* 55(1991): 497-510.
- Lang, J.K., Schillaci, M. and Irvin, B. Vitamin E. Eds. De Leenheer, A.P., Lambert, W.E., Nelis, H.J. In *Modern Chromatographic Analysis of Vitamins*. pp. 153–195. Marcel Dekker, New York. 1992.
- Law, M. Plant sterol and stanol margarines and health. *Br Med J.* 320(2000):861–864.
- Lees, A.M., Mok, H.Y.I., Lees, R.S., McCluskey, S. and Grundy, S.M. Plant sterols as cholesterol-lowering agents: clinical trials in patients with hypercholesterolemia and studies of sterol balance. *Atherosclerosis* 28(1977): 325–338.
- Lichtenstein, A.H., Ausman, L.M., Carrasco, W., Gualtieri, L.J., Jenner, J.L., Ordovas, J.M., Nicolosi, R.J., Goldin, B.R. and Schaefer, E.J. Rice bran oil consumption and plasma lipid levels in moderately hypercholesterolemic humans. *Arterioscler Thromb.* 14(1994): 549-556.
- Ling, W. H. and Jones, P.J.H. Dietary phytosterols: a review of metabolism, benefits and side effects. *Life Sci.* 57(1995): 195–206.
- Lu, S. and Luh, B.S. Properties of the rice caryopsis. Ed. Luh, B.S., In *Rice: Production*. Vol. 1, pp. 389-419. Van Nostrand Reinhold, New York, NY. 1991.
- Lyon, M. A. and Brown, A. J. 7-Ketocholesterol. *Int. J. Biochem. Cell Biol.* 31(1999): 369–375.
- Macheix, J.-J., Fleuriet, A. and Billot, J. The main phenolics of fruit. In *Fruit Phenolics*. pp. 1-98. CRC Press, Boca Raton, USA. 1990.
- Mattson, F.H., Grundy, S. M., and Crouse, J.R. Optimizing the effect of plant sterols on cholesterol absorption in man. *Am J Clin Nutr* 35(1982):697–700.

- McCaskill, D.R. and Orthoefer, F.T. Storage stability of extrusion stabilized and parboiled rice bran. (Chapt. 3) Eds. Marshall, W. E., Wadsworth, J. I. In Rice Science and Technology pp. 37-48. Marcel Dekker: New York, 1994.
- McCaskill, D.R. and Zhang, F. Use of rice bran oil in foods. Food Technol 53(1999): 50-53.
- Miettinen, T. A. and Gylling, H. Regulation of cholesterol metabolism by dietary plant sterols. Curr. Opin. Lipidol. 10(1999): 9-14.
- Miettinen, T.A., Puska, P., Gylling, H., Vanhanen, H., and Vartiainen, E. Reduction of Serum Cholesterol with Sitostanol-Ester Margarine in a Mildly Hypercholesterolemic Population. Engl. J. Med. 333(1995): 1308 - 1312.
- Miettinen, T.A., Vuoristo, M., Nissinen, M., Jarvinen, H.J., and Gylling, H. Serum, biliary, and fecal cholesterol and plant sterols in cholectomized patients before and during consumption of stanol ester margarine. Am J Clin Nutr 71(2001): 1095 - 1102.
- Miller, A., Frenzel, T., Schmarr, H. and Engel, K. Coupled liquid chromatography-gas chromatography for the rapid analysis of γ -oryzanol in rice lipids. J. Chromatogr. A. 985(2003): 403-410.
- Miller, A., Majauskaite, L. and Engel, K.-H. Enzyme-catalyzed hydrolysis of γ -oryzanol. Eur Food Res Technol 218(2004): 349-354.
- Moldenhauer, K.A., Champagne, E.T., McCaskill, D.R. and Guraya, H. Functional products from rice (Chapt. 3). Ed Mazza, G., In Functional Foods: Biochemical and Processing Aspects pp.71-89. Technomic Pub.: Lancaster, PA, 1998.
- Moreau, R.A. and Hicks, K.B. The in vitro hydrolysis of phytosterol conjugates in food matrices by mammalian digestive enzymes. Lipids. 39(2004):,769-776.
- Moreau, R. A., Powell, M. I., and Hicks, K. B. Extraction and quantitative analysis of oil from commercial corn fiber. J. Agric. Food Chem 44(1996): 2149-2154.
- Moreau, R.A., Singh, V., Eckhoff, S.R., Powell, M.J., Hicks, K.B. and Norton, R.A. Comparison of yield composition of oil extracted from corn fiber and corn bran. Cereal Chem. 76(1999): 449-451.
- Morel, D. W. and Lin, C. Y. Cellular biochemistry of oxysterols derived from the diet or oxidation in vivo. J. Nutr. Biochem. 7(1996): 495-506.

- Nicolosi, R. J., Rogers, E. J., Ausman, L. M. and Orthoefer, F. T. Rice bran oil and its health benefits. (Chapt. 18) Eds. Marshall, W. E., Wadsworth, J. I. In Rice Science and Technology pp. 421-437. Marcel Dekker: New York, 1994.
- Noboru, K. and Yusho, T. Oryzanol containing cosmetics. JP patent number 70:32078. 1970.
- Normén, L., Johnsson, M., Andersson, H., van Gameren, Y. and Dutta, P. Plant sterols in vegetables and fruits commonly consumed in Sweden. Eur J Nutr 38(1999): 84-89.
- Norton, R. A. Isolation and identification of steryl cinnamic acid derivatives from corn bran. Cereal Chem. 71(1994): 111-117.
- Norton, R. A. Quantitation of steryl ferulate and *p*-coumarate esters from corn and rice bran. Lipids. 30(1995): 269-274.
- Nyström, L., Makinen, M., Lampi, A.M., Piironen, V. Antioxidant activity of steryl ferulate extracts from rye and wheat bran. J Agric Food Chem. 53(2005): 2503-2510.
- Odomi, M., Ishigami, M., Shimizu, T. Probucol prevents the progression of atherosclerosis in Watanabe heritable hyperlipidemic rabbit, an animal model for familial hypercholesterolemia. Otsuka Pharmaceut. In-House Rep. 64(1983): 592-598.
- Ohta, G. and Shimizu, M. Studies on the constituents of rice bran oil: Structure of oryzanol-A. Pharm Bull. 5(1957): 40-44.
- Okada, T. and Yamaguchi, N. Antioxidative effect and pharmacology of oryzanol. Yakugaku (J. Jap. Oil Chem. Soc.). 32(1983): 305-311.
- Orthoefer, F.T. Rice bran oil: Healthy lipid source. Food Technoloay 50(1996): 62-64.
- Ostlund, R.E., Spilburg, C.A., and Stenson, W.F. Sitostanol administered in lecithin micelles potently reduces cholesterol absorption in humans. Am J Clin Nutr 70(1999): 826-831.
- Paniangvait, P., King, A. J., Jones, A. D., and German, B. G. Cholesterol oxides in foods of animal origin. J. Food Sci. 60(1995): 1159-1174.
- Parrado, J., Miramontes, E., Jover, M., Marquez, J.C., Angeles Mejias. M., Collantes De Teran, L., Absi, E. and Bautista, J. Prevention of brain protein and lipid oxidation

- elicited by a water-soluble oryzanol enzymatic extract derived from rice bran. *Eur J Nutr.* 42(2003): 307-314.
- Piironen, V., Syvaloja, E., Varo, P., Salminen, K. and Koivistoinen, P. Tocopherols and tocotrienols in cereal products from Finland. *Cereal Chem.* 63(1986): 78-81.
- Piironen, V., Toivo, J. and Lampi, A.-M. Natural Sources of Dietary Plant Sterols. *J Food Compo. Anal.* 13(2000): 619-624(6).
- Plat, J., Kerckhoffs, D. A., and Mensink, R. P. Therapeutic potential of plant sterols and stanols. *Curr. Opin. Lipidol.* 11(2000): 571-576.
- Plat, J. and Mensink, R. P. Effects of plant stanol esters on LDL receptor protein expression and on LDL receptor and HMG-CoA reductase mRNA expression on mononuclear blood cells of healthy men and women. *FASEB J.* 16(2002): 258-260.
- Plat, J. and Mensink, R. P. Effects of plant sterols and stanols on lipid metabolism and cardiovascular risk. *Nutr. Metab. Cardiovasc. Dis.* 11(2001): 31-40.
- Pouteau, E.B., Monnard, I.E., Piguet-Welsch, C., Groux, M.J.A., Sagalowicz, L. and Berger, A. Non-esterified plant sterols solubilized in low fat milks inhibit cholesterol absorption: A stable isotope double-blind crossover study. *Eur J Nutr* 42(2003): 154-164.
- Qiu, Y. and Tang, B. Determination of ferulates in suppository by HPLC. *Huaxi Yaoxue Zazhi.* 13(1998): 36-38.
- Qureshi, A.A., Mo, H., Packer, L. and Peterson, D. M. Isolation and identification of novel tocotrienols from rice bran with hypocholesterolemic, antioxidant, and antitumor properties. *J. Agric. Food Chem.* 48(2000): 3130-3140.
- Qureshi, A.A., Sami, S.A. and Khan, F.A. Effects of stabilized rice bran, its soluble and fiber fractions on blood glucose levels and serum lipid parameters in humans with diabetes mellitus Types I and II. *J Nutr Biochem.* 13(2002): 175-187.
- Raghuram, T.C., Rao, U.B. and Rukmini, C. Studies on hypolipidemic effects of dietary rice bran oil in human subjects. *Nutr. Rep. Intl.* 39(1989): 889-895.
- Raicht, R. F., Cohen, L. I., Fazzini, E. P., Sarwal, A. N. and Takahashi, M. Protective effect of plant sterols against chemically induced colon tumors in rats. *Cancer Res.* 40(1980): 403-405.

- Ramarathnam, N., Osawa, T., Namiki, M. and Kawakishi, S. Chemical studies on novel rice hull antioxidants. 1. Isolation, fractionation, and partial characterization. J. Agric. Food Chem. 36(1988): 732-737.
- Randall, J.M., Sayre, R.N., Schultz, W.G., Fong, R.Y., Mossman, A.P., Trihethorn, R.E., and Saunders, R.M. Rice bran stabilization by extrusion cooking for extraction of edible oil. J Food Sci. 50(1985): 361-368.
- Rao, A.V., and Janezic, S.A. The role of dietary phytosterols in colon carcinogenesis. Nutr Cancer. 18(1992): 43-52.
- Rao, K.V.S.A., Rao, B.V.S.K. and Kaimal, T.N.B. Process for the isolation of oryzanol from rice bran oil soap stock. US patent number 6,410,762B1. 2002.
- Rendell, N., Misso, N.L.A. and Goad, L.J. Biosynthesis of 24-methylcholest-5-en-3 β -ol and 24-ethylcholest-5-en-3 β -ol in *Zea mays*. Lipids. 21(1986): 63-68.
- Robbins R.J. Phenolic acids in food: an overview of analytical methodology. J Agric Food Chem. 51(2003), 2866-87.
- Rogers, E. J., Rice S. M., Nicolosi, R. J., Carpenter, D. R., McClelland, C. A. and Romanczyk, L. J. Identification and quantitation of γ -oryzanol components and simultaneous assessment of tocols in rice bran oil. J. Am. Oil Chem. Soc. 70(1993): 301-307.
- Rohrer, C. C., and Siebenmorgen, T. J. Nutraceutical concentrations within the bran of various rice kernel thickness fractions. Biosystems Eng. 88(2004): 453-460.
- Rong, N., Ausman, L.M. and Nicolosi, R.J. Oryzanol decreases cholesterol absorption and aortic fatty steaks in hamsters. Lipids 32(1997): 303-309.
- Rubinstein, I., Goad, L.J., Clague, A.D.H. and Mulheim, L.J. The 220 MHz NMR spectra of phytosterols. Phytochem. 15(1976): 195-200.
- Rukmini, C. Chemical, nutritional and toxicological studies of rice bran oil. Food Chem. 30(1988): 257-268.
- Rukmini, C. and Raghuram, T.C. Nutritional and biochemical aspects of the hypolipidemic action of rice bran oil: a review. J Am Coll Nutr. 10(1991): 593-601.
- Salen, G., Kwiterovich Jr, P.O., Shefer, S., Tint, G.S., Horak, I., Shore, V., Dayal, B. and Horak, E. Increased plasma cholestanol and 5 alpha-saturated plant sterol

- derivatives in subjects with sitosterolemia and xanthomatosis. *J. Lipid Res.* 26(1985): 203-209.
- Sasaki, J., Takada, Y., Handa, K., Kusuda, M., Tanabe, Y., Matsunaga, A. and Arakawa, K. Effects of Gamma-Oryzanol on Serum Lipids and Apolipoproteins in Dyslipidemic Schizophrenics Receiving Major Tranquilizers. *Clin. Ther.* 12(1990): 263-268.
- Sayre, B. and Saunders, R. Rice bran and rice bran oil. *Lipid Technol.* 2(1990): 72-76.
- Schaller, H., Bouvier-Navé, P. and Benveniste, P. Overexpression of an *Arabidopsis* cDNA encoding a sterol-C24¹-methyltransferase in tobacco modifies the ratio of 24-methyl cholesterol to sitosterol and is associated with growth reduction. *Plant Physiol.* 118(1998): 461-469.
- Seetharamaiah, G.S. and Chandrasekhara, N. Hypocholesterolemic activity of oryzanol in rats. *Nutr. Rep. Intl.* 38(1988): 927-935.
- Seetharamaiah, G.S. and Chandrasekhara, N. Studies on hypocholesterolemic activity of rice bran oil. *Atherosclerosis*, 78(1989): 219-223.
- Seetharamaiah, G.S., Krishnakantha, T.P. and Chandrasekhara, N. Influence of oryzanol on platelet aggregation in rats. *J. Nutr. Sci. Vit.* 36(1990): 291-297.
- Seetharamaiah, G.S. and Prabhakar, J.V. Oryzanol content of Indian rice bran oil and its extraction from soap stock. *J. Food Sci. Technol.* 23(1986): 270-273.
- Seitz, L. M. Stanol and sterol esters of ferulic and p-coumaric acids in wheat, corn, rye, and triticale. *J. Agric. Food Chem.* 37(1989): 662-667.
- Shen, Z., Palmer, M. V., Ting, S. S. T. and Fairclough, R. J. Pilot scale extraction and fractionation of rice bran oil using supercritical carbon dioxide. *J. Agric. Food Chem.* 45(1997): 4540-4544.
- Shin, T.-S., J. S. Godber, D. E. Martin, and J. H. Wells. Hydrolytic stability and changes in vitamin E vitamers and oryzanol of extruded rice bran during storage. *J. Food Sci.* 62(1997): 704-708, 728.
- Shugo, M. Anti-dandruff and anti-itching shampoo. Japanese patent number 79:36306. 1979.

- Sierksma, A., Weststrate, J. A. and Meijer, G. W. Spreads enriched with plant sterols, either esterified 4,4-dimethylsterols or free 4-desmethylsterols, and plasma total- and LDL-cholesterol concentrations. Br. J. Nutr. 82(1999): 273–282.
- Slota, T., Kozlov, N.A., and Ammon, H.V. Comparison of cholesterol and β -sitosterol: effects on jejunal fluid secretion induced by oleate, and absorption from mixed micellar solutions. Gut 24(1983): 653-658.
- Strack, D. Phenolic metabolism. In Eds Dey, P.M. and Harborne, J.B. Plant biochemistry. (Vol. 1) pp. 387-416. Academic Press, New York. 1997.
- Suga, K. Improvements in and relating to oryzanol solutions and compositions. GB patent number 1,134,878. 1968.
- Sugano, M., Morioka, H. and Ikeda, I. A comparison of hypocholesterolemic activity of β -sitosterol, and β -sitostanol in rats. J Nutr. 107(1977): 2011–2019.
- Sugano, M. and Tsuji, E. Rice bran oil and cholesterol metabolism. J Nutr. 127(1997): 521S-524S.
- Sunitha, T., Manorama, R. and Rukmini, C. Lipid profile of rats fed blends of rice bran oil in combination with sunflower and safflower oil. Plants Foods Human Nutr. 51(1997): 219-230.
- Suzuki, A., Kagawa, D., Fujii, A., Ochiai, R., Tokimitsu, I. and Saito, I. Short- and long-term effects of ferulic acid on blood pressure in spontaneously hypertensive rats. Am J Hypertens. 15(2002): 351-357.
- Suzuki, S. and Osima. Influence of blending of edible fats and oils on human serum cholesterol level (Part 1 and 2). Jap. J. Nutr. 28(1970): 3-6 and 194-198.
- Swell, L., Field, H. and Treadwell, C.R. Sterol specificity of pancreatic cholesterol esterase. Proc Soc Exp Biol Med 87(1954): 216–218.
- Takemoto, T., Miyoshi, H. and Nagashima, H. Clinical trial of Hi-Z fine granules (gamma-oryzanol) on gastrointestinal symptoms at 375 hospitals (Japan). Shinyaku To Rinsho [in Japanese] 11(1977): 26-32.
- Tamagawa, M., Otaki, Y., Takahashi, T., Otaka, T., Kimura, S. and Miwa, T. Carcinogenicity study of γ -oryzanol in B6C3F₁ mice. Food Chem. Toxicol. 30(1992a): 41-48.

- Tamagawa, M., Shimizo, Y., Takahashi, T., Otaka, T., Kimura, S., Kadowaki, H., Uda, F. and Miwa, T. Carcinogenicity study of γ -oryzanol in F344 rats. Food Chem. Toxicol. 30(1992a): 41-48.
- Tammi, A., Ro"nnemaa, T., Gylling, H., Rask Nissila", L., Viikari, J., Tuominen, J., Pulkki, K. and Simell, O. Plant stanol ester margarine lowers serum total and low-density lipoprotein cholesterol concentrations of healthy children: the STRIP project. Special Turku Coronary Risk Factors Intervention Project. J. Pediatr. 136(2000): 503-510.
- Taylor, J. B., Richar, T. M., Wilhelm, C. L., Chrysam, M. M., Otterburn, M. and Leveille, G. A. Rice bran oil antioxidant. US patent number 5,552,167. 1996.
- Tsuchiya, T., Kaneko, R. and Okubo, O. Solvent extraction process of oryzanol. Tokyo Kogyo Shikensho Hokoku. 52(1957): 1-3.
- Tsushima, G., Shibahara, T., Awogi, T., Kaneko, E., Sotou, S., Yamamoto, K. and Shirakawa, H. DNA-damaging, mutagenic, clastogenic and cell-cell communication inhibitory properties of γ -oryzanol. J. Toxicol. Sci. 16(1991): 191-202.
- Umehara, K., Shimokawa, Y. and Miyamoto, G. Effect of gamma-oryzanol on cytochrome P450 activities in human liver microsomes. Biol Pharm Bull. 27(2004): 1151-1153.
- Vissers, M. N., Zock, P. L. Meijer, G. W. and Katan, M. B. Effect of plant sterols from rice bran oil and triterpene alcohols from sheanut oil on serum lipoprotein concentrations in humans. Am. J. Clin. Nutr. 72(2000): 1510-1505.
- Vuorio, A. F., Gylling, H., Turtola, H., Kontula, K., Ketonen, P. and Miettinen, T. A. Stanol ester margarine alone and with simvastatin lowers serum cholesterol in families with familial hypercholesterolemia caused by the FH-North Karelia mutation. Arterioscler Thromb Vasc Biol. 20(2000): 500-506.
- Wang, T., Hicks, K.B. and Moreau, R. Antioxidant activity of phytosterols, oryzanol and other phytosterol conjugates. J. Am. Oil Chem. Soc. 79(2002): 1201-1204.
- Watanabe, K., Nakamura, R. and Hosono, A. Mutagenic activity of heat-induced cholesterol-degradation products. J. Food Sci. 38(1988): 119-123.

- Weststrate, J. A. and Meijer, G. W. Plant sterol-enriched margarines and reduction of plasma total- and LDL-cholesterol concentrations in normocholesterolaemic and mildly hypercholesterolaemic subject. *Eur. J. Clin. Nutr.* 52(1998): 334-343.
- Wheeler, K.B. and Garleb, K.A. Gamma oryzanol-plant sterol supplementation: metabolic, endocrine, and physiologic effects. *Int J Sport Nutr.* 1(1991): 170-177.
- Wilson, A. M., Sisk, R. M. and O'Brien, N. M. Modulation of cholestan-3 β ,5 $\alpha,6\beta$ -triol toxicity by butylated hydroxytoluene, α -tocopherol and β -carotene in newborn rat kidney cells in vitro. *Br. J. Nutr.* 78(1997): 479-492.
- Winter, M. and Herrmann, K. Esters and glucosides of hydroxycinnamic acids in vegetables. *J Agric Food Chem.* 34(1986): 616-620.
- Woods, J. A. and O'Brien, N. M. Investigation of the potential genotoxicity of cholesterol oxidation products in two mammalian fibroblast cell lines. *Nutr. Cancer* 31(1998): 192-198.
- Wu, H., Haig, T., Pratley, J., Lemerle, D., and An, M. Distribution and exudation of allelochemicals in wheat (*Triticum aestivum*). *J. Chem. Ecol.* 26(2000): 2141-2154.
- Wu, H., Haig, T., Pratley, J., Lemerle, D., and An, M. Simultaneous determination of phenolic acids and 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one in wheat (*Triticum aestivum*) by gas chromatography-tandem mass spectrometry. *J. Chromatogr. A* 864(1999): 315-321.
- Xu, Z. and Godber, J.S. Antioxidant activities of major components of γ -oryzanol from rice bran using a linoleic acid model. *J. Am. Oil Chem. Soc.* 78(2001): 645-649.
- Xu, Z. and Godber, J.S. Purification and identification of components of γ -oryzanol in rice bran oil. *J. Agri. Food Chem.* 47(1999): 2724-2728.
- Xu, Z., Hua, N. and Godber, J.S. Antioxidant activity of tocopherols, tocotrienols, and gamma-oryzanol components from rice bran against cholesterol oxidation accelerated by 2,2'-azobis (2-methylpropionamidine) dihydrochloride. *J. Agri. Food Chem.* 49(2001): 2077-2081.
- Yagi, K. and Ohishi, N. Action of ferulic acid and its derivatives as antioxidants. *J. Nutr. Sci. Vitaminol. (Tokyo)*. 25(1979): 127-130.

Yasukawa, K. and Akihisa, T. Anti-inflammatory and antitumor-promoting activities of sterols, triterpene alcohols and their derivatives. Recent Res. Devel. Oil Chem. 1(1997): 115-125.

Yasukawa, K., Takido, M., Matsumoto, T., Takeuchi, M. and Nakagawa, S. Sterol and triterpene derivatives from plants inhibit the effects of a tumor promoter, and sitosterol and betulinic acid inhibit tumor formation in mouse skin two-stage carcinogenesis. Oncology. 48(1991): 72-76.

Yasukawa, K., Akihisa, T., Kimura, Y., Tamura, T. and Takido, M. Inhibitory effect of cycloartenol ferulate, a component of rice bran, on tumor promotion in two-stage carcinogenesis in mouse skin. Biol. Pharm. Bull. 21(1998): 1072-1076.

Yoshino, G., Kazumi, T., Amano, M., Tateiwa, M., Yamasaki, T., Takashima, S., Iwali, M., Hatanaka, H. and Baba, S. Effects of gamma-oryzanol on hyperlipidemic subjects. Curr. Ther. Res. 45(1989): 543-552.

APPENDICES

Appendix A




Figure 1 ¹H-NMR spectrum of standard oryzanol




Figure 2 IR spectrum of oryzanol mixture PF1




Figure 3 ¹H-NMR spectrum of oryzanol mixture PF1




Figure 4 MS spectrum of oryzanol mixture PF1




Figure 5 MS/MS spectrum of oryzanol mixture PF1 at m/z 589




Figure 6 MS/MS spectrum of oryzanol mixture PF1 at m/z 615




Figure 7 MS/MS spectrum of oryzanol mixture PF1 at m/z 617




Figure 8 MS/MS spectrum of oryzanol mixture PF1 at m/z 633




Figure 9 IR spectrum of oryzanol mixture PF2




Figure 10 ¹H-NMR spectrum of oryzanol mixture PF2




Figure 11 MS spectrum of oryzanol mixture PF2




Figure 12 MS/MS spectrum of oryzanol mixture PF2 at m/z 561




Figure 13 MS/MS spectrum of oryzanol mixture PF2 at m/z 573




Figure 14 MS/MS spectrum of oryzanol mixture PF2 at m/z 575




Figure 15 MS/MS spectrum of oryzanol mixture PF2 at m/z 577




Figure 16 MS/MS spectrum of oryzanol mixture PF2 at m/z 589




Figure 17 MS/MS spectrum of oryzanol mixture PF2 at m/z 591




Figure 18 MS/MS spectrum of oryzanol mixture PF2 at m/z 601




Figure 19 MS/MS spectrum of oryzanol mixture PF2 at m/z 615




Figure 20 IR spectrum of oryzanol mixture WF1




Figure 21 ¹H-NMR spectrum of oryzanol mixture WF1




Figure 22 MS spectrum of oryzanol mixture WF1




Figure 23 MS/MS spectrum of oryzanol mixture WF1 at m/z 573




Figure 24 MS/MS spectrum of oryzanol mixture WF1 at m/z 575




Figure 25 MS/MS spectrum of oryzanol mixture WF1 at m/z 577




Figure 26 MS/MS spectrum of oryzanol mixture WF1 at m/z 587




Figure 27 MS/MS spectrum of oryzanol mixture WF1 at m/z 589




Figure 28 MS/MS spectrum of oryzanol mixture WF1 at m/z 601




Figure 29 MS/MS spectrum of oryzanol mixture WF1 at m/z 615




Figure 30 IR spectrum of oryzanol mixture SF1




Figure 31 ¹H-NMR spectrum of oryzanol mixture SF1





Figure 32 MS spectrum of oryzanol mixture SF1

Figure 33 MS/MS spectrum of oryzanol mixture SF1 at m/z 575




Figure 34 MS/MS spectrum of oryzanol mixture SF1 at m/z 589




Figure 35 MS/MS spectrum of oryzanol mixture SF1 at m/z 591




Figure 36 MS/MS spectrum of oryzanol mixture SF1 at m/z 601




Figure 37 MS/MS spectrum of oryzanol mixture SF1 at m/z 615




Figure 38 IR spectrum of oryzanol mixture CF1




Figure 39 ¹H-NMR spectrum of oryzanol mixture CF1




Figure 40 MS spectrum of oryzanol mixture CF1




Figure 41 MS/MS spectrum of oryzanol mixture CF1 at m/z 601




Figure 42 MS/MS spectrum of oryzanol mixture CF1 at m/z 615




Figure 43 IR spectrum of standard oryzanol




Figure 44 ¹H-NMR spectrum of standard oryzanol

Appendix B




Figure 1 Antioxidant activity of Pathumthani 1 extracts




Figure 2 Antioxidant activity of Kao Dok Mali 105 extracts




Figure 3 Antioxidant activity of Sunpatong 1 extracts




Figure 4 Antioxidant activity of Go Ko Chai Nat 1 extracts




Figure 5 Antioxidant activity of standard oryzanol




Figure 6 Antioxidant activity of α -tocopherol

BIOGRAPHY

Miss Lek Vejjanukroh was born on August 25, 1980, in Bangkok, Thailand. From Chulalongkorn University Demonstration School (Satit Chula), she has possessed the Certificate of High School since March 1998. She attended at the Department of Biochemistry, Faculty of Science, Chulalongkorn University and has graduated with Bachelor of Science Degree with Second Class Honors in Biochemistry since July, 2002. Since then, she had been studying for a Master Degree of Science in Biotechnology, Faculty of Science, Chulalongkorn University.

