

REFERENCE

- Ahn J, S., Choi, H.K., Lee, K.H., Nahm, J.H., and Cho, C. S. (2001) Novel mucoadhesive polymer prepared by template polymerization of acrylic acid in the presence of silk sericin. *Journal of Applied Polymer Science*, 80, 274-280.
- Ajibola, C.F., Fashakin, J.B., Fagbemi, T.N. and Aluko, R.E. (2011) Effect of Peptide Size on Antioxidant Properties of African Yam Bean Seed (*Sphenostylis stenocarpa*) Protein Hydrolysate Fractions. *International Journal of Molecular Sciences*, 12, 6685-6702.
- Akturk, O., Tezcaner, A., Bilgili, H., Deveci, M.S., Gecit, M.R. and Keskin, D. (2011) Evaluation of sericin/collagen membranes as prospective wound dressing biomaterial. *Journal of Bioscience and Bioengineering*, 112, 279-288.
- Alvarez, O.M., Patel, M., Booker, J., and Markowitz, L. (2004) Effectiveness of a biocellulose wound dressing for the treatment of chronic venous leg ulcers: Results of a single center randomized study involving 24 patients. *Wounds*, 16, 224-233.
- Ang-atikarnkul, P. (2011) Preparation and Characterization of Cellulose Whisker/Chitin Whisker/Silk Sericin Bionanocomposite Sponges for Potential Wound Dressing Applications. M.S. Thesis, The Petroleum and Petrochemical College, Chulalongkorn University, Bangkok, Thailand.
- Aramwit, P., and Sangcakul, A. (2007) The effects of sericin cream on wound healing in rats. *Bioscience, Biotechnology, and Biochemistry*, 71, 2473-2477.
- Aramwit, P., Kanokpanont, S., De-Eknamkul, W. and. Srichana, T. (2009) Monitoring of inflammatory mediators induced by silk sericin. *Journal of Bioscience and Bioengineering*, 107, 556-561.

- Aramwit, P., Kanokpanont, S., De-Eknakul, W., and Srichana, T. (2009). Monitoring of inflammatory mediators induced by silk sericin. *Journal of Bioscience and Bioengineering*, 107, 556-561.
- Aramwit, P., Kanokpanont, S., Nakpheng, T., and Srichana, T. (2010). The Effect of Sericin from Various Extraction Methods on Cell Viability and Collagen Production. *International Journal of Molecular Sciences*, 11, 2200-2211.
- Aramwit, P., Kanokpanont, S., Punyarit, P., Srichana, T. (2009). Effectiveness of Inflammatory Cytokines Induced by Sericin Compared to Sericin in Combination With Silver Sulfadiazine Cream on Wound Healing. *Wounds*, 21, 198–206.
- Aramwit, P., Sirintientonga, T., Kanokpanontb, S., Srichanac, T. (2010) Formulation and characterization of silk sericin–PVA scaffold crosslinked with genipin. *International Journal of Biological Macromolecules*, 47, 668–675.
- Baklagina, Y.G., Khripunov, A.K., Tkachenko, A.A., Kopeikin, V.V., Matveeva, N.A., Lavrent'ev, V.K., Nilova, V.K., Sukhanova, T.E., Smyslov, R.Y., Zanaveskina, I.S., Klechkovskaya, and V.V., Feigin, L.A. (2005) Sorption Properties of Gel Films of Bacterial Cellulose. *Russian Journal of Applied Chemistry*, 78, 1176-1181.
- Bharathi, R.V., Veni, B.K., Jayashree, Suseela, L., and Thiruma, M. (2010) Antioxidant and wound healing studies on different extracts of *Stereospermum colais* leaf. *International Journal of Research in Pharmaceutical Sciences*. 1, 435-439.
- Bhuvanesh, G. (2010) Textile-based smart dressing. *Indian Journal of Fibre & Textile Research*, 35, 174-187.
- Bielecki S, Krystynowicz A, Turkiewicz M, Kalinowska H. (Ed.) (2002) *Bacterial Cellulose: A Biopolymers Polysaccharides I* (pp. 37-90). Germany: Steinbuchel Munster.
- Boateng, J.S., Matthews, K.H., Stewens , H.N.E., and Eccleston G.M. (2008) Wound healing dressings and drug delivery systems: A Review. *Journal of Pharmaceutical Sciences*, 97, 2892-2923.

- Brett, D. W. (2006) A Review of Moisture-Control Dressings in Wound Care. *Journal of Wound Ostomy Continence Nursing*, 33, S3-S8.
- Brown, A. J., (1886) An acetic ferment which forms cellulose. *Journal of Chemical Society*, 49, 432-439.
- Byrom, D. (Ed.) (1991) Microbial cellulose, Biomaterials: Novel materials from biological sources (pp. 263-284). New York: Stockton Press.
- Chaplin, M. (2009). "Water structure and science." Lsbu. 23 Apr 2013. <<http://www1.lsbu.ac.uk/water/hycmc.html>>, 23 April 2013.
- Chen, H.M., Muramoto, K., Yamauchi, F., and Nokihara, K. (1996) Antioxidant activity of designed peptides based on the antioxidative peptide isolated from digests of a soybean protein. *Journal of Agricultural and Food Chemistry*, 44, 2619–2623.
- Chen, W.Y.J., Rogers, A.A., and Lydon, M.J. (1992). Characterization of biologic properties of wound fluid collected during early stages of wound healing. *Journal of Investigative Dermatology*, 99, 559-564.
- Czaja, W., Krystynowicz, A., Bielecki, S., and Brown, R.M. (2006) Microbial cellulose—the natural power to heal wounds. *Biomaterials*, 27, 145–151.
- Czaja, W., Krystynowicz, A., Kawecki, M., Wysota, K., Sakiel, S., Wróblewski, P., Glik, J., Nowak, M., and Bielecki, S. (2007) Biomedical Applications of Microbial Cellulose in Burn Wound Recovery. *Cellulose: Molecular and Structural Biology*, 307–321.
- Czaja, W., Romanowicz, D., and Brown, R.M. (2004) Structural investigations of microbial cellulose produced in stationary and agitated culture. *Cellulose*, 11,403-411.
- Czaja, W., Young, D. J., Kawecki, M., and Brown, R.M. (2007) The Future prospects of microbial cellulose in biomedical applications. *Biomacromolecules*, 8, 1-12.
- Dash, R., Acharya, C., Bindu, P.C., and Kundu S.C. (2007) Antioxidant potential of silk protein sericin against hydrogen peroxide-induced oxidative stress in skin fibroblasts. *BMB reports*, 236-241.

- De Jesus-Gallardo, E., Andres, R.M., and Magno, E.T. (1971) A study on the isolation and screening of microorganism for production of diversed textured Nata. Philippine Journal of Science, 100, 41-51.
- Eming, S.A., Smola, H., Krieg, T. (2002) Treatment of chronic wounds: state of the art and future concepts. Cells Tissues Organs, 172, 105–117.
- Farah, L.F.X. (1990) Process for the preparation of cellulose film, cellulose film produced thereby, artificial skin graft and its use. U.S. Patent, 4912049.
- Fontana, J.D., de Sousa, A.M., Fontana, C.K., Torriani, I.L., Moreschi ,J.C, Gallotti, B.J., de Souza, S.J., Narcisco, G.P., Bichara, J.A., and Farah, L.F.X. (1990) Acetobacter cellulose pellicle as a temporary skin substitute. Applied Biochemistry and Biotechnology, 24-25, 253–264.
- Gulcin, I. (2007) Comparison of in vitro antioxidant and antiradical activities of L-tyrosine and L-Dopa. Amino Acids, 32, 431-438.
- Halliwell, B. (1991) Reactive oxygen species in living systems: Source biochemistry and role in human disease. American Journal of Medicine, 91, 14-22.
- Hestrin, S. and Schramm, M., (1954) Synthesis of cellulose by Acetobacter xylinum: preparation of freeze dried cells capable of polymerizing glucose to cellulose. Biochemical Journal, 58, 345-352.
- Horii, F., Hirai, A., and Kitamaru, R. (1987) Transformation of native cellulose crystals from cellulose I(β) to I(α) through solid-state chemical reactions. Macromolecules, 20, 1440-1442.
- Houghton P.J., Hylands P.J., Mensah A.Y., Hensel A., and Deters A.M. (2005). In vitro tests and ethnopharmacological investigations: Wound healing as an example. Journal of Ethnopharmacology, 100, 100–107.
- Iguchi, M., Yamanaka, S., and Budhiono, A., (2000) Review Bacterial Cellulose—a masterpiece of nature's arts. Journal of Materials Science, 35, 261-270.
- Ilango, K. and Chitra, V. (2010) Wound Healing and Anti-oxidant Activities of the Fruit Pulp of *Limonia Acidissima* Linn (Rutaceae) in Rats Tropical. Journal of Pharmaceutical, 9, 223-230.

- Jaehwan K., Zhijiang C., Hyun S. L., Gwang S. C., Don H. L., and Chulhee J. (2011) Preparation and characterization of a Bacterial cellulose/Chitosan composite for potential biomedical application. *Journal of Polymer Research*, 18,739-744.
- Jae-Young, J. and Se-Kwon, K. (2012) Chitosan as Potential Marine Nutraceutical. *Advances in Food and Nutrition Research*, 65, 121-135.
- Jen Ming, Y. and Hao Tzu, L. (2004) Properties of chitosan containing PP-g-AA-g-NIPAAm bigraft nonwoven fabric for wound dressing. *Journal of Membrane Science*, 243, 1-7.
- Jonkman, M.F., Hoeksma, E.A., and Nieuwenhuis, P. (1990) Accelerated epithelialization under a highly vapor-permeable wound dressing is associated with increased precipitation of fibrin (ogen) and fibronectin. *Journal of Investigative Dermatology*, 94, 477-484.
- Jyh-Ping, C., Gwo-Yun C., and Jan-Kan C. (2008) Electrospun collagen/chitosan nanofibrous membrane as wound dressing. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 313–314,183–188.
- Kato, N., Sato, S., Yamanaka A, Yamada H, Fuwa N, Nomura M. (1998) Silk protein, sericin, inhibits lipid peroxidation and tyrosinase activity. *Bioscience, Biotechnology, and Biochemistry*, 62, 145-147.
- Klemm, D., Schumann, D., Udhardt, U., and Marsch, S. (2001) Bacterial synthesized cellulose - artificial blood vessels for microsurgery. *Progress in Polymer Science*, 26, 1561-1603.
- Kong, J., and Yu, S. (2007). Fourier Transform Infrared Spectroscopic Analysis of Protein Secondary Structures. *Acta Biochimica et Biophysica Sinica*, 39,549-559.
- Korsmeyer, R.W., Gurny, R., Doelker, E., Buri, P., and Peppas, N.A., 1983. Mechanism of solute release from porous hydrophilic polymers. *International Journal of Pharmaceutics*, 15, 25-35.

- Kundu, S.C., Dash, B.C., Dash R., and Kaplan D.L. (2008) Natural protective glue protein, sericin bioengineered by silkworms: Potential for biomedical and biotechnological applications. *Progress in Polymer Science*, 33, 998-1012.
- Lagana, G., and Anderson, E.H. (2010). Moisture Dressings: The New Standard in Wound Care. *The Journal for Nurse Practitioners*, 6, 366-370.
- Lehninger, A. L., (3rd Ed.) (1971). *Biochemistry: the molecular basis of cell structure and function*. Worth Publishers.
- Lou, C.W. (2008) Process Technology and Properties Evaluation of a Chitosan-coated Tencel/cotton Nonwoven Fabric as a Wound Dressing. *Fibers and Polymers*, 9,286-292.
- Majeti N.V. Ravi Kumar (2000) A review of chitin and chitosan applications. *Reactive & Functional Polymers*, 46, 1-27.
- Mandal, B.B., Ghosh, B., and Kundu, S.C. (2011) Non-mulberry silk sericin/poly (vinyl alcohol) hydrogel matrices for potential biotechnological applications. *International Journal of Biological Macromolecules*, 49, 125-133.
- Mandal, B.B., Priya, A.S., and Kundu, S.C. (2009) Novel silk sericin/gelatin 3-D scaffolds and 2-D films: Fabrication and characterization for potential tissue engineering applications. *Acta Biomaterialia*, 5, 3007-3020.
- Maneerung, S., Tokura, S., and Rujiravanit, R. (2008) Impregnation of silver nanoparticles into bacterial cellulose for antimicrobial wound dressing. *Carbohydrate Polymers*, 72, 43-51.
- Maneerung, T. (2007) Impregnation of silver nanoparticles into bacterial cellulose for antimicrobial wound dressing. M.S. Thesis, The Petroleum and Petrochemical College, Chulalongkorn University, Bangkok, Thailand.
- Mendis, E., Rajapakse, N., Byun, H. G., and Kim, S. K. (2005) Investigation of Jumbo squid (*Dosidicus gigas*) skin gelatin peptides for their in vitro antioxidant effects. *Life Sciences*, 77, 2166–2178.
- Meyer, K., Rajanahalli, P., Ahamed, M, Rowe, J.J., and Hong, Y. (2011) ZnO nanoparticles induce apoptosis in human dermal fibroblasts via p53 and p38 pathways. *Toxicology in Vitro*, 25, 1721-1726.

- Moustafa M.G. Fouda, R. Wittke, D. Knittel, and E. Schollmeyer (2009) Use of chitosan/polyamine biopolymers based cotton as a model system to prepare antimicrobial wound dressing. *International Journal of Diabetes Mellitus*, 1, 61–64.
- Muenduen, P., and Nirun, J. (2008) Biosynthesis and characterization of bacteria cellulose–chitosan film. *Carbohydrate Polymers*, 74, 482–488.
- Nakayama, A., Kakugo, A., Gong , J.P., and Osada, Y. (2004) High mechanical strength double-network hydrogel with bacterial cellulose. *Advance Functional Materials*, 14, 1124–1128.
- Namviriyachote, N., Bang, N., and Aramwit, P. (2009) Sericin film: Influence of concentration on its physical properties. *World Academy of Science, Engineering and Technology*, 56, 7-9.
- Nishiyama. T., Tsunenaga, M., Akutsu, N., Horii, I., Nakayama, Y., Adachi, E., Yamato, M., and Hayashi, T. (1993) Dissociation of actin microfilament organization from acquisition and maintenance of elongated shape of human dermal fibroblasts in three-dimensional collagen gel. *Matrix*, 13, 447-455.
- Padamwar, M.N., and Pawar, A.P. (2004) Silk sericin and its applications: A review. *Journal of Scientific & Industrial Research*, 63, 323-329.
- Padamwar, M.N., Pawar, A.P., Daithankar, A.V., and Mahadik, K.R. (2005) Silk sericin as a moisturizer: an in vivo study. *Journal of Cosmetic Dermatology*, 4, 250–257.
- Philip, M. (2004) Use of DPPH to estimate antioxidant activity. *The Songklanakarin Journal of Science and Technology*, 26(2), 211-219.
- Phisalaphong, M., Jatupaiboon, N. (2008) Biosynthesis and characterization of bacteria cellulose-chitosan film. *Carbohydrate Polymers*, 74, 482-488.
- Phisalaphong, M., Suwanmajo, T., and Thamarat P. (2008) Synthesis and Characterization of Bacterial Cellulose/ Alginate Blend Membrane. *Journal of Applied Polymer Science*, 107, 3419-3424.
- Pittayaapipon, P. (2012) Preparation of Bacterial Cellulose Containing Silk Sericin for Wound Dressing Application. M.S. Thesis, The Petroleum and Petrochemical College, Chulalongkorn University, Bangkok, Thailand.

- Razzak, M.T., Darwis, D., Zainuddin, Sukirno. (2001) Irradiation of polyvinyl alcohol and polyvinyl pyrrolidone blended hydrogel for wound dressing. Radiation Physics and Chemistry, 62, 107-113.
- Rejane C. Goy, Douglas de Britto, & Odilio B. G. Assis (2009) A Review of the Antimicrobial Activity of Chitosan. Polímeros: Ciência e Tecnologia, 19, 241-247.
- Rigueiro, J.P., Elices, M., Llorca, J. and Viney, C. (2001) Tensile Properties of Silkworm Silk obtained by Forced Silking, Journal of Applied Polymer Science, 82, 1928-1935.
- Ronge, X., Song, L., Zhanyong, G., Huahua, Y., Pibo, W., Cuiping, L., Zhien, L., and Pengcheng, L. (2005) Relevance of molecular weight of chitosan and its derivatives and their antioxidant activities in vitro. Bioorganic & Medicinal Chemistry, 13, 1573–1577.
- Ross, P., Mayer, R., and Benziman, M. (1991) Cellulose biosynthesis and function in bacterial. Microbiological Review, 55, 35-58.
- Saibuatong, O. A., and Phisalaphong, M. (2010) Novo aloe vera-bacterial cellulose composite film from biosynthesis. Carbohydrate Polymers, 79, 455-460.
- Sakkawet, Y., Orathai, P., and Ratana, R. (2012) Preparation and Characterization of Chitosan-Coated DBD Plasma-Treated Natural Rubber Latex Medical Surgical Gloves with Antibacterial Activities. Plasma Chemistry and Plasma Processing, 32, 1275–1292.
- Sarovart, S., Sudatis, B., Meesilpa, P., Grady, B.P., and Magaraphan, R. (2003) The use of sericin as an antioxidant and antimicrobial for polluted air treatment. Reviews on Advanced Materials Science, 5, 193-198.
- Sarovart, S., Sudatis, B., Meesilpa, P., Grady, B.P., and Magaraphan, R. (2003) The use of sericin as an antioxidant and antimicrobial for polluted air treatment. Reviews on Advanced Materials Science, 5, 193-198.
- Senakoon, W., Nuchadomrong, S., Sirimungkararat, S., Senawong, T., and Kitikoon, P. (2009) Antibacterial action of eri (samia ricini) sericin against escherichia coli and staphylococcus aureus. Journal of Food and Agro-Industry, Special issue, 222-228.

- Senakoon, W., Nuchadomrong, S., Sirimungkararat, S., Senawong, T., and Kitikoon, P. (2009) Antibacterial action of eri (*samia ricini*) sericin against *escherichia coli* and *staphylococcus aureus*. Journal of Food and Agro-Industry, Special issue, 222-228.
- Shin-Yeu Ong, Jian Wu, Shabbir M. Moochhala, Mui-Hong Tan, and Jia Lu (2008) Development of a chitosan-based wound dressing with improved hemostatic and antimicrobial properties. Biomaterials, 29, 4323-4332.
- Shuichi Aoyagi, Hiraku Onishi, and Yoshiharu Machida (2007) Novel chitosan wound dressing loaded with minocycline for the treatment of severe burn wounds. International Journal of Pharmaceutics, 330, 138–145.
- Sjöestrom, E. (2nd Ed.) (1993) Wood chemistry: Fundamentals and Applications.
- Stojadinovic, A., Carlson, J. W., Schultz G.S, Davis, T.A., and Elster. E.A. (2008) Topical advances in wound care. Gynecologic Oncology, 111, 70-80.
- Surma-Slusarska, B., Presler, S., and Danielewicz, D. (2008) Characteristics of bacterial cellulose obtained from *Acetobacter xylinum* culture for application in papermaking. Fibres & Textiles, 16,108-111.
- T. Takei, H. Nakahara, H. Ijima, and K. Kawakami (2012). Synthesis of a chitosan derivative soluble at neutral pH and gellable by freeze-thawing, and its application in wound care. Acta Biomaterialia, 8, 686–693.
- Tao, W., Xiao-Kang, Z., Xu-Ting, X., and Da-Yang, W. (2012) Hydrogel sheets of chitosan, honey and gelatin as burn wound dressings. Carbohydrate Polymers, 88, 75–83.
- Tao, W., Li, M., and Xie, R. (2005) Preparation and Structure of Porous Silk Sericin Materials. Macromolecular Materials and Engineering, 290, 188-194.
- Terada, S., Nishimura, T., Sasaki, M., Yamada, H., and Miki, M. (2002) Sericin, a protein derived from silkworms, accelerates the proliferation of several mammalian cell lines including a hybridoma. Cytotechnology, 40, 3–12.

- Teramoto, H., Kameda, T., and Tamada, Y., (2008) Preparation of gel film from *Bombyx mori* silk sericin and its characterization as a wound dressing. Bioscience, Biotechnology, and Biochemistry, 72, 3189-3196.
- Teramoto, H., Nakajima, K., and Takabayashi, C. (2005) Preparation of elastic silk sericin hydrogel. Bioscience, Biotechnology, and Biochemistry, 69, 845-847.
- Tokoh, C., Takabe, K., Fujita, M., and Saiki, H. (1998) Cellulose synthesized by *Acetobacter xylinum* in the presence of acetyl glucosaminan. Cellulose, 5, 249–261.
- Tsubouchi, K., Igarashi, Y., Takasu, Y., and Yamada, H. (2005). Sericin enhances attachment of cultured human skin fibroblasts. Bioscience, Biotechnology, and Biochemistry, 69, 403-405.
- Twentyman, P.R. and Luscombe, M. (1987) A study of some variables in a tetrazolium dye (MTT) based assay for cell growth and chemosensitivity. British Journal of Cancer, 56, 279-285.
- Valla, S., and Kjosbakken J (1982) Cellulose-negative mutants of *Acetobacter xylinum*. Journal of General Microbiology. 128, 1401 – 1408.
- Vandamme, E. J., Beats, S. D., Vanbalen , A., Joris, K. and Wulf, P. D. (1998) Improved production of bacterial cellulose and its application potential. Polymer Degradation and Stability, 59, 93-99.
- Velnar, T., Bailey, T., and Smrkolj, V. (2009) The wound healing process: an overview of the cellular and molecular mechanisms. The Journal of International Medical Research, 37, 1528-1542.
- Voegeli, R., Meier, J. and Blust, R. (1993) Sericin silk protein: unique structure and properties. Cosmetics & Toiletries, 108, 101-108.
- Watanabe, K., Tabuchi, M., Morinaga, Y., and Yoshinaga, F. (1998) Structural features and properties of bacterial cellulose produced in agitated culture. Cellulose, 5, 187-200.
- Wei, B., Yang,G., and Hong, F. (2011) Preparation and evaluation of a kind of bacterial cellulose dry films with antibacterial properties. Carbohydrate Polymers, 84, 533-538.

- Wild, T., Rahbarnia, A., Kellner, M., Sobotka, L., and Thomas E.M.D. (2010) Basics in nutrition and wound healing. *Nutrition*, 26, 862–866.
- Williams, W.S. and Cannon, R.E., (1989) Alternative environmental roles for cellulose produced by *Acetobacter xylinum*. *Applied and Environmental Microbiology*, 55, 2448-2452.
- Winter, G.D. (1962) Formation of the Scab and the Rate of Epithelization of Superficial Wounds in the Skin of the Young Domestic Pig. *Nature*, 193, 293-294.
- Witthayaprapakorn, C. (2011) Design and Preparation of Synthetic Hydrogels Via Photopolymerisation for Biomedical Use as Wound Dressings. *Procedia Engineering*, 8,286-291.
- Wu, J.H., Wang, Z. and Xu, S.Y. (2007) Preparation and Characterization of Sericin Powder Extracted from Silk Industry Wastewater. *Food Chemistry*, 103 1255–1262.
- Wu, P., Fisher, AC., Foo, P.P., Queen, D. and Gaylor, J.D.S. (1995). In Vitro assessment of water vapour transmission of synthetic wound dressings. *Biomaterials*, 16, 171–175.
- Yamanaka, S., Watanabe, K., Kitamura, N., Iguchi, M., Mitsuhashi, S., Nishi, Y. and Uryu, M. (1989) The structure and mechanical properties of sheets prepared from bacterial cellulose. *Journal of Materials Science*, 24, 3141-3145.
- Yamanaka, S., Watanabe, K., Kitamura, N., Iguchi, M., Mitsuhashi, S., Nishi, Y., and Uryu, M. (1989). The structure and mechanical properties of sheets prepared from bacterial cellulose. *Journal of Materials Science*, 24, 3141-3145.
- Yang, Y.K., Park, S.H., Hwang, J.W., Pyun, Y.R., and Kim, Y.S. (1997) Cellulose production by *Acetobacter xylinum* BRCS under agitated condition. *Journal of Fermentation and Bioengineering*, 85, 312-317.
- Yoshinaga, F., Tonouchi, N., and Watanabe, K. (1997) Research progress in production of bacterial cellulose by aeration and agitation culture and its application as a new industrial material. *Bioscience, Biotechnology, and Biochemistry*, 61, 219-224.

- Zhang, Y.Q. (2002) Applications of natural silk protein sericin in biomaterials. Biotechnology Advances, 20, 91-100.
- Zhijiang, C., and Guang, Y. (2011) Bacterial cellulose/collagen composite: Characterization and first evaluation of cytocompatibility. Journal of Applied Polymer Science, 120, 2938-2944.
- Zhijiang, C., Chengwei, H., and Guang, Y. (2011) Preparation and characterization of a Bacterial cellulose/Chitosan composite for potential biomedical application. Journal of Applied Polymer Science, 121, 1488-1494.
- Zhu, L. J, Arai, M., and Hirabayashi, K. (1996) Sol-gel transition of sericin. Nippon Sanshigaku Zasshi, 65(4) 270-274.
- Zhu, L.J., Yao, J., and Youlu, L. (1998) Structural transformation of sericin dissolved from cocoon layer in hot water. Zhejiang Nongye Daxue Xuebao, 24, 268-272.

APPENDICES

Appendix A Nitrogen Content of Different Chitosan/Sericin Blend Ratio in Bacterial Cellulose

The amount of chitosan and sericin incorporated into bacterial cellulose pellicle was determined by using Kjeldahl method for determining the nitrogen contents of chitosan and sericin.

Table A1 Nitrogen content of chitosan-sericin solution different chitosan-sericin blend ratio

Chitosan/Sericin Blend Ratio	Nitrogen Content in Solution (mg)				
	1	2	3	Average	S.D.
100/0	12.71	10.15	16.40	13.09	3.14
75/25	15.98	16.69	10.86	14.51	3.17
50/50	16.61	18.53	23.71	19.62	3.67
25/75	21.37	27.26	18.60	22.41	4.42
0/100	24.64	27.69	21.16	24.50	3.26

Table A2 Nitrogen content of chitosan-sericin different chitosan-sericin blend ratio in bacterial cellulose

Chitosan/Sericin Blend Ratio	Nitrogen Content in Solution (mg)				
	1	2	3	Average	S.D.
100/0	10.72	13.56	7.32	10.53	3.12
75/25	14.41	15.69	8.882	12.99	3.62
50/50	15.69	17.68	20.38	17.92	2.35
25/75	19.10	20.94	22.79	20.94	1.84
0/100	21.23	22.36	19.67	21.09	1.35

The chitosan and sericin-incorporated bacterial cellulose was immersed in 1% NaOH to remove sericin out from bacterial cellulose matrix. The bacterial cellulose pellicles are containing chitosan and sericin dissolved in NaOH solution. The amount of chitosan and sericin was determined by using Kjeldahl method for determining the nitrogen contents of chitosan and sericin.

Table A3 Chitosan content in different chitosan-sericin blend ratio in bacterial cellulose

Chitosan/Sericin Blend Ratio	Nitrogen Content in Solution (mg)				
	1	2	3	Average	S.D.
100/0	11.32	9.98	9.53	10.28	0.92
75/25	10.09	8.51	9.78	9.46	0.83
50/50	8.80	9.23	8.57	8.86	0.33
25/75	5.32	4.97	5.54	5.27	0.28
0/100	0	0	0	0	0

Table A4 Sericin content in NaOH solution in different chitosan-sericin blend ratio

Chitosan/Sericin Blend Ratio	Nitrogen Content in Solution (mg)				
	1	2	3	Average	S.D.
100/0	0	0	0	0	0
75/25	2.98	3.21	3.07	3.08	0.11
50/50	8.32	8.99	9.51	8.94	0.59
25/75	15.79	14.09	16.21	15.36	1.12
0/100	21.97	21.03	19.87	20.95	1.05

Appendix B Antioxidant Activity of Chitosan and Sericin-incorporated Bacterial Cellulose

The antioxidant activity of chitosan and sericin-incorporated bacterial cellulose was determined based on the 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical scavenging activity method.

Table B1 Antioxidant activity of chitosan and sericin-incorporated bacterial cellulose

Chitosan/Sericin Blend Ratio		Scavenging activity (%)				
		1	2	3	Average	S.D.
Chitosan A in BC	100/0	7.38	9.04	8.76	8.39	0.89
	75/25	11.03	12.65	10.72	11.47	1.03
	50/50	13.46	13.50	13.67	13.54	0.11
	25/75	15.02	15.50	15.70	15.41	0.35
Chitosan B in BC	100/0	6.92	5.30	6.01	6.08	0.80
	75/25	9.65	10.08	9.54	9.76	0.28
	50/50	11.78	12.09	11.98	11.95	0.15
	25/75	13.27	12.49	12.48	12.75	0.44
Sericin in BC	0/100	17.81	18.60	18.03	18.15	0.40

Appendix C Antibacterial Activity of Chitosan and Sericin-incorporated Bacterial Cellulose

The antibacterial activity of chitosan and sericin-incorporated bacterial cellulose was determined based on colony forming count method.

Table C1 Antibacterial Activity of chitosan and sericin-incorporated bacterial cellulose against *S. aureus* at 1.8×10^{-8} cfu/mL.

Chitosan/Sericin Blend Ratio		Bacterial Reduction Rate (%)				
		1	2	3	Average	S.D.
Chitosan A in BC	100/0	77.14	66.28	70.85	71.42	5.45
	75/25	60.00	61.14	52.00	57.71	4.61
	50/50	42.28	44.00	49.14	45.14	3.56
	25/75	28.57	26.28	36.57	30.47	5.40
Chitosan B in BC	100/0	54.28	45.71	49.142	49.71	4.31
	75/25	42.28	30.85	27.42	33.52	7.77
	50/50	23.42	26.28	14.85	21.52	5.94
	25/75	8.00	10.28	17.14	11.80	4.75
Sericin in BC	0/100	9.14	7.42	15.42	10.66	4.21

Table C2 Antibacterial Activity of chitosan and sericin-incorporated bacterial cellulose against *E. coli* at 1.8×10^{-8} cfu/mL.

Chitosan/Sericin Blend Ratio		Bacterial Reduction Rate (%)				
		1	2	3	Average	S.D.
Chitosan A in BC	100/0	71.64	54.63	65.40	63.89	8.60
	75/25	46.12	44.42	55.19	48.58	5.43
	50/50	41.02	38.18	36.48	38.56	2.29
	25/75	34.21	28.54	34.78	32.51	3.44
Chitosan B in BC	100/0	37.61	46.12	29.11	37.61	8.50
	75/25	29.11	22.87	21.17	24.38	4.180
	50/50	23.44	15.50	10.96	16.63	6.31
	25/75	6.42	14.93	12.09	11.15	4.33
Sericin in BC	0/100	7.561	12.66	1.89	7.37	5.39

Appendix D Water Vapor Transmission Rate

Table D1 Water vapor transmission rate of chitosan and sericin-incorporated bacterial cellulose

Chitosan/Sericin Blend Ratio		Water vapor transmission rate (g/m ² /hour)				
		1	2	3	Average	S.D.
Pure BC		1834.61	1861.61	2331.05	2009.09	279.15
Chitosan A in BC	100/0	1554.11	1793.99	1508.87	1618.99	153.23
	75/25	1440.26	1608.30	1506.39	1518.32	84.65
	50/50	1590.90	1395.64	1364.82	1450.46	122.60
	25/75	1391.92	1363.08	1372.15	1365.72	14.74
Chitosan B in BC	100/0	1165.54	1201.23	1535.67	1300.81	204.17
	75/25	1034.45	1294.56	1357.9	1228.97	171.41
	50/50	1096.65	1236.08	1145.83	1159.52	70.71
	25/75	1022.42	1129.53	1134.09	1095.34	63.19
SS in BC	0/100	1466.12	1253.83	1337.10	1352.3555	106.96

Appendix E Sericin Releasing Behavior

Table E1 Sericin releasing profile of 100/0 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
0.166	-0.0275	-0.0534	-0.0369	0.0159
	-0.0104	-0.0216		
	-0.0179	-0.0356		
0.333	-0.0111	-0.0247	-0.0165	0.0074
	-0.0040	-0.0105		
	-0.0059	-0.0143		
0.5	-0.0108	-0.0241	-0.0086	0.0149
	0.0041	0.0056		
	-0.0024	-0.0073		
0.75	-0.0043	-0.0111	-0.0086	0.0144
	-0.0095	-0.0215		
	0.0047	0.0068		
1	-0.0084	-0.0193	-0.0028	0.0149
	0.0019	0.0012		
	0.0061	0.0096		
1.30	-0.0030	-0.0085	-0.0031	0.0080
	-0.0021	-0.0067		
	0.0043	0.0060		
2	0.0028	0.0030	0.0123	0.0082
	0.0088	0.0150		
	0.0107	0.0188		
3	0.0085	0.0144	0.0091	0.0053
	0.0032	0.0038		
	0.0057	0.0088		

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
5	-0.0063	-0.0151	0.0042	0.0168
	0.0090	0.0154		
	0.0074	0.0122		
8	-0.0053	-0.0131	0.0028	0.0139
	0.0073	0.0120		
	0.0060	0.0094		
12	-0.0045	-0.0115	-0.0002	0.0108
	0.0018	0.0010		
	0.0062	0.0098		
24	0.0022	0.0018	0.0060	0.0099
	0.0099	0.0172		
	0.0007	-0.0011		
36	0.0017	0.0008	0.0103	0.0084
	0.0078	0.0130		
	0.0098	0.0170		
48	0.0058	0.0090	0.0165	0.0099
	0.0151	0.0276		
	0.0076	0.0126		
60	0.0080	0.0130	0.0225	0.0088
	0.0133	0.0240		
	0.0165	0.0304		
72	0.0110	0.0188	0.0271	0.0080
	0.0150	0.0274		
	0.0187	0.0348		

Table E2 Sericin releasing profile of 75/25 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
0.166	-0.0040	-0.0098	0.0137	0.0206
	0.0165	0.0283		
	0.0135	0.0227		
0.333	0.0079	0.0132	0.0259	0.0110
	0.0169	0.0312		
	0.0179	0.0332		
0.5	0.0111	0.0196	0.0283	0.0115
	0.0132	0.0238		
	0.0219	0.0412		
0.75	0.0122	0.0218	0.0288	0.0134
	0.0114	0.0202		
	0.0234	0.0442		
1	0.0155	0.0284	0.0467	0.0159
	0.0287	0.0548		
	0.0297	0.0568		
1.30	0.0170	0.0314	0.0384	0.0171
	0.0142	0.0258		
	0.0302	0.0578		
2	0.0126	0.0226	0.0405	0.0166
	0.0230	0.0434		
	0.0290	0.0554		
3	0.0487	0.0948	0.0761	0.0163
	0.0341	0.0656		
	0.0352	0.0678		

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
5	0.0834	0.1642	0.1038	0.0524
	0.0375	0.0724		
	0.0386	0.0746		
8	0.0772	0.1518	0.1183	0.0293
	0.0505	0.098		
	0.0535	0.1044		
12	0.1306	0.2586	0.1961	0.0545
	0.0805	0.1584		
	0.0869	0.1712		
24	0.1206	0.2786	0.2708	0.0288
	0.1407	0.2388		
	0.1487	0.2948		
36	0.1569	0.3112	0.3271	0.0141
	0.1673	0.3320		
	0.1703	0.3380		
48	0.1670	0.3314	0.3367	0.0045
	0.1710	0.3394		
	0.1708	0.3390		
60	0.1750	0.3474	0.3489	0.0080
	0.1721	0.3416		
	0.1800	0.3574		
72	0.1760	0.3494	0.3513	0.0094
	0.1727	0.3428		
	0.182	0.3614		

Table E3 Sericin releasing profile of 50/50 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
0.166	0.0060	0.0088	0.0233	0.0153
	0.0223	0.0392		
	0.0130	0.0218		
0.333	0.0101	0.0177	0.0339	0.0171
	0.0271	0.0517		
	0.0174	0.0323		
0.5	0.0193	0.0361	0.0546	0.0192
	0.0385	0.0745		
	0.0279	0.0533		
0.75	0.0309	0.0593	0.0740	0.0257
	0.0531	0.1037		
	0.0308	0.0591		
1	0.0284	0.0543	0.0745	0.0228
	0.05090	0.0993		
	0.0362	0.0699		
1.30	0.03930	0.0761	0.0848	0.0077
	0.0451	0.0877		
	0.0466	0.0907		
2	0.0422	0.0819	0.1054	0.0204
	0.0604	0.1183		
	0.0593	0.1161		
3	0.0854	0.1683	0.1421	0.0227
	0.0654	0.1283		
	0.0661	0.1297		

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
5	0.1672	0.3319	0.2552	0.0665
	0.1118	0.2211		
	0.1076	0.2127		
8	0.1920	0.3815	0.3121	0.0646
	0.15189	0.3013		
	0.1281	0.2537		
12	0.2555	0.5087	0.4250	0.0791
	0.2087	0.4149		
	0.1770	0.3515		
24	0.2741	0.5457	0.5135	0.0381
	0.2630	0.5235		
	0.2370	0.4715		
36	0.2753	0.5483	0.5675	0.0457
	0.3111	0.6197		
	0.2685	0.5345		
48	0.2863	0.5701	0.5948	0.0492
	0.3270	0.6515		
	0.2827	0.5629		
60	0.2860	0.5695	0.6149	0.0411
	0.3260	0.6495		
	0.3142	0.6259		
72	0.2840	0.5655	0.6401	0.0658
	0.3340	0.6655		
	0.3460	0.6895		

Table E4 Sericin releasing profile of 25/75 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
0.166	0.0168	0.0288	0.0296	0.0061
	0.0142	0.0240		
	0.0207	0.0361		
0.333	0.0618	0.1211	0.1167	0.0079
	0.0551	0.1077		
	0.0620	0.1215		
0.5	0.0872	0.1719	0.1710	0.0067
	0.0899	0.1773		
	0.0832	0.1639		
0.75	0.1183	0.2341	0.2595	0.0223
	0.1393	0.2761		
	0.1354	0.2683		
1	0.1250	0.2475	0.2451	0.0176
	0.1145	0.2265		
	0.1320	0.2615		
1.30	0.1190	0.2355	0.2531	0.0268
	0.1212	0.2399		
	0.1432	0.2839		
2	0.1332	0.2639	0.2799	0.0246
	0.1351	0.2677		
	0.1554	0.3083		
3	0.1878	0.3731	0.3397	0.0376
	0.1507	0.2989		
	0.1748	0.3471		

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
5	0.1893	0.3761	0.3382	0.0378
	0.1515	0.3005		
	0.1703	0.3381		
8	0.2839	0.5653	0.4950	0.0637
	0.2406	0.4787		
	0.2218	0.4411		
12	0.3534	0.7043	0.6491	0.0571
	0.3276	0.6527		
	0.2964	0.5903		
24	0.3758	0.7491	0.7761	0.0246
	0.3999	0.7973		
	0.3922	0.7819		
36	0.3869	0.7713	0.8060	0.0306
	0.4102	0.8179		
	0.4157	0.8289		
48	0.3844	0.7663	0.8113	0.0409
	0.4121	0.8217		
	0.4243	0.8461		
60	0.3860	0.7695	0.8229	0.0504
	0.4161	0.8297		
	0.4361	0.8697		
72	0.3971	0.7917	0.8337	0.0415
	0.4187	0.8349		
	0.4386	0.8747		

Table E5 Sericin releasing profile of 0/100 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
0.166	0.0154	0.0262	0.0328	0.0056
	0.0207	0.0361		
	0.0207	0.0361		
0.333	0.0393	0.0761	0.0729	0.0103
	0.0320	0.0615		
	0.0420	0.0815		
0.5	0.0799	0.1573	0.1549	0.0101
	0.0732	0.1439		
	0.0832	0.1639		
0.75	0.1108	0.2191	0.2518	0.0284
	0.1354	0.2683		
	0.1354	0.2683		
1	0.1534	0.3043	0.3327	0.0247
	0.1748	0.3471		
	0.1748	0.3471		
1.30	0.1677	0.3329	0.3363	0.0030
	0.1703	0.3381		
	0.1703	0.3381		
2	0.1720	0.3415	0.3545	0.0113
	0.1818	0.3611		
	0.1818	0.3611		
3	0.2587	0.5149	0.5651	0.0435
	0.2964	0.5903		
	0.2964	0.5903		

Time (hr)	Absorbance at 562 nm	Protein Concentration (mg)	Average	SD
5	0.3774	0.7523	0.7853	0.0286
	0.4022	0.8019		
	0.4022	0.8019		
8	0.3961	0.7897	0.8825	0.0804
	0.4657	0.9289		
	0.4657	0.9289		
12	0.4465	0.8905	0.9542	0.0552
	0.4943	0.9861		
	0.4943	0.9861		
24	0.4854	0.9683	0.9892	0.0207
	0.5061	1.0097		
	0.4961	0.9897		
36	0.4991	0.9957	1.0069	0.0100
	0.5086	1.0147		
	0.5065	1.0105		
48	0.5093	1.0161	1.0249	0.0076
	0.5157	1.0289		
	0.5161	1.0297		
60	0.5152	1.0279	1.0323	0.0076
	0.5152	1.0279		
	0.5218	1.0411		
72	0.5191	1.0357	1.0406	0.0122
	0.5171	1.0317		
	0.5285	1.0545		

Appendix F Chitosan Releasing Behavior

Table F1 Chitosan releasing profile of 100/0 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
0.166	0.0275	3.5694	3.8380	0.2665
	0.0316	4.1024		
	0.0296	3.8424		
0.333	0.0357	4.9919	5.6733	0.9155
	0.0380	5.3139		
	0.0480	6.7139		
0.5	0.0372	5.2019	5.5473	1.0514
	0.0337	4.7119		
	0.0481	6.7279		
0.75	0.0385	5.3839	5.1833	1.1403
	0.0444	6.2099		
	0.0283	3.9559		
1	0.0433	6.0559	7.6939	1.4199
	0.0604	8.4499		
	0.0613	8.5759		
1.30	0.0527	7.3719	7.3953	0.6793
	0.0578	8.0859		
	0.0481	6.7279		
2	0.0778	10.8859	12.3933	1.6450
	0.0868	12.1459		
	0.1011	14.1479		
3	0.0929	12.9999	12.8879	0.6098
	0.0960	13.4339		
	0.0874	12.2299		

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
5	0.1096	15.3379	15.6506	0.4593
	0.1103	15.4359		
	0.1156	16.1779		
8	0.1304	18.2499	17.5079	1.7400
	0.1109	15.5199		
	0.1340	18.7539		
12	0.1379	19.2999	21.2973	1.7480
	0.1611	22.5479		
	0.1575	22.0439		
24	0.1793	25.0959	25.2826	0.5160
	0.1778	24.8859		
	0.1848	25.8659		
36	0.2033	28.4559	28.9133	0.4483
	0.2067	28.9319		
	0.2097	29.3519		
48	0.2167	30.3319	30.9806	0.8493
	0.2191	30.6679		
	0.2282	31.9419		
60	0.2168	30.3459	31.0413	0.8465
	0.2200	30.7939		
	0.2285	31.9839		
72	0.2191	30.6679	30.8546	1.3537
	0.2307	32.2919		
	0.2115	29.6039		

Table F2 Chitosan releasing profile of 75/25 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
0.166	0.0170	2.2044	3.3094	0.9596
	0.0303	3.9334		
	0.0292	3.7904		
0.333	0.0281	3.9279	4.5906	0.6796
	0.0326	4.5579		
	0.0378	5.2859		
0.5	0.0343	4.7959	4.8286	0.1773
	0.0334	4.6699		
	0.0359	5.0199		
0.75	0.0381	5.3279	5.0199	0.3229
	0.0361	5.0479		
	0.0335	4.6839		
1	0.0357	4.9919	5.7666	0.7118
	0.0423	5.9159		
	0.0457	6.3919		
1.30	0.0455	6.3639	7.0686	0.6170
	0.0524	7.3299		
	0.0537	7.5119		
2	0.0639	8.9399	9.2386	1.2589
	0.0759	10.6199		
	0.0583	8.1559		
3	0.0670	9.3739	10.3259	0.8508
	0.0757	10.5919		
	0.0787	11.0119		

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
5	0.0908	12.7059	13.0046	0.3515
	0.0957	13.3919		
	0.0923	12.9159		
8	0.1054	14.7499	16.1406	1.2850
	0.1171	16.3879		
	0.1235	17.2839		
12	0.1216	17.0179	18.8566	1.6660
	0.1378	19.2859		
	0.1448	20.2659		
24	0.1470	20.5739	22.3473	1.6208
	0.1697	23.7519		
	0.1623	22.7159		
36	0.1758	24.6059	25.0773	0.4990
	0.1788	25.0259		
	0.1829	25.5999		
48	0.1869	26.1599	26.5659	0.4628
	0.1934	27.0699		
	0.1891	26.4679		
60	0.1907	26.6919	26.7899	0.4354
	0.1887	26.4119		
	0.1948	27.2659		
72	0.2017	28.2319	27.2146	0.8811
	0.1907	26.6919		
	0.1909	26.7199		

Table F3 Chitosan releasing profile of 50/50 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
0.166	0.0068	0.8784	2.0917	1.0765
	0.0226	2.9324		
	0.0190	2.4644		
0.333	0.0168	2.3459	3.1813	0.8698
	0.0223	3.1159		
	0.0292	4.0819		
0.5	0.0237	3.3119	4.4599	1.0083
	0.0372	5.2019		
	0.0348	4.8659		
0.75	0.0304	4.2499	4.4179	0.3790
	0.0347	4.8519		
	0.0297	4.1519		
1	0.0315	4.4039	4.7773	0.3260
	0.0352	4.9219		
	0.0358	5.0059		
1.30	0.0318	4.4459	4.4179	0.3508
	0.0290	4.0539		
	0.0340	4.7539		
2	0.0347	4.8519	5.5753	0.6404
	0.0415	5.8039		
	0.0434	6.0699		
3	0.0465	6.5039	7.0266	0.6175
	0.0551	7.7079		
	0.0491	6.8679		

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
5	0.0573	8.0159	7.2599	0.6642
	0.0500	6.9939		
	0.0484	6.7699		
8	0.0770	10.7739	11.2639	0.4447
	0.0832	11.6419		
	0.0813	11.3759		
12	0.0857	11.9919	12.1553	0.2709
	0.0891	12.4679		
	0.0858	12.0059		
24	0.1263	17.6759	17.8066	0.2263
	0.1263	17.6759		
	0.1291	18.0679		
36	0.1405	19.6639	20.1679	0.7211
	0.1418	19.8459		
	0.1500	20.9939		
48	0.1613	22.5759	22.7066	0.7921
	0.1571	21.9879		
	0.1683	23.5559		
60	0.1667	23.3319	22.7829	0.6361
	0.1578	22.0859		
	0.1638	22.9310		
72	0.1552	21.7219	22.6366	0.8604
	0.1626	22.7579		
	0.1674	23.4299		

Table F4 Chitosan releasing profile of 25/75 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
0.166	-0.0043	-0.5646	0.1417	0.6621
	0.0058	0.7484		
	0.0019	0.2414		
0.333	0.0156	2.1779	2.4346	0.4446
	0.0156	2.1779		
	0.0211	2.9479		
0.5	0.0335	4.6839	4.7679	1.7935
	0.0216	3.0179		
	0.0472	6.6019		
0.75	0.0292	4.0819	4.3666	0.2737
	0.0331	4.6279		
	0.0314	4.3899		
1	0.0413	5.7759	5.4819	0.4385
	0.0407	5.6919		
	0.0356	4.9779		
1.30	0.0518	7.2459	7.5866	0.3017
	0.0559	7.8199		
	0.0550	7.6939		
2	0.0554	7.7499	7.9693	0.3559
	0.0556	7.7779		
	0.0599	8.3799		
3	0.0552	7.7219	7.9599	0.4002
	0.0602	8.4219		
	0.0553	7.7359		

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
5	0.0685	9.5839	9.1826	0.7815
	0.0592	8.2819		
	0.0692	9.6819		
8	0.0725	10.1439	10.6339	0.4766
	0.0793	11.0959		
	0.0762	10.6619		
12	0.0729	10.1999	10.9093	2.2014
	0.0654	9.1499		
	0.0956	13.3779		
24	0.0956	13.3779	13.9277	0.5053
	0.1002	14.0331		
	0.1027	14.3719		
36	0.1108	15.5059	15.2166	0.3554
	0.1095	15.3239		
	0.1059	14.8199		
48	0.1226	17.1649	15.9096	1.2636
	0.1046	14.6379		
	0.1138	15.9259		
60	0.1134	15.8699	16.0379	1.1711
	0.1235	17.2839		
	0.1069	14.9599		
72	0.1103	15.4359	16.0239	0.6691
	0.1135	15.8839		
	0.1197	16.7519		

Table F5 Chitosan releasing profile of 0/100 chitosan-sericin blend ratio in acetic/acetate buffer at pH 5.5

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
0.166	-0.0013	-0.1746	-0.3003	0.3366
	-0.0003	-0.0446		
	-0.0052	-0.6816		
0.333	-0.0003	-0.0481	-0.4541	0.3632
	-0.0053	-0.7481		
	-0.0040	-0.5661		
0.5	-0.0012	-0.1741	0.0177	0.3071
	0.0027	0.3719		
	-0.0010	-0.1447		
0.75	0.0008	0.1059	0.2413	0.1402
	0.0017	0.2319		
	0.0028	0.3859		
1	0.0016	0.2179	0.1993	0.0583
	0.0010	0.1339		
	0.0018	0.2459		
1.30	0.0019	0.2711	0.2777	0.0493
	0.0017	0.2319		
	0.0024	0.3299		
2	0.0019	0.2599	0.2693	0.1542
	0.0009	0.1199		
	0.0031	0.4279		
3	0.0024	0.3299	0.2506	0.0718
	0.0014	0.1899		
	0.0017	0.2319		

Time (hr)	Absorbance at 570 nm	Chitosan Concentration (mg)	Average	SD
5	0.0031	0.4279	0.3859	0.0420
	0.0025	0.3439		
	0.0028	0.3859		
8	0.0068	0.9459	0.8806	0.0796
	0.0065	0.9039		
	0.0057	0.7919		
12	0.0078	1.0859	1.1233	0.0771
	0.0077	1.0719		
	0.0087	1.2119		
24	0.0115	1.6039	1.4406	0.3200
	0.0077	1.0719		
	0.0118	1.6459		
36	0.0128	1.7859	1.6039	0.1590
	0.0110	1.5339		
	0.0107	1.4919		
48	0.0118	1.6459	1.6926	0.0808
	0.0118	1.6459		
	0.0128	1.7859		
60	0.0117	1.6319	1.6926	0.0932
	0.0118	1.6459		
	0.0129	1.7999		
72	0.0107	1.4919	1.7206	0.2040
	0.0128	1.7859		
	0.0135	1.8839		

CURRICULUM VITAE

Name: Mr. Thus La-ongnuan

Date of Birth: July 3, 1989

Nationality: Thai

University Education:

2007–2010 Bachelor Degree of Science in Chemistry, Faculty of Science,
Naresuan University, Phitsanulok, Thailand

Proceedings:

1. La-ongnuan, T.; and Rujiravanit, R. (2013, April 23rd) Development of Wound Dressing with Enhanced Wound Healing Properties: Preparation of Chitosan/Silk Serecin-incorporated Bacterial Cellulose. Proceedings of the 4th Research Symposium on Petrochemical and Materials Technology and The 19th PPC Symposium on Petroleum, Petrochemicals, and Polymers, Bangkok, Thailand.

Presentations:

1. La-ongnuan, T.; and Rujiravanit, R. (2013, April 23rd) Development of Wound Dressing with Enhanced Wound Healing Properties: Preparation of Chitosan/Silk Serecin-incorporated Bacterial Cellulose. Paper presented at the 4th Research Symposium on Petroleum, Petrochemicals, and Advanced Materials and The 19th PPC Symposium on Petroleum, Petrochemicals, and Polymers, Bangkok, Thailand.