

REFERENCES

- Ajavakoma, A., Supsvetson, S., Somboot, A. and Sukwattanasinitta, M. (2012). Products from microwave and ultrasonic wave assisted acid hydrolysis of chitin. Carbohydrate Polymers 90(1), 73-77.
- Allan, G.G. and Peyron, M. (1995). Molecular weight manipulation of chitosan I: kinetics of depolymerization by nitrous acid. Carbohydr Res 277(2), 257-272.
- Benhabiles, M.S., Salah, R. and Lounici, H. (2012). Antibacterial activity of chitin, chitosan, and its oligomers prepared from shrimp shell waste. Food Hydrocolloids 29(1), 48-56.
- Cabrera, J.C. and Cutsem, P.V. (2005). Preparation of chitooligosaccharides with degree of polymerization higher than 6 by acid or enzymatic degradation of chitosan Biochemical Engineering Journal 25(2), 165-172.
- Chang, K.L.B., Tsai, G., Lee, J. and Fu, W.-R. (1997). Heterogeneous N-deacetylation of chitin in alkaline solution. Carbohydr Res 303(3), 327-332.
- Cohen, E. (2010). Chitin Biochemistry. 38, 5-74.
- Einbu, A., Grasdalen, H. and Varum, K.M. (2007). Kinetics of hydrolysis of chitin/chitosan oligomers in concentrated hydrochloric acid. Carbohydr Res 342(8), 1055-1062.
- Fernandes, J.C., Eaton, P., Nascimento, H., Gião, M.S., Ramos, O.S., Belo, L., Santos-Silva, A., Pintado, M.E. and Malcata, F.X. (2010). Antioxidant activity of chitooligosaccharides upon two biological systems: Erythrocytes and bacteriophages. Carbohydrate Polymers 79(4), 1101-1106.
- Hasegawa, M., Isogai, A. and Onabe, F. (1993). Preparation of low-molecular-weight chitosan using phosphoric acid. Carbohydrate Polymers 20(4), 279-283.

- He, G., Wang, Z., Zheng, H., Yin, Y., Xiong, X. and Lin, R. (2012). Preparation, characterization and properties of aminoethyl chitin hydrogels. Carbohydr Polym 90(4), 1614-1619.
- Huang, K.-S., Wu, W.-J., Chen, J.-B. and Lian, H.-S. (2008). Application of low-molecular-weight chitosan in durable press finishing. Carbohydrate Polymers 73(2), 254-260.
- Huang, L., Zhai, M., Peng, J., Li, J. and Wei, G. (2007). Radiation-induced degradation of carboxymethylated chitosan in aqueous solution. Carbohydrate Polymers 67(3), 305-312.
- Huang, R., Mendis, E., Rajapakse, N. and Kim, S.-K. (2006). Strong electronic charge as an important factor for anticancer activity of chitooligosaccharides (COS). Life Sciences 78(20), 2399-2408.
- Il'ina, A.V. and Varlamov, V.P. (2004). Hydrolysis of Chitosan in Lactic Acid. Applied Biochemistry and Microbiology 40 (3), 300-303.
- Jung, J. and Zhao, Y. (2011). Characteristics of deacetylation and depolymerization of beta-chitin from jumbo squid (*Dosidicus gigas*) pens. Carbohydr Res 346(13), 1876-1884.
- Lieberman, A.M.. and Lichtenberg, J.A. (2005). Principles of plasma discharges and materials processing. Canada, John Wiley and Sons.
- Lia, J., Dua, Y. and Yanga, J. (2005). Preparation and characterisation of low molecular weight chitosan and chito-oligomers by a commercial enzyme. Polymer Degradation and Stability 87(3), 441-448.
- Mohammed, M.H., Williams, P.A. and Tverezovskaya, O. (2013). Extraction of chitin from prawn shells and conversion to low molecular mass chitosan. Food Hydrocolloids 31(2), 166-171.
- Mohan, C.O., Ravishankar, C.N., Lalitha, K.V. and Srinivasa Gopal, T.K. (2012). Effect of chitosan edible coating on the quality of double filleted Indian oil sardine (*Sardinella longiceps*) during chilled storage. Food Hydrocolloids 26(1), 167-174.
- Nagahama, H., Rani, V.V., Shalumon, K.T., Jayakumar, R., Nair, S.V.. Koiwa, S.,

- Furuike, T. and Tamura, H. (2009). Preparation, characterization, bioactive and cell attachment studies of alpha-chitin/gelatin composite membranes. *Int J Biol Macromol* 44(4), 333-337.
- Peter, M., Sudheesh Kumar, P.T., Binulal, N.S., Nair, S.V., Tamura, H. and Jayakumar, R. (2009). Development of novel α -chitin/nanobioactive glass ceramic composite scaffolds for tissue engineering applications. *Carbohydrate Polymers* 78(4), 926-931.
- Potocký, Š., Saito, N. and Takai, O. (2009). Needle electrode erosion in water plasma discharge. *Thin Solid Films* 518(3), 918-923.
- Prasertsung, I., Damrongsakkul, S., Terashima, C., Saito, N. and Takai, O. (2012). Preparation of low molecular weight chitosan using solution plasma system. *Carbohydrate Polymers* 87(4), 2745-2749.
- Prashanth, K.V.H. and Tharanathan, R.N. (2005). Depolymerized products of chitosan as potent inhibitors of tumor-induced angiogenesis *Biochim Biophys Acta* 1722 (1), 22– 29.
- Quan, H., Zhu, F. and Han, X. (2009). Mechanism of anti-angiogenic activities of chitooligosaccharides may be through inhibiting heparanase activity *Medical Hypotheses* 73(2), 205-206.
- Saito, G., Hosokai, S. and Akiyama, T. (2011). Synthesis of ZnO nanoflowers by solution plasma. *Materials Chemistry and Physics* 130(1-2), 79-83.
- Sjoholm, K.H., Cooney, M. and Minteer, S.D. (2009). Effects of degree of deacetylation on enzyme immobilization in hydrophobically modified chitosan. *Carbohydrate Polymers* 77(2), 420-424.
- Sugiarto, A.T. and Sato, M. (2001). Pulsed plasma processing of organic compounds in aqueous solution. *Thin Solid Films* 386(2), 295-299.
- Takai, O. (2008). Solution plasma. *Pure Apply Chem* 80(9), 2003-2011
- Tamura, H., Furuike, T. and Nair, S.V. (2011). Biomedical applications of chitin hydrogel membranes and scaffolds. *Carbohydrate Polymers* 84(2), 820-824.
- Tamura, H., Nagahama, H. and Tokura, S. (2006). Preparation of chitin hydrogel under mild conditions. *Cellulose* 13(2006), 357-364.
- Tolaimate, A., Desbrieres, J. and Rhazi, M. (2000). On the influence of

- deacetylation process on the physicochemical characteristics of chitosan from squid chitin. Polymer 41(7), 2463-2469.
- Tolaimate, A., Desbrieres, J., Rhazi, M. and Alagui, A. (2003). Contribution to the preparation of chitins and chitosans with controlled physico-chemical properties. Polymer 44(26), 7939-7952.
- Tong, D.G., Luo, Y.Y., Chu, W., Guo, Y.C. and Tian, W. (2010). Cutting Of Carbon Nanotubes Via Solution Plasma Processing. Plasma Chemistry and Plasma Processing 30(6), 897-905.
- Varum, K.M. and Ottoy O.Smidrod, M.H. (2001). Carbohydrate Polymers. Acid hydrolysis of chitosans 46(1), 89-98.
- Vishu Kumar, A. (2004). Low molecular weight chitosans: preparation with the aid of papain and characterization. Biochimica et Biophysica Acta (BBA) - General Subjects 1670(2), 137-146.
- Wasikiewicz, J.M., Yoshii, F., Nagasawa, N., Wach, R.A. and Mitomo, H. (2005). Degradation of chitosan and sodium alginate by gamma radiation, sonochemical and ultraviolet methods. Radiation Physics and Chemistry 73(5), 287-295.
- Watthanaphanit, A. and Saito, N. (2013). Effect of polymer concentration on the depolymerization of sodium alginate by the solution plasma process. Polymer Degradation and Stability 98(5), 1072-1080.
- Wu, P., Sun, J.H., Huang, Y.Y., Gu, G.F. and Tong, D.G. (2012). Solution plasma synthesized nickel oxide nanoflowers: An effective NO₂ sensor. Materials Letters 82(0), 191-194.
- Zhu, X., Cai, J., Yang, J. and Su, Q. (2005). Determination of glucosamine in impure chitin samples by high-performance liquid chromatography. Carbohydr Res 340(10), 1732-1738.

APPENDICES

Appendix A Characterization of raw material

Table A1 Percent yield of chitin

Samples	Dry weight (%)
Dry shrimp shell	100.00
Decalification product	30.48
Deproteinization product	22.87
Chitin	22.87

Table A2 Percent yield of chitin hydrogel

Number of samples	Chitin solution (g)	Chitin hydrogel (g)
1	100	50.85
2	100	52.12
3	100	54.6
Average	100	52.53
SD	0	1.907258

Table A3 Percent of chitin and water contents in chitin hydrogel

Number of samples	Wet weight (g)	Dry weight (g)	Chitin (%)	Water (%)
1	1.7320	0.06	3.464203	96.5358
2	1.7179	0.0588	3.422784	96.57722
3	1.7008	0.0495	2.910395	97.0896
Average	1.7169	0.0561	3.267517	96.73248
SD	0.015624	0.0057	0.30848	0.30848

Table A3 Degree of deacetylation of chitin powder and chitin hydrogel

Number of samples	Degree of deacetylation (%DD)				SD
	1	2	3	Average	
Type of chitin					
Chitin powder	35.26	36.61	34.3	35.39	1.160474
Chitin hydrogel	38.19	37.98	35.91	37.36	1.260119

Appendix B Deacetylation of chitin hydrogel with KOH/MeOH

Table B1 Effect of amount of chitin hydrogel in KOH solution on the percent of degree of deacetylation

Amount of chitin hydrogel (g)	Degree of deacetylation (%DD)				SD
	1	2	3	Average	
10	60.76	62.33	61.14	61.41	0.819085
20	57.58	56.68	58.88	57.71333	1.106044
30	52.89	53.64	55.24	53.92333	1.200347

Table B2 Effect of amount of chitin hydrogel in KOH solution on the percent of solubility

Amount of chitin hydrogel (g)	% Solubility				SD
	1	2	3	Average	
10	30.26	31.51	28.44	30.07	1.543794
20	22.42	24.48	23.64	23.51333	1.035825
30	16.56	15.84	18.26	16.88667	1.242632

Table B1 Effect of 0% w/v KOH concentration in methanol solvent on degree of deacetylation of chitin hydrogel by using solution plasma technique

0% (w/v) KOH concentration					
Reaction time (hour)	Degree of deacetylation (%DD)				
	1	2	3	Average	SD
0	38.19	37.98	35.91	37.36	1.260119
1	40.15	44.71	43.26	42.71	2.329814
2	47.22	46.88	48.92	47.67	1.092947
3	48.12	49.82	50.12	49.35	1.078579
4	52.88	51.78	53.84	52.83	1.030793
5	56.23	55.44	54.26	55.31	0.991413

Table B2 Effect of 7% w/v KOH concentration in methanol solvent on degree of deacetylation of chitin hydrogel by using solution plasma technique

7% (w/v) KOH concentration					
Reaction time (hour)	Degree of deacetylation (%DD)				
	1	2	3	Average	SD
0	38.19	37.98	35.91	37.36	1.260119
1	53.82	55.18	54.12	54.37	0.714516
2	62.74	61.9	60.11	61.58	1.343292
3	66.79	63.95	63.28	64.67	1.863447
4	67.95	66.91	65.64	66.83	1.156907
5	70.12	68.55	67.82	68.83	1.175287

Table B3 Effect of 10% w/v KOH concentration in methanol solvent on degree of deacetylation of chitin hydrogel by using solution plasma technique

10% (w/v) KOH concentration					
Reaction time (hour)	Degree of deacetylation (%DD)				
	1	2	3	Average	SD
0	38.19	37.98	35.91	37.36	1.260119
1	60.76	62.33	61.14	61.41	2.227308
2	66.61	68.51	68.25	67.79	1.030146
3	68.54	70.61	69.26	69.47	1.050857
4	72.25	71.78	73.25	72.26	0.750755
5	72.26	70.67	75.39	74.44	2.401506

Table B4 Effect of 0% w/v KOH concentration in methanol solvent on degree of deacetylation of chitin hydrogel by using conventional heat treatment

0% (w/v) KOH concentration					
Reaction time (hour)	Degree of deacetylation (%DD)				
	1	2	3	Average	SD
0	38.19	37.98	35.91	37.36	1.260119
1	38.12	37.92	38.98	38.34	0.563205
2	39.21	39.4	40.62	39.74333	0.765136
3	40.88	40.56	41.4	40.94667	0.42395
4	42.11	42.84	43.26	42.73667	0.581922
5	43.21	44.56	45.71	44.49333	1.251333

Table B5 Effect of 7% w/v KOH concentration in methanol solvent on degree of deacetylation of chitin hydrogel by using conventional heat treatment

7% (w/v) KOH concentration					
Reaction time (hour)	Degree of deacetylation (%DD)				
	1	2	3	Average	SD
0	38.19	37.98	35.91	37.36	1.260119
1	40.13	39.22	38.14	39.16333	0.996209
2	43.88	41.71	45.39	43.66	1.849838
3	47.88	48.71	50.39	48.99333	1.278762
4	50.23	52.88	53.65	52.25333	1.794055
5	55.42	56.64	55.53	55.86333	0.674858

Table B6 Effect of 10% w/v KOH concentration in methanol solvent on degree of deacetylation of chitin hydrogel by using conventional heat treatment

10% (w/v) KOH concentration					
Reaction time (hour)	Degree of deacetylation (%DD)				
	1	2	3	Average	SD
0	38.19	37.98	35.91	37.36	1.260119
1	42.13	41.22	40.14	41.16333	0.996209
2	47.11	46.25	45.66	46.34	0.729178
3	52.88	51.71	50.39	51.66	1.245753
4	52.23	54.88	55.65	54.25	1.794055
5	57.42	58.64	53.53	56.53	2.668726

Appendix C Solubility of chitin hydrogel by plasma treated

Table C1 Effect of 7% w/v KOH concentration in methanol solvent on solubility of plasma treated chitin hydrogel

7% (w/v) KOH concentration					
Reaction time (hour)	7% solubility of plasma treated chitin hydrogel				
	1	2	3	Average	SD
0	0	0	0	0	0
1	19.74	20.26	22.64	20.88	1.546221
2	24.29	24.86	26.52	25.23	1.158548
3	41.24	42.18	44.56	42.66	1.711257
4	46.6	51.54	52.88	50.24	3.307507
5	55.5	58.5	59.46	57.82	2.06572

Table C2 Effect of 10% w/v KOH concentration in methanol solvent on solubility of plasma treated chitin hydrogel

10% (w/v) KOH concentration					
Reaction time (hour)	% solubility of plasma treated chitin hydrogel				
	1	2	3	Average	SD
0	0	0	0	0	0
1	42.1	38.3	39.81	40.07	1.913296
2	47.83	53.22	50.36	50.47	2.696683
3	52.34	61.5	62.73	58.85667	5.677009
4	88.33	85.08	90.62	88.01	2.783828
5	90.94	89	92.49	90.81	1.748628

Appendix D Yield of plasma treated chitin hydrogel

Table D1 Yield of plasma treated chitin hydrogel with 0% w/v KOH concentration

0% (w/v) KOH concentration					
Reaction time (hour)	% Yield of plasma treated chitin hydrogel				
	1	2	3	Average	SD
0	100	100	100	100	0
1	80.53	81.36	85.71	82.53	2.782199
2	75.55	76.43	73.22	75.06	1.658684
3	70.56	71.61	69.31	70.49	1.151448
4	66.12	65.71	67.78	66.54	1.096099
5	60.34	61.22	59.23	60.26	0.997213

Table D2 Yield of plasma treated chitin hydrogel with 7% w/v KOH concentration

7% (w/v) KOH concentration					
Reaction time (hour)	% Yield of plasma treated chitin hydrogel				
	1	2	3	Average	SD
0	100	100	100	100	0
1	77.89	78.91	79.59	78.79667	0.855648
2	66.48	67.12	65.99	66.53	0.566657
3	44.19	42.26	45.21	43.88667	1.49821
4	35.13	32.22	36.76	34.70333	2.299877
5	30.12	25.71	28.25	28.02667	2.213466

Table D3 Yield of plasma treated chitin hydrogel with 10% w/v KOH concentration

10% (w/v) KOH concentration					
Reaction time (hour)	-% Yield of plasma treated chitin hydrogel				
	1	2	3	Average	SD
0	100	100	100	100	0
1	81.51	74.44	75.14	77.03	3.895549
2	61.43	62.95	60.19	61.52333	1.382365
3	46.085	43.63	40.12	43.27833	2.998009
4	37.5456	28.0485	30.51	32.0347	4.928718
5	30.3041	25.61	24.57	26.82803	3.054943

Table D4 Yield of conventional heat treatment chitin hydrogel with 10% w/v KOH concentration

10% (w/v) KOH concentration					
Reaction time (hour)	% Yield of plasma treated chitin hydrogel				
	1	2	3	Average	SD -
0	100	100	100	100	0
1	95.36	96.78	92.19	94.77667	2.349943
2	81.51	80.24	78.26	80.00333	1.637875
3	56.48	58.34	59.77	58.19667	1.649677
4	49.69	50.12	52.98	50.93	1.788323
5	45.17	44.56	46.23	45.32	0.845044

Appendix E Acid hydrolysis of chitin hydrogel by using solution plasma in acetic acid solution

Table E1 Degree of deacetylation of depolymerized chitin hydrogel

Number of samples	Degree of deacetylation (%DD)				SD
	1	2	3	Average	
Time (hours)					
3	45.68	43.12	45.89	44.89667	1.542217
5	47.89	48.56	48.26	48.23667	0.335609

Table E1 Effect of depolymerized chitin hydrogel at 3 hours in 1% v/v acetic acid solution and deacetylated in 10 % KOH/MeOH on the degree of deacetylation.

3 hours of depolymerization in 1%v/v acetic acid and deacetylation at 10% (w/v) KOH concentration					
Reaction time (hour)	Degree of deacetylation (%DD)				
	1	2	3	Average	SD
0	45.68	43.12	45.89	44.90	1.542217
1	65.53	67.53	66.24	66.43	1.01392
2	72.64	71.399	69.51	71.18	1.57614
3	74.22	73.78	71.84	73.28	1.266333
4	75.08	75.96	74.88	75.64	0.574572
5	74.63	76.67	77.18	76.16	1.349333

Table E2 Effect of depolymerized chitin hydrogel at 5 hours in 1% v/v acetic acid solution and deacetylated in 10 % KOH/MeOH on the degree of deacetylation.

5 hours of depolymerization in 1%v/v acetic acid and deacetylation at 10% (w/v) KOH concentration					
Reaction time (hour)	Degree of deacetylation (%DD)				
	1	2	3	Average	SD
0	47.89	48.56	48.26	48.23667	0.335609
1	68.23	69.81	70.24	69.42667	1.058411
2	74.73	72.66	73.26	73.88333	1.561292
3	73.44	75.62	73.12	74.06	1.360441
4	76.12	76.48	76.66	76.42	0.274955
5	80.28	76.36	78.88	78.50667	1.986488

Table E3 Effect of depolymerized chitin hydrogel at 3 hours in 1% v/v acetic acid solution and deacetylated in 10 % KOH/MeOH on the solubility.

3 hours of depolymerization in 1%v/v acetic acid and deacetylation at 10% (w/v) KOH concentration					
Reaction time (hour)	% Solubility				
	1	- 2	3	Average	SD
1	59.16	60.77	63.38	61.10333	2.129656
2	69.82	73.06	76.5	73.12667	3.340499
3	79.73	79.62	82.96	80.77	1.897393
4	88.91	90.86	92.35	90.70667	1.725118
5	91.04	94.3	96.3	93.88	2.655033

Table E4 Effect of depolymerized chitin hydrogel at 5 hours in 1% v/v acetic acid solution and deacetylated in 10 % KOH/MeOH on the solubility.

5 hours of depolymerization in 1%v/v acetic acid and deacetylation at 10% (w/v) KOH concentration					
Reaction time (hour)	% Solubility				
	1	2	3	Average	SD
1	67.42	68.2	69.28	67.81	0.934024
2	76.06	76.5	80.4	76.28	1.084866
3	88.47	74.88	75.56	81.675	1.192644
4	93.55	95.03	95.36	94.29	0.963967
5	97.32	98.64	92.49	97.98	1.034601


Appendix FDetermination of Molecular weight by GPC**Figure F1 Calibration curve for GPC measurement using pullulan standard**

Table F2The molecular weight of deacetylation chitin hydrogel in 10% w/v KOH/MeOH with different times by using solution plasma

Conditions	Treated chitin hydrogel 3 hours	Treated chitin hydrogel 4 hours	Treated chitin hydrogel 5 hours
M_w	340,124	324,709	321,624
M_n	116,703	124,551	128,701
PDI	2.91	2.61	2.50
M_z	738,640	633,138	610,783

Table F3The molecular weight of depolymerization chitin hydrogel in 1%v/v acetic solution and deacetylation in 10% w/v KOH/MeOH with different times by using solution plasma

Conditions	Treated chitin hydrogel 3 hours	Treated chitin hydrogel 4 hours	Treated chitin hydrogel 5 hours
M_w	281,454	272,856	251,567
M_n	117,292	128,705	123,631
PDI	2.40	2.12	2.03
M_z	636,744	620,890	615,871

CURRICULUM VITAE

Name : Ms. Kuntinee Somboonying

Date of Birth : March 12, 1988

Nationality : Thai

University Education :

2007–2010 Bachelor Degree of Science in Chemistry, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

Proceedings

1. Somboonying, K.; and Rujiravanit, R. (2014, April 22) Solution Plasma Process for Deacetylation and Depolymerization of Chitin Hydrogel . Proceedings of the 5th Research Symposium on Petrochemical and Materials Technology and the 20th PPC Symposium on Petroleum, Petrochemicals, and Polymers. Bangkok, Thailand.

Presentation

1. Somboonying, K.; and Rujiravanit, R. (2014, April 22) Solution Plasma Process for Deacetylation and Depolymerization of Chitin Hydrogel . Poster presented at The 5th Research Symposium on Petrochemical and Materials Technology and the 20th PPC Symposium on Petroleum, Petrochemicals, and Polymers. Bangkok, Thailand.
2. Somboonying, K.; and Rjuravanit, R. (2014, May 18-23) Deacetylation and Depolymerization of Chitin Hydrogel via Solution Plasma Process. Poster presented at 5th International Conference on Plasma Medicine (ICPM) 2014, Nara, Japan.