

## REFERENCES

- Abram SE, Kostreva DR. Spinal cord metabolic response to noxious radiant heat stimulation of the cat hind footpad. **Brain Res** 1986;385:143-7.
- Adham H, Romanienko P, Hartig P, Weinshank RL, Branchek T. The rat 5-hydroxytryptamine<sub>1B</sub> receptor is the species homologue of the human 5-hydroxytryptamine<sub>1Dβ</sub> receptor. **Mol Pharmacol** 1992;41:1-7.
- Aghajanian GK, Sprouse JS, Sheldon P, Rasmussen K. Electrophysiology of brain serotonin receptors: subtype specificity for effector mechanisms. In: Paoletti R, Vanhoutte PM, Brunello N, Maggi FM, editors. **Serotonin: From cell biology to pharmacology and therapeutics**. Dordrecht: Kluwer, 1990:277-82.
- Albert PR, Zhou QY, Van Tol HHM, Bunzow JR, Civelli O. Cloning, functional expression and messenger RNA tissue distribution of the rat 5-hydroxytryptamine<sub>1A</sub> receptor gene. **J Biol Chem** 1990;265:5825-32.
- Alhaider AA. Antinociceptive effect of ketanserin in mice: involvement of supraspinal 5-HT<sub>2</sub> receptors in nociceptive transmission. **Brain Res** 1991;543:335-40.
- Alhaider AA, Wilcox GL. Spinal 5-HT<sub>3</sub> receptor-mediated antinociception : possible release of GABA. **J Neurosci** 1991;18:81-8.
- Alhaider AA, Wilcox GL. Differential roles of 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptor subtypes in modulating spinal nociceptive transmission in mice. **J Pharmacol Exp Ther** 1993;265:378-85.
- Ali Z, Wu G, Kozlov A, Barasi S. The activation of 5-HT<sub>1</sub> agonists and antagonists on nociceptive processing in the rat spinal cord: results from behavioral and electrophysiological studies. **Brain Res** 1994;661:83-90.
- Archer T, Arwestrom E, Minor BG, Persson ML, Post C, Sundstrom E, et al. (+)-8-OH-DPAT and 5-MeODMT induced analgesia is antagonized by noradrenaline depletion. **Physiol Behav** 1987;39:95-102.

- Armstrong RC, Montining MR. Transsynaptic control of gene expression. *Ann Rev Neurosci* 1993;16:17-29.
- Bach AW, Unger L, Sprengel R, Mengod G, Palacios J, Seeburg PH. Structure, functional expression and spatial distribution of a cloned cDNA encoding a rat 5-HT<sub>1D</sub>-like receptor. *J Recept Res* 1993;13:479-502.
- Baez M, Yu L, Cohen ML. Pharmacological and molecular evidence that the contractive response to serotonin in rat stomach fundus is not mediated by activation of the 5-hydroxytryptamine(1C) receptor. *Mol Pharmacol* 1990;38:31-7.
- Banks WA, Kastin AJ, Trentman TL, Haynes HS, Johnson BG, Galina ZH. Mediation of serotonin-induced analgesia by the 5-HT<sub>2</sub> receptor in the pentobarbital anesthetized mouse model. *Brain Res Bull* 1988;21:887-91.
- Barber A, Harting J, Wolf HP. Antinociceptive effects of the 5-HT<sub>2</sub> antagonist ritanserin in rats: evidence for an activation of descending monoaminergic pathways in the spinal cord. *Neurosci Lett* 1989;99:234-8.
- Basbaum AI, Fields HL. Endogenous pain control system: Brainstem spinal pathways and endorphin circuitry. *Ann Rev Neurosci* 1984;7:309-38.
- Baskin DS, Mehler WR, Hosobuchi Y, Richardson DE. Autopsy analysis of the safety, efficacy, and cartography of electrical stimulation of the central gray in humans. *Brain Res* 1986;371:231-6.
- Beer MS, Middlemiss DN. Serotonin-5-O-carboxymethyl-glycyl [<sup>125</sup>I] tyrosinamide labels the 5-HT<sub>1Dβ</sub> receptor subtype in human cortex. *Eur Pharmacol* 1993;242:195-8.
- Beer MS, Stanton JA, Bevan Y, Chauhan NS, Middlemiss DN. An investigation of the 5-HT<sub>1D</sub> receptor binding affinity of 5-hydroxytryptamine, 5-carboxamidotryptamine and sumatriptan in the central nervous system of seven species. *Eur J Pharmacol* 1992;213:193-7.

- Clineschmidt BV, Reiss DR, Pettibone DJ, Robinson JL. Characterization of 5-hydroxytryptamine receptors in rat stomach fundus. *J Pharmacol Exp Ther* 1985;235:696-708.
- Coderre TJ, Melzack R. Increased pain sensitivity following heat injury involves a central mechanism. *Brain Res* 1985;15:259-62.
- Cohen ML, Wittenauer LA. Serotonin receptor activation of phosphoinositide turnover in uterine, fundal, vascular and tracheal smooth muscle. *J Cardiovasc Pharmacol* 1987;10:176-81.
- Coleman RA, Humphey PPA. Prostaglandins-receptors and mode of action. In: Vane JR, Grady JO, editors. *Clinical Application of Prostaglandin*. Seven Oaks : Edward Arnold Publisher, 1993:15-36.
- Conn PJ, Sanders-Bush E. Selective 5-HT<sub>2</sub> antagonists inhibit serotonin stimulated phosphatidylinositol metabolism in cerebral cortex. *Neuropharmacol* 1984;23:993-6.
- Con PJ, Sanders-Bush E. Serotonin-stimulated phosphoinositide turnover : mediation by the S2 binding site in rat cerebral cortex but not in subcortical regions. *J Pharmacol Exp Ther* 1985;234:195-203.
- Conn PJ, Sanders-Bush E, Hoffman BJ, Hartig PR. A unique serotonin receptor in choroid plexus is linked to phosphotidylinositol turnover. *Proc Natl Acad Sci* 1986;4086-8.
- Crisp T, Stafinsky JL, Boja JW, Schechter MD. The antinociceptive effects of 3,4-methylenedioxymethamphetamine (MDMA) in the rat. *Pharmacol Biochem Behav* 1989;34:497-501.
- Crisp T, Stafinsky JL, Spanos LS. Analgesic effects of serotonin and receptor-selective serotonin agonists in the rat spinal cord. *Gen Pharmacol* 1991;22:247-51.
- Cross AJ, Crow TJ, Gerrier IN, Johnson JA, Bloom SR, Corsellis JAN. Serotonin receptor changes in dementia of the Alzheimer type. *J Neurochem* 1984;43:1574-81.
- Cross AJ, Slater P, Perry EK, Perry RM. An autoradiographic analysis of

- serotonin receptors in human temporal cortex changes in Alzheimer-type dementia. *Neurochem Int* 1988;13:89-96.
- Curzon G, Kennett GA. MCPP: a tool for studying behavioral responses associated with 5-HT<sub>(1C)</sub> receptors. *Trend Pharmacol Sci* 1991;11:181-2.
- D'Amour FE, Smith D. A method for determining loss of pain sensation. *J Pharmacol Exp Ther* 1941;72:74-9.
- Daval G, Verge D, Basbaum AI, Bourgoin S, Hamon M. Autoradiographic evidence of serotonin binding sites on primary afferent fibers in the dorsal horn of the rats spinal cord. *Neurosci* 1987;83:71-6.
- Davies MF, Deisz RA, Prince DA, Peroutka SJ. Two distinct effects of 5-hydroxytryptamine on single cortical neurons. *Brain Res* 1987;423:347-52.
- Davies JE, Roberts MH. 5-Hydroxytryptamine reduces substance P responses on dorsal horn interneurons : A possible interaction of neurotransmitter. *Brain Res* 1981;399:404.
- De Chaffoy DE, Courcelles D, Leysen JE, De Clerck F, Van Belle H, Janssen PAJ. Phospholipid turnover is the signal transducting system coupled to serotonin-S2 receptors. *J Biol Chem* 1985;260:7603-8.
- De Montigny C, Blier P. Electrophysiological evidence for the distinct properties of presynaptic and postsynaptic 5-HT<sub>1A</sub> receptors: possible clinical relevance. In: Langer SZ, Brunello N, Racagni G, Mendlewicz G, editors. *Serotonin Receptor Subtypes: Pharmacological Significance and Clinical Implications*. J Int Acad Biomed Drug Res: Karger, 1992:80-8.
- De vivo M, Maayani S. Inhibition of Forskolin-stimulated adenylate cyclase activity by 5-HT receptor agonists. *Eur J Pharmacol* 1985;119:231-4.
- De vivo M, Maayani S. Characterization of 5-hydroxytryptamine<sub>1A</sub>-receptor-mediated inhibition of forskolin-stimulated adenylate cyclase activity in

- guinea-pig and rat hippocampal membranes. **J Pharmacol Exp Ther** 1986;238:248-52.
- Dhasmana KM, Zhu YN, Cruz SL, Villalon CM. Gastrointestinal effects of 5-hydroxytryptamine and related drugs. **Life Sciences** 1993;53:1651-61.
- Dickenson AH, Sullivan AF. Peripheral origins and central modulation of subcutaneous formalin-induced activity of rat dorsal horn neurons. **Neurosci Lett** 1987a;83:207-11.
- Dickenson AH, Sullivan AF. Subcutaneous formalin-induced activity of dorsal horn neurons in the rat : Differential response to an intrathecal opiate administered pre or post formalin. **Pain** 1987b;30:349-60.
- Doods HN, Boddeke HW, Kalkman HO, Hoyer D, Mathy MJ, van Zweiten PA. Central 5-HT<sub>1A</sub> receptors and the mechanism of the central hypotensive effect of (+)8-OH-DPAT, DP-5-CT, R28935 and urapidil. **J Cardiovasc Pharmacol** 1988;11:432-7.
- Douglas WW. Histamine and 5-hydroxytryptamine(serotonin) and their antagonists. In: Gilman AG, Goodman LS, Gilman A, editors. **Goodman and Gilman's The pharmacological basis of therapeutics**. New York: Macmillan, 1980: 609-46.
- Doyle VM, Creba JA, Ruegg UT, Hoyer D. Serotonin increases the production of inositol phosphates and mobilizes calcium via the 5-HT<sub>2</sub> receptor in A7r5 smooth muscle cells. **Naunyn Schmiedebergs Arch Pharmacol** 1986;333:98-103.
- Dreteler GH, Wouters W, Saxena PR. Comparison of the cardiovascular effects of the 5-HT<sub>(1A)</sub> receptor agonist flesinoxan with that of 8-OH-DPAT in the rat. **Eur J Pharmacol** 1990;180:339-49.
- Dreteler GH, Wouters W, Toorop GP, Jansen JAP, Saxena PR. Systemic and regional hemodynamic effects of the 8-hydroxy-2(di-N-propyl amino)tretalin in the conscious rat. **J Cardiovasc Pharmacol** 1991;17:488-93.
- Dubner R. Pain research in animals. **Ann NY Acad Sci** 1983;406:128-32.
- Dubner R. Methods of assessing pain in animals. In: Wall PD, Melzack R,

- editors. **Text book of pain.** New York:Churchill Livingstone, 1994;295-302.
- Dubner R, Beitel RE, Brown FJ. A behavioral animal model for the study of pain mechanisms in primates. In : Weisenberg M, Tursky B, editors. **Pain : new perspectives in therapy and research.** New york : Plenum ;1976:155-70.
- Dubner R, Sumino R, Wood W. Peripheral 'cold' fiber population responsive to innocuous and noxious thermal stimuli applied to monkey's face. **J Neurophysiol** 1975;38:1373-1389.
- Dubuisson D, Dennis SG. The formalin test : a quantitative study of the analgesic effects of morphine, meperidine, and brain stem stimulation in rats and cats. **Pain** 1977;4:161-74.
- Eide PK. Stimulation of 5-HT<sub>1</sub> receptors in the spinal cord changes substance P-induced behavior. **Neuropharmacology** 1992;31:541-5.
- Eide PK, Berge OG, Tjølsen A, Hole K. Apparent hyperalgesia in the mouse tail-flick test due to increased tail skin temperture after lesioning of serotonergic pathway. **Acta Physio Scand** 1988;34:413-20.
- Eide PK, Hole K. Subsensitivity of serotonin and substance P receptors involved in nociception after repeated administration of a serotonin receptor agonists. **J Neural Transm** 1989;77:1-10.
- Eide PK, Hole K. Different role of 5-HT<sub>1A</sub> and 5-HT<sub>2</sub> receptors in spinal cord in the control of nociceptive responsiveness. **Neuropharmacology** 1991a;30:727-31.
- Eide PK, Hole K. Interactions between serotonin and substance P in the spinal regulation of nociception. **Brain Res** 1991b;550:225-30.
- Eide PK, Hole K, Berge O-G, Broch OJ. 5-HT depletion with 5,7-DHT, PCA and PCPA in mice: differential effects on the sensitivity to 5-MeODMT, 8-OH-DPAT and 5-HTP as measured by two nociceptive tests. **Brain Res** 1988;440:42-52.
- Eide PK, Joly NM, Hole K. The role of spinal cord 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub>

- receptors in the modulation of a spinal nociceptive reflex. **Brain res** 1990;36:195-200.
- Eide PK, Tjølsen A. Effects of serotonin receptor antagonists and agonists on the tail-flick response in mice involve altered tail-skin temperature. **Neuropharmacology** 1988;27:889-93.
- El-Yassir N, Fleetwood-Walker SM, Mitchell R. Heterogeneous effects of serotonin in the dorsal horn of rat: the involvement of 5-HT<sub>1</sub> receptor subtypes. **Brain Res** 1988;456:147-58.
- El-Yassir N, Fleetwood-Walker SM. A 5-HT<sub>1</sub>-type receptor mediates the antinociceptive effect of nucleus raphe magnus stimulation in the rat. **Brain Res** 1990;523:92-9.
- Engel G, Gothert M, Hoyer D, Schlicker E, Hillenbrand K. Identity of inhibitory presynaptic 5-hydroxytryptamine (5-HT) autoreceptors in the rat brain cortex with 5-HT<sub>1B</sub> binding sites. **Naunyn Schmiedebergs Arch Pharmacol** 1986;120:51-4.
- Fanelli RJ, Mc Monagle-Strucko K. Alteration of 5-HT<sub>1A</sub> receptor binding sites following chronic treatment with ipsapirone measured by quantitative autoradiography. **Synapse** 1992;12:75-80.
- Fargin A, Raymond JR, Lohse MJ, Kobilka BK, Caron MG, Lefkowitz RT. The genomic clone G-21 which resembles a β-adrenergic receptor sequence encodes the 5-HT<sub>1A</sub> receptor. **Nature** 1988;335:358-60.
- Fasmer OB, Berge O-G, Post C, Hole K. Effects of the putative 5-HT<sub>1A</sub> receptor agonist 8-OH-2-(di-n-propyl-amino)tetralin on nociceptive sensitivity in mice. **Pharmacol Biochem Behav** 1986;25:883-8.
- Fields HL, Basbaum AI. Central Nervous system Mechanisms of Pain Modulation In: Wall PD, Melzack R, editors. **Textbook of pain**. New York: Churchill livingstone, 1994:243.
- Fields H, Bry J, Hentall I, Zorman G. The activity of neurons in the rostral medulla of the rat during withdrawal from noxious heat. **J Neurosci** 1983;3:545-52.

- Fields HL , Heinricher MM , Mason P. Neurotransmitters in nociceptive modulatory circuits. *Ann Rev Neurosci* 1991;14:219-45.
- Foguet M, Hoyer D, Pardo LA, Parekh A, Kluxen FW, Kalkman HO, et al., Cloning and functional characterization of the rat stomach fundus serotonin receptor. *EMBO J* 1992b;11:3481-7.
- Foguet M, Nguyen H, Le H, Lubbert H. Structure of the mouse 5-HT<sub>1C</sub>, 5-HT<sub>2</sub> and stomach fundus serotonin receptor genes. *Neurorept* 1992a;3:345-8.
- Fone KCF, Robinson AJ, Marsden CA. Characterization of the 5-HT receptor subtypes involved in the motor behaviors produced by intrathecal administration of 5-HT agonist in rats. *Br J Pharmacol* 1991;103:1547-55.
- Fozard JR, Gray JA. 5-HT<sub>1C</sub> receptor activation: a key step in the initiation of migraine? *Trends Pharmacol Sci* 1989;10:307-9.
- Frazer A, Maayani S, Wolfe BB. Subtypes of receptors for serotonin. *Annu Rev Pharmacol Toxicol* 1990;30:307-48.
- Furchtgott RF, Vanhoutte PM. Endothelium-derived relaxing and contracting factors. *FASEB J* 1989;3-2007-18.
- Gershon MD. Serotonin, its role and receptors in enteric neurotransmission. In: Chwarcz R, editor. *Advances in Experimental Medicine and Biology*. New York : Plenum Press, 1991:221-30.
- Gibbon GH, Dzau VJ. The emerging concept of vascular remodeling. *N Eng J Med* 1994;330:1431-8.
- Gillis CN. Peripheral metabolism of serotonin. In: Vanhoutte PM, editer. *Serotonin and the Cardiovascular System*. New York: Raven Press, 1985:27-42.
- Gilman AG, Goodman LS, Gilman A, editors. *Goodman and Gilman's The pharmacological basis of therapeutics*. New York : Macmilian , 1980:609-46.

- Giordano J. Analgesic profile of centrally administered 2-methylserotonin against acute pain in rats. *Eur J Pharmacol* 1991;233-6.
- Giordano J, Dyche J. Differential analgesic actions of serotonin 5-HT<sub>3</sub> receptor antagonists in the mouse. *Neuropharmacol* 1989;28:423-7.
- Giordano J, Rogers LV. Peripherally administered serotonin 5-HT<sub>3</sub> receptor antagonists reduce inflammatory pain in rats. *Eur J Pharmacol* 1989;170:83-6.
- Gjerstad J, Tjølsen A, Hole K. The effects of 5-HT<sub>1A</sub> receptor stimulation on nociceptive dorsal horn neurons innervation. *Eur J Pharmacol* 1996;318:315-21.
- Goelet P, Castellucci VF, Schacher S, Kandel ER. The long and the short of long-term memory molecular framework. *Nature* 1986;322:419-22.
- Gogas KR, Presley RW, Levine JD, Basbaum AI. The antinociceptive action of a supraspinal opioids results from an increase in descending inhibitory control: correlation of nociceptive behavior and c-fos expression. *Neuroscience* 1991;42:617-28.
- Greenberg ME, Greene LA, Ziff EB. Nerve growth factor and epidermal growth factor induce rapid transient changes in proto-oncogene transcription in PC12 cells. *J biol Chem* 1985;14:101-14.
- Griersmith BT, Duggan AW. Prolonged depression of spinal transmission of nociceptive information by 5-HT administered in the substantia gelatinosa : Antagonism by methysergide. *Bain Res* 1980;187:231-6.
- Grunberg SM, Hesketh PJ. Control of Chemotherapy-induced emesis. *N Eng J Med* 1993;329:1790-96.
- Grubb BD, McQueen DS, Iggo A, Birrell GJ, Dutia MB. A study of 5-HT-receptors associated with afferent nerves located in normal and inflamed rat ankle joints. *Agents Actions* 1988;25:216-18.
- Grumbach L. The prediction of analgesic activity in man by animal testing. In: Knighton RS, Dumke PR, editors. *Pain*. New York: little brown, 1966;163-82.

- Belcher G, Ryall RW, Schaffner R. The differential effects of 5-hydroxytryptamine, noradrenaline and raphe stimulation on nociceptive and non-nociceptive dorsal horn interneurons in the cat. *Brain Res* 1978;151:307-21.
- Berge O-G. Effects of 5-HT receptor agonists and antagonists on a reflex response to radiant heat in normal and spinally transected rats. *Pain* 1982;13:253-66.
- Berge O-G, Hole K, Dahle H. Nociception is enhanced after low doses and reduced after high doses of the serotonin receptor agonist, 5-methoxy-N,N-dimethyltryptamine. *Neurosci Lett* 1980;19:219-23.
- Berridge M. Second messenger dualism in neuromodulation and memory. *Nature* 1986;323:294-5.
- Bervoets K, Rivet J-M, Millan MJ. 5-HT<sub>1A</sub> Receptors and the tail-flick response. IV. Spinally-localized 5-HT<sub>1A</sub> receptors postsynaptic to serotonergic neurons mediate spontaneous tail-flick in the rat. *J Pharmacol Exp Ther* 1987;67:67-186.
- Besson JM, Chaouch A. Peripheral and spinal mechanisms of nociception. *Physiol Rev* 1987;67:178-186.
- Bill AM, Knight M, Forster EA, Fletcher A. Direct evidence for an important species difference in the mechanism of 8-OH-DPAT-induced hypothermia. *Br J Pharmacol* 1991;103:1857-64.
- Bobker DH, Williams JT. Ion conductances by 5-HT receptor subtypes in mammalian neurons. *Trend Neurosci* 1990;13:169-73.
- Bonnett KA, Peterson KE. A modification of the jump-flinch technique for measuring pain sensitivity in rats. *Pharmacol Biochem Behav* 1975;3:1-47.
- Bouhelal R, Smounya L, Bockaert J. 5-HT<sub>1B</sub> receptors are negatively coupled with adenylate cyclase in rat substantia nigra. *Eur J Pharmacol* 1988;151:189-96.
- Boulenguez P, Chauveau J, Segu L, Morel A, Lanoir J, Delaage M.

- Biochemical and pharmacological characterization of serotonin-O-carboxymethylglycyl [<sup>125</sup>] iodotyrosinamide, a new radioiodinated probe for 5-HT<sub>1B</sub> and 5-HT<sub>1D</sub> binding sites. **J Neurochem** 1992;58:951-9.
- Bradley PB, Engel G, Feniuk W, Fozard JR, Humphrey PPA, Middlemiss DN, et al. Proposals for the classification and nomenclature of functional receptor for 5-hydroxytryptamine. **Neuropharmacology** 1986;25:563-76.
- Bragin EO, Korneev AY, Vasilenko GF. Buspirone effect on the development of antinociceptive reactions. **Pain** 1989;36:257-61.
- Brandao ML, Lopez-Garcia JA, Graeff FG, Roberts MHT. Electrophysiological evidence for excitatory 5-HT<sub>2</sub> and depressant 5-HT<sub>1A</sub> receptors in neurons of the rat midbrain tectum. **Brain Res** 1991;556:259-66.
- Brewerton TD, Murphy DL, Mueller EA, Jimerson I. Induction of migraine-like headaches by the serotonin agonists m-chlorophenylpiperazine. **Clin Pharmacol Therap** 1988;43:605-9.
- Bruunvels AT, Landwehrmeyer B, Palacios JM, Moskowitz MA, Hoyer D. Localization of 5-HT<sub>1D</sub> and 5-HT<sub>1B</sub> receptor messenger RNA in rat brain and trigeminal ganglia. **Br J Pharmacol** 1993b;108:95P.
- Bruunvels AT, Palacios JM, Hoyer D. Pharmacological characterization and distribution of serotonin 5-HT<sub>1D</sub>-like and 5-HT<sub>1D</sub> binding sites in rat brain. **Naunyn Schmiedebergs Arch Pharmacol** 1993a;108:95P.
- Bullit E. Expression of c-fos-like protein as a marker for neuronal activity following noxious stimulation in the rat. **J Comp Neurol** 1990;296:517-30.
- Cervo L, Grignashi G, Samanin R. 8-Hydroxy-2-(di-n-propyl-amino) tetralin, a selective serotonin<sub>1A</sub> receptor agonist, reduces the immobility of rats in the forced swimming test by acting on the nucleus raphe dorsalis. **Eur J Pharmacol** 1988;158:53-9.
- Chapman CR, Casey KL, Dubner R, Foley KM, Gracely RH, Reading AE. Pain measurement : an overview. **Pain** 1985;22:1-31.

- Haleem DJ, Kennett GA, Whitton PS, Curzon G. 8-OH-DPAT increase coticosterone but not other 5-HT<sub>1A</sub> receptor-dependent responses more in females. *Eur J Pharmacol* 1989;164:435-51.
- Hamblin MW, Ac Guffin RW, Metcalf MA, Dorsa DM, Merchan KM. Distinct 5-HT<sub>1B</sub> and 5-HT<sub>1D</sub> serotonin receptors in rat : structure and pharmacological comparison of the two cloned receptors. *Mol Cell Neurosci* 1992;3:578-87.
- Hamblin MW, Metcalf MA. Primary structure and functional characterization of human 5-HT<sub>1D</sub> type serotonin receptor. *Mol Pharmacol* 1991;40:143-8.
- Hamel E, Fan E, Linville D, Ying V, Villemure JG, Chia LS. Expression of mRNA for the serotonin 5-hydroxytryptamine<sub>1Dβ</sub> receptor subtype in human and bovine cerebral arteries. *Mol Pharmacol* 1993b;44:242-6.
- Hamel E, Gregoire L, Lau B. 5-HT<sub>1</sub> receptors mediating contraction in bovine cerebral arteries : a model for human cerebrovascular 5-HT<sub>1Dβ</sub> receptors. *Eur J Pharmacol* 1993a;242:75-82.
- Hammond DL , Tyce GM , Taksh TL. Efflux of 5-hydroxytryptamine and noradrenaline into spinal cord super fusates during stimulation of the rat medulla. *J Physiol* 1985;359:151-62.
- Hamon M, Collin E, Chantrel D, Daval G, Verge D, Bourgoin S, et al. Serotonin receptors and the modulation of pain. In: Besson J-M, editor. *Serotonin and pain*. Amsterdam: Excerpta Medica, 1990:53-72.
- Harel-Dupas C, Cloez I, Fillion G. The inhibitory effect of trifluoromethyl phenylpiperazine on [<sup>3</sup>H]acetylcholine release in guinea-pig hippocampal synaptosomes is mediated by 5-hydroxytryptamine receptor distinct from 1A,1B and 1C subtypes. *J Neurochem* 1991;56:221-7.
- Harris GD, Zemlan FD, Murphy RM, Behbehani MM. Spinal cord 5-HT 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptor subtypes: relation to pain transmission. *Neurosci Abstr* 1986;12:1015.

- Hartig PR. Molecular biology of 5-HT receptors. **Trends Pharmacol Sci** 1989;10:64-9.
- Hartig PR , Branchek TA , Weinshank RL. A subfamily of 5HT<sub>1D</sub> receptor genes. **Trends Pharmacol sci** 1992;13:152-9.
- Hawiger J. Repertoire of platelet receptors. In: Hawiger J, editor. **Method of Enzymology**. San Diego: Academic Press, 1992;131-6.
- Headley PM , Duggan AW , Griersmith BT. Selective reduction by noradrenaline and 5-HT of nociceptive responses of cat dorsal horn neurons. **Brain Res** 1978;145:185-9.
- Herrick-Davies K, Titeler M. Detection and characterization of the serotonin 5-HT<sub>1D</sub> receptor in rat and human brain. **J Neurochem** 1988;50:1624-31.
- Heuring RE, Peroutka SJ. Characterization of a novel <sup>3</sup>H-5-hydroxytryptamine binding site subtype in bovine brain membranes. **J Neurosci** 1987;7:894-903.
- Hjorth S. Single dose 8-OH-DPAT pretreatment does not induce tachyphylaxis to the 5-HT release-reducing effect of 5-HT autoreceptor agonists. **Eur J Pharmacol** 1991;199:237-82.
- Hoffman BJ, Mezey E. Distribution of serotonin 5-HT<sub>1C</sub> receptor mRNA in adult rat brain. **FEBS Lett** 1989;247:453-62.
- Hole K, Berge O-G, Eide PK, Fasmer OB, Hunskaar S, Lund A, et al. Modifications in nociceptive reactivity following various serotonergic treatments: methodological difficulties. In: Besson J-M, editor. **Serotonin and pain**. Amsterdam: Excepta Medica, 1990:137-51.
- Hoyer D. Functional correlates of serotonin 5-HT<sub>1</sub> recognition sites. **J Recept Res** 1988a;8:59-81.
- Hoyer D , Clarke DE , Fozard JR , Harting PR , Martin GR , Mylecharane EJ , et al. International Union of Pharmacology classification of Receptors for 5-hydroxytryptamine (serotonin). **Pharmacol Rev** 1994;46:157-203.
- Hoyer D, Middlemiss DN. Species differences in the pharmacology of terminal

- 5-HT autoreceptors in mammalian brain. **Trends Pharmacol Sci** 1989a;10:131-2.
- Hoyer D, Pazos A, Probst A, Palacios JM. Serotonin receptors in the human brain. I. Characterization and autoradiographic localization of 5-HT<sub>1A</sub> recognition sites. Apparent absence of 5-HT<sub>1B</sub> recognition sites. **Brain Res** 1986a;376:85-96.
- Hoyer D, Schoeffter P. 5-HT<sub>1D</sub> receptor-mediated inhibition of Forskolin-stimulated adenylate cyclase activity in calf substantia nigra. **Eur J Pharmacol** 1988b;147:145-7.
- Hoyer D, Schoeffter P, Waeber C, Palacios JM. Serotonin 5-HT<sub>1D</sub> receptors. In: Whitaker-Azmitia PM, Peroutka SJ, editors. **The neuropharmacology of serotonin**. New York: NY Acad Sci 1990:168-81.
- Hoyer D, Waeber C, Karpf A, Neijt H, Palacios JM. [<sup>3</sup>H]ICS 205930 labels 5HT<sub>3</sub> recognition sites in membranes of cat and rabbit in pig choroid plexus; a pharmacological characterization. **Naunyn Schmiedebergs Arch Pharmacol** 1989b;339:252-8.
- Humphrey PPA, Hartig P, Hoyer D. A proposed new nomenclature for 5-HT receptors. **Trends Pharmacol Sci** 1993;14:223-36.
- Hunt SP, Pini A, Evan C. Introduction of c-fos like protein in spinal cord neurons following sensory stimulation. **Nature** 1987;328:632-4.
- Hutson PH, Tricklebank MD, Curzon G. Analgesia induced by brief footshock: Blockade by fenfluramine and 5-methoxy-N,N-dimethyltryptamine and prevention of blockade by 5-HT antagonist. **Brain Res** 1983;279:105-10.
- Hyden JLK, Wilcox GL. Intrathecal serotonin in mice : analgesia and inhibition of a spinal action of substance P. **Life sci** 1983;33:789-95.

- Iadarola MJ, Douglass J, Civelli O, Naranjo JR. Differential activation of spinal cord dynorphin and enkephalin neurons during hyperalgesia : evidence using cDNA hybridization. *Brain Res* 1988a;455:313-26.
- Iadarola MJ, Brady LS, Draisici G, Dubner R. Enhancement of dynorphin gene expression in spinal cord following experimental inflammation : stimulus specificity, behavioral parameters and opioid receptor binding. *Pain* 1988b; 45:313-26.
- Iverfeldt K, Perterson L-L, Brodin E, Ogren S-O, Bartfai T. Serotonin type-2 mediated regulation of substance P release in the ventral spinal cord and the effects of chronic antidepressant treatment. *Naunyn Schmiedebergs Arch Pharmacol* 1986;333:1-6.
- Iverson SD. 5-HT and anxiety. *Neuropharmacology* 1984;23:1553-60.
- Jin H, Oksenborg D, Askenazi A, Peroutka SJ, Duncan AMV, Rozmahel R, et al., Characterization of the human 5-hydroxytryptamine <sub>1B</sub> receptor. *J Biol Chem* 1992;267:5735-5738.
- Jordan LM , Kenshalo DR jr , Martin RF, Haber LH, Willis WD. Two populations of spinothalamic tract neurons with opposite responses to 5-HT. *Brain Res* 1979;164:342-6.
- Julius D, MacDermott AB, Axel R, Jessell TM. Molecular characterization of a functional cDNA encoding the serotonin receptor. *Science* 1988;214:558-64.
- Kalkman HO, Fozard JR. Further definition of the 5-HT receptor mediating contraction of rat stomach fundus : relation to 5-HT<sub>1D</sub> recognition sites. In : Fozard JR, Saxena PR, editors. *Serotonin : Molecular biology, Receptors and Functional Effects*. Birkhauser Verlag : Basel , 1991a:153-60.
- Kalkman HO, Fozard JR. 5-HT receptor types and their role in disease. *Curr Opin Neurol Neurosurg* 1991b;4:560-5.
- Katori M, Ikeda K, Harada Y, Uchida Y, Tanaka K, Ohishi S. A possible role

- of prostaglandins and bradykinin as trigger of exudation in carreinin-induced rat pleurisy. *Agents Actions* 1978;8:108-112.
- Kauman AJ, Parsons AA, Brown AM. Human arterial constrictor serotonin receptor. *Cardiovasc Res* 1993;27:2094-103.
- Kelder D, Ross SB. Long lasting attenuation of 8-OH-DPAT-induced corticosterone secretion after a single injection of a 5-HT<sub>1A</sub> presynaptic, but not postsynaptic receptor mediated responses ; relationship to antidepressant-like action. *Eur J Pharmacol* 1992;138:53-61.
- Kennett GA, Curzon G. Evidence that mCPP may have behavior effects mediated by central 5-HT<sub>1C</sub> receptors. *Br J Pharmacol* 1988;94:137-47.
- Kennett GA, Whitton P, Shah K , Curzon G. Anxiogenic-like effects of mCPP and TFMPP in animal models are opposed by 5-HT<sub>1C</sub> receptor antagonists. *Eur J Pharmacol* 1989;164:445-54.
- Kiefel JM , Rossi GC , Bodnar RJ. Medullary mu and delta opioid receptors modulate mesencephalic morphine analgesia in rats. *Brain Res* 1993;624:151-61.
- Kjøsvik A, Størkson r, Tjølsen a, Hole K. Differential effects of activation of lumbar and thoracic 5-HT<sub>2A/2C</sub> receptors on nociception in rats. *Pharmacol Biochem Behav* 1997;56:523-7.
- Kobilka BK, Frielle T, Collins S, Yang-Feng T, Kobilka TS, Francke U, et al., An intronless gene encoding a potential member of the family of receptors coupled to guanine nucleotide regulatory protein. *Nature* 1987;329:75-9.
- Koek W, Jackson A, Colpaert F. Behavioral pharmacology of antagonist at 5-HT<sub>2</sub>/5-HT<sub>1C</sub> receptors. *Neurosci Behav Rev* 1992;16:95-105.
- Kursar JD, Nelson DL, Wainscott DB, Cohen ML, Baez M. Molecular cloning, functional expression and pharmacological characterization of a novel serotonin receptor (5-hydroxytryptamine<sub>2F</sub>) from rat stomach fundus. *Mol Pharmacol* 1992;42:549-57.

- Le Bars D. serotonin and pain. In: Osborne NN , Hamon M, editors. **Neuronal serotonin.** New York : John Wiley 1988:171-229.
- Leysen JE, Niemegeers CJE, Van Nueten JM, Laduron PM. [<sup>3</sup>H] Ketanserin (R 41 468), a selective 3H-ligand for serotonin<sub>2</sub> receptor binding sites. **Mol Pharmacol** 1981;21:301-14.
- Leysen JE, Van Gompel P, Verwimp M, Niemegeers CJE. Role and localization of serotonin (S<sub>2</sub>) - receptor binding sites : effects of neuronal lesions. In: Mandel P, de Feudis FV, editors. **CNS-Receptors-from molecular pharmacology to behavior.** New York, Raven Press 1983:373-83.
- Levy FO, Gudermann T, Peres-Reyes E, Birnbaumer M, Kaumann AJ, Birnbaumer L. Molecular cloning of a human serotonin receptor (S12) with a pharmacological profile resembling that of the 5-HT<sub>1D</sub> subtype. **J Biol Chem** 1992;267:7553-62.
- Libert F, Parmentier M, Lefort A, Dinsart C, Van Sande J, Maenhaut C, et al. Selective amplification and cloning of four new members of the G-protein-coupled receptor family. **Science** 1989;244:569-72.
- Light AR, Perl ER. Spinal termination of functionally identified primary afferent neurons with slowly conducting myelinated fibers. **J Comp Neurol** 1979;186:133-50.
- Limberger N, Deicher R, Starke K. Species differences in presynaptic serotonin autoreceptors: mainly 5-HT<sub>1B</sub> but possibly in addition 5-HT<sub>1D</sub> in the rat, 5-HT<sub>1B</sub> in the rabbit and guinea-pig brain cortex. **Naunyn Schmiedebergs Arch Pharmacol** 1991;343:353-64.
- Loric S, Launay JM, Colas JF, Maroteaux L. New mouse 5-HT<sub>2</sub>-like receptor. Expression in brain, heart and intestine. **FEBS Lett** 1992;312:203-7.
- Lubbert H, Hoffman Bj, Snutch TP, VanDyke t, Levine AJ, Hartig PR, et al. cDNA cloning of a serotonin 5-HT<sub>1C</sub> receptor by electrophysiological assays of mRNA-injected *Xenopus* oocytes. **Proc Natl Acad Sci** 1987a;84:4322-36.

- Lubbert H, Snutch TP, Dascal N, Lester HA, Davidson N. Rat brain 5-HT<sub>1C</sub> receptors are encoded by 5-6 kbase mRNA size class and are functionally expressed in injected *Xenopus* oocytes. **J Neurosci** 1987b;7:1159-65.
- Lucki I. 5-HT<sub>1</sub> receptors and behavior. **Neurosci Biobehav Rev** 1992;16:83-93.
- Maenhaut C, Van Sande J, Massart C, Dinsart C, Libert F, Monferini E, et al. The orphan receptor cDNA RDC4 encodes a 5-HT<sub>1D</sub> serotonin receptor. **Biochem Biophys Res Commun** 1991;180:1460-8.
- Magnaini E, Berribi JI, Morgan JI, Fos-like immunoreactivity induced by seizure in mice is specifically associated with enchromatin in neurons. **Eur J Neurosci** 1989;1:46-52.
- Manson P, Feilds HL. Central Modulation of Nociception. In: Olesen J, Tfelt-Hansen P, editors. **The headache**. New York : Raven Press , 1993:79-85.
- Marcinkiewicz M, Verge D, Gozlan H, Pichat L, Hamon M. Autoradiographic evidence for the heterogeneity of 5-HT<sub>1</sub> sites in the rat brain. **Brain Res** 1984;291:159-63.
- Marlier L, Teilhac JR, Cerruti C, Privat A. Autoradiographic mapping of 5-HT<sub>1</sub>, 5-HT<sub>1A</sub>, 5-HT<sub>1B</sub> and 5-HT<sub>2</sub> receptors in the rat spinal cord. **Brain Res** 1991;550:15-23.
- Maroteaux L, Saudou F, Amlaiky N, Boschert U, Plassat JL, Hen R. The mouse 5-HT<sub>1B</sub> serotonin receptor: cloning, functional expression and localization in motor control centers. **Proc Natl Acad Sci** 1992;89:3020-4.
- Matsuda T, Kanda T, Seong YH, Bada A, Iwata H. *p*-chlorophenylalanine attenuates the pituitary-adrenocortical response to 5-HT<sub>1A</sub> receptor agonists in mice. **Eur J Pharmacol** 1990;181:295-60.
- Maura G, Raiteri M. Cholinergic terminals in rat hippocampus possess 5-HT<sub>1B</sub> receptors mediating inhibition of acetylcholine release. **Eur J Pharmacol** 1986;129:333-7.

- Maura G, Roccatagliata E, Ulivi M, Rateri M. Serotonin-glutamate interaction in rat cerebellum : involvement of 5-HT<sub>1</sub> and 5-HT<sub>2</sub> receptors. **Eur J Pharmacol** 1988;145:31-8.
- McGrath PA, Sharav Y, Dubner R, Gracely RH. Masseter inhibitory periods and sensations evoked by electrical tooth pulp stimulation. **Pain** 1981;10:1-17.
- McKearney JW. Apparent antinociceptive properties of piperazine-type Serotonin agonists:trifluoromethyl-phenyl-piperazine, Chlorophenyl-piperazine, and MK-212. **Pharmacol Biochem Behav** 1989;32:657-60.
- Meibach RC, Maayani S, Green JP. Characterization and radioautography of [<sup>3</sup>H]LSD binding by rat brain slices in vitro : the effect of 5-hydroxytryptamine. **Eur J Pharmacol** 1980;67:371-82.
- Meller E, Bohmker K. Differential receptor reserve for 5-HT<sub>1A</sub> receptor mediated regulation of plasma neuroendocrine hormones. **J Pharmacol Exp ther** 1994;274:1246-53.
- Meller E, Goldstein M, Bohmker K. Receptor reserve for 5-hydroxytryptamine<sub>1A</sub>-mediated inhibition of serotonin synthesis : possible relationship to anxiolytic properties of 5-hydroxytryptamine agonists. **Mol Pharmacol** 1990;37:231-6.
- Meller E, Chalfin M, Bohmker K. Serotonin 5-HT<sub>1A</sub> receptor-mediated hypothermia in mice : absence of spare receptors and rapid induction of tolerance. **Pharmacol Biochem Behav** 1992;43:405-52.
- Menetrey D, Gannon A, Levine JD, Basbaum AI. Expression of c-fos protein in interneurons and projection neurons somatic, articular and visceral stimulation. **J Comp Neurol** 1989;285:177-95.
- Mengod G, Nguyen H, Waeber C, Lubbert H, Palacios JM. The distribution and cellular localization of the serotonin<sub>1C</sub> receptor mRNA in the rodent brain examined by *in situ* hybridization histochemistry. Comparison with receptor binding distribution. **Neuroscience** 1990a;35:577-91.
- Messing RB , Lytte LD. Serotonin containing neurons: their possible role in

- pain and analgesia. *Pain* 1977;4:1-21.
- Middlemiss DN. Blockade of the central 5-HT autoreceptor by -adrenoceptor antagonist. *Eur J Pharmacol* 1986;120:51-4.
- Middlemiss DN, Fozard JR. 8-hydroxy-2-(di-n-propyl-amino)-tetralin discriminates between subtypes of the 5-HT<sub>1</sub> recognition site. *Eur J Pharmacol* 1983;90:151-3.
- Millan MJ. Serotonin and pain : evidence that activation of 5-HT<sub>1A</sub> receptors does not elicit antinociception against noxious thermal, mechanical and chemical stimuli in mice. *Pain* 1994;58:45-61.
- Millan MJ, Colpaert FC. 5-hydroxytryptamine (5-HT)<sub>1A</sub> receptors and the tail-flick response. II. High efficacy 5-HT<sub>1A</sub> agonists attenuate morphine-induced antinociception in mice in a competitive-like manner. *J Pharmacol Exp Ther* 1991;256:983-92.
- Mitchell CL. A comparison of drug effects upon the jaw jerk response to electrical stimulation of the tooth pulp in dogs and cats. *J Pharmacol Exp ther* 1964;146:1-6.
- Mitchell R, Riley S. Characterization receptors for 5-hydroxytryptamine in spinal cord. *Trans Biochem Soc* 1985;13:955.
- Mjellem N, Lund A, Eide PK, Storkson R, Tjølsen A. The role of 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptors in spinal nociceptive transmission and in the modulation of NMDA-induced behavior. *Neuroreport* 1992;3:1061-4.
- Molander C, Grant G. Cutaneous projections from the rat hindlimb foot to the substantia gelatinosa of the spinal cord studied by transganglionic transport of WGA-HRP conjugate. *J Comp Neurol* 1985;237:476-84.
- Monroe PJ, Smith DJ. Characterization of [<sup>3</sup>H] 5-hydroxytryptamine binding sites in rat spinal cord tissue. *J Neurochem* 1983;41:349-55.
- Morgan JI, Cohen DR, Hemstead JL, Curran T. Mapping patterns of c-fos expression in central nervous system after seizure. *Science* 1987;237:192-7.
- Morgan JI, Curran T. Role of ion influx in the control of c-fos expression.

- Nature 1986;322:522-55.
- Mugnaini E, Beribi JI, Morgan JL. Fos-like immunoreactivity induced by seizure in mice is specifically associated with euchmatin in neurons. Eur J Neurosci 1989;1:46-52.
- Murphy RM, Zemlan FP. Differential effects of substance P on serotonin-modulated spinal nociceptive reflexes. Psychopharmacology 1987;93:118-21.
- Murphy RM, Zemlan FP. Selective serotonin<sub>1A/1B</sub> agonists differentially affect spinal nociceptive reflexes. Neuropharmacology 1990;29:463-8.
- North RA, Uchimura N. 5-Hydroxytryptamine acts at 5-HT<sub>2</sub> receptors to decrease potassium conductance in rat nucleus accumbens neurons. J Physiol 1989;417:1-12.
- Nozaki K, Boccalini P, Moskowitz M. Expression of c-fos like immunoreactivity in brainstem after meningeal irritation by blood in the subarachnoid space. Neuroscience 1992a;49:669-80.
- Nozaki K, Moskowitz MA, Boccalini P. CP-93,129,sumatriptan, dihydro-ergotamine block c-fos expression within rat trigeminal nucleus caudalis caused by chemical stimulation of the meninges. Br J Pharmacol 1992b;106:409-15.
- Oksenberg K, Marsters SA, O'Dowd BF, Jin H, Havlik S, Peroutka SJ, Ashkenazi A. A single amino acid difference confers major pharmacological variation between human and rodent 5-HT<sub>1B</sub> receptors. Nature 1992;360:161-3.
- Oliveras JL, Redjemi G, Guibaud G, Besson JM. Analgesia induced by electrical stimulation of the inferior centralis nucleus of the raphe in the cat. Pain 1975;1:139-45.
- Oliveras JL, Maixner W, Dubner R. The medullary dorsal horn : a target for the expression of opiate effects on the perceived intensity of noxious heat. J Neurosci 1986;6:3086-93.
- Palacios JM, Waeber C, Bruinvels AT, Hoyer D. Direct visualization of

- serotonin<sub>1D</sub> receptors in the human brain using a new iodinated ligand.  
*Mol Brain Res* 1992;346:175-9.
- Pan ZZ , Feilds HL. Endogenous opioids acting at a medullary mu opioids receptor contribute to the behavioral antinociception produced by GABA antagonism in the midbrain periaqueductal gray. 1996;in press.
- Paxinos G, Watson C, editors. *The Rat Brain in Stereotaxic Coordinates*. Sydney : Academic Press, 1982.
- Pazos A, Cortes M, Palacios JM. Quantitative autoradiographic mapping of serotonin receptors in the rat brain. II.Serotonin2- receptors. *Brain Res* 1985;346:231-49.
- Pazos A, Hoyer D, Palacios JM. The binding of serotonergic ligands to the porcine choroid plexus : characterization of a new type of serotonin recognition site. *Eur J Pharmacol* 1984;106:539-46.
- Pazos A, Palacios JM. Quantitative autoradiographic mapping of serotonin receptors in the rat brain. I. Serotonin-1 receptors. *Brain Res* 1985;346:205-30.
- Pazos A, Probst A, Palacios JM. Serotonin receptors in the human brain. III. Autoradiographic mapping of serotonin-1 receptors. *Neuroscience* 1987a;21:97-122.
- Pazos A, Probst A, Palacios JM. Serotonin receptors in the human brain. IV. Autoradiographic mapping of serotonin-2-receptors. *Neuroscience* 1987b;21:123-39.
- Pernow B. Substance P. *Pharmacol Rev* 1983;35:85-141.
- Peroutka SJ. 5-Hydroxytryptamine receptors. *J Neurochem* 1993;60:408-16.
- Peroutka SJ, Snyder SH. Multiple serotonin receptors: differential binding of [<sup>3</sup>H]5-HT, [<sup>3</sup>H]lysergic acid diethylamide and [<sup>3</sup>H]spiroperidol. *Mol Pharmacol* 1979;16:687-99.
- Post C, Archer T. Interactions between 5-HT and noradrenaline in analgesia. In: Besson JM, editor. *Serotonin and Pain*. Amsterdam:Excerpta, 1990:153-74.

Pritchett DB, Bach AEJ, Wozny M, Taleb O, Dal Toso R, Shin JC, et al.

Structure and functional expression of cloned rat serotonin 5-HT<sub>2</sub> receptor. **EMBO J** 1988;7:4135-40.

Radja F, Laporte AM, Daval G, Verge D, Gozlan H, Hamon M.

Autoradiography of serotonin receptor subtypes in the central nervous system. **Neurochem Int** 1991;18:1-15.

Raiteri M, Maura G, Bonanno G, Pittaluga A. Differential pharmacology and function of two 5-HT<sub>1</sub> receptors modulating transmitter release in rat cerebellum. **J Pharmacol Exp Ther** 1986;237:644-8.

Randic M, Yu HH. Effects of 5-hydroxytryptamine and bradykinin in cat dorsal horn neurons activated by noxious stimuli. **Brain Res** 1976;111:197-203.

Ren MF, Han JS. Rat tail flick acupuncture analgesia model. **Chinese Med J** 1979;92:576-82.

Rexed B. A cytoarchitectonic atlas of the spinal cord in the cat. **J Comp Neurol** 1954;100(2):297-379.

Reynolds DV. Surgery in the rat during electrical analgesia induced by focal brain stimulation. **Science** 1969;164:444-5.

Rivot JP, Chiang CY, Besson JM. Increase of serotonin metabolism within the dorsal horn of the spinal cord during nucleus raphe magnus stimulation, as revealed by in vivo electrochemical detection. **Brain Res** 1980;238:117-26.

Robert MHL. Pharmacology of putative neurotransmitters and receptors : 5-hydroxytryptamine. In: Feilds HL, Besson J-M, editors. **Pain modulation**. Amsterdam : Elsevier, 1988:329-38.

Robert MHT, Davies M. In vivo electrophysiology of receptors mediating the central nervous system actions of 5-hydroxytryptamine. In : Mylecharane EJ, Angus JA, de la Lande IS, Humphrey PPA, editors. **Serotonin : Actions , Receptors and Pathophysiology**. Macmillan : London, 1989:70-6.

- Rodgers RJ, Shepherd JK. Prevention of the analgesic consequences of social defeat in male mice by 5-HT<sub>1A</sub> anxiolytics, buspirone, gepirone and psapirone. *Psychopharmacology* 1989;33:674-80.
- Roth B, Nakaki T, Chuang DM, Costa E. Aortic recognition sites for serotonin (5-HT) are coupled to phospholipase C and modulate phosphatidylinositol turnover. *Neuropharmacol* 1984;23:1223-5.
- Roychowdhury SM, Feilds HL. Endogenous opioids acting at a medullary mu opioids receptor contribute to the behavioral antinociception produced by GABA antagonism in the midbrain periaqueductal gray. 1996;in press.
- Sagar SM, Sharp FR, Curran T. Expression of c-fos protein in brain : Metabolic mapping at the cellular level. *Science* 1988;240:1328-31.
- Sahin-Erdemli L, Schoeffter P, Hoyer D. Competitive antagonism by recognized 5-HT<sub>2</sub> receptor antagonists a 5-HT<sub>1C</sub> receptors in pig choroid plexus. *Naunyn Schmiedebergs Ach Pharmacol* 1991;344:137-42.
- Saltzmann AG, Morse B, Whitman MM, Ivanschchenko Y, Jaye M, Felder S. Cloning of the human serotonin 5-HT<sub>2</sub> and 5-HT<sub>1C</sub> receptor subtypes. *Biochem Biophys Res Commun* 1991;181:1469-78.
- Sandkuhler J, Gebhart GF. Characterization of inhibition of a spinal nociceptive reflex by stimulation medially and laterally in the midbrain and medulla in the pentobarbital-anesthetized rat. *Brain Res* 1984;305:67-76.
- Sambucetti LC, Curran T. The fos protein complex is associated with DNA in isolated nuclei and binds to DNA cellulose. *Science* 1986;234:1417-9.
- Sawynok J. The role of ascending and descending noreadrenergic and serotonergic pathways in opioid and nonopioid antinociception as revealed by lesion studies. *Can J Physio Pharmacol* 1989;87:975-88.
- Sawynok J, Reid A. Spinal supersensitivity to 5-HT<sub>1</sub>, 5-HT<sub>2</sub> and 5-HT<sub>3</sub> receptor agonists following 5,7-dihydroxytryptamine. *Eur J Pharmacol* 1994;264:249-57.

- Saxena PR, Villalon CM. Cardiovascular effects of serotonin agonists and antagonists. *J Cardiovasc Pharmacol* 1990;15 suppl 7:S17-S34.
- Schmauss C, Hammond DL, Ochi JW. Pharmacological antagonism of the antinociceptive effects of serotonin in the rat and spinal cord. *Eur J Pharmacol* 1983;90:349-57.
- Schmauss C, Hammond DL, Ochi JW, Yasksh TL. Pharmacological antagonism of the antinociceptive effects of serotonin in the rat spinal cord. *Eur J Pharmacol* 1983;90:349-57.
- Schlicker E, Fink K, Gothert M, Hoyer D, Middlemiss G, Roschke I, et al. The pharmacological properties of the presynaptic 5-HT autoreceptor in the pig brain cortex conform to the 5-HT<sub>1D</sub> receptor subtype. *Naunyn Schmiedebergs Ach Pharmacol* 1989;340:45-51.
- Schoeffter P, Hoyer D. Centrally acting hypotensive agents with affinity for 5-HT<sub>1A</sub> binding sites inhibit forskolin-stimulated adenylate cyclase activity in calf hippocampus. *Br J Pharmacol* 1988;95:975-85.
- Schoeffter P, Hoyer D. 5-hydroxytryptamine 5-HT<sub>1B</sub> and 5-HT<sub>1D</sub> receptors mediating inhibition of adenylate cyclase activity. Pharmacological comparison with special reference to the effects of yohimbine, rauwolscine and some β-adrenoreceptor antagonist. *Naunyn Schmiedebergs Ach Pharmacol* 1989a;340:285-92.
- Schoeffter P, Hoyer D. Interaction of arylpiperazine with 5-HT<sub>1A</sub>, 5-HT<sub>1B</sub>, 5-HT<sub>1C</sub> and 5-HT<sub>1D</sub> receptors: do discriminatory 5-HT<sub>1B</sub> receptor ligand exist? *Naunyn Schmiedebergs Arch Pharmacol* 1989b;339:675-83.
- Schoeffter P, Waeber C, Palacios JM, Hoyer D. The serotonin 5-HT<sub>1D</sub> receptor subtype is negative coupled to adenylate cyclase in calf substantia nigra. *Naunyn schmiedebergs Arch Pharmacol* 1988;337:602-8.
- Secrest RJ, Lucaites VL, Mendelsohn LG, Cohen ML. Protein kinase C translocation in rat stomach fundus : effects of serotonin, carbamylcholine and phorbol dibutyrate. *J Pharmacol Exp Ther* 1991;256:103-9.

- Segu L, Chauveau J, Boulenguez P, Morel A, Lanoir J, Delaage M. Synthesis and pharmacological study of radioiodinated serotonin derivative specific for 5-HT<sub>1B</sub> and 5-HT<sub>1D</sub> binding sites in the central nervous system. *CR Acad Sci* 1991;312:655-61.
- Shenker A, Maayani S, Weinstein H, Green JP. Enhanced serotonin-stimulated adenylate cyclase activity in membranes from adult guinea-pig hippocampus. *Life sci* 1983;32:2335-42.
- Shenker A, Maayani S, Weinstein H, Green JP. Two 5-HT receptors linked to adenylate cyclase in guinea-pig hippocampus are discriminated by 5-carboxaminotryptamine and spiperone. *Eur J Pharmacol* 1985;109:427-9.
- Shenker A, Maayani S, Weinstein H, Green JP. Pharmacological characterization of two 5-hydroxytryptamine receptors coupled to adenylate cyclase in guinea-pig hippocampus membrane. *Mol Pharmacol* 1987;31:357-67.
- Shibata M, Ohkubo Y, Takahashi H, Inoki R. Interaction of bradykinin with substance P on vascular permeability and pain response. *Jpn J Pharmacol* 1986;41:427-9.
- Shibata M, Ohkubo Y, Takahashi H, Inoki R. Modified formalin test : characteristic biphasic pain response. *Pain* 1989;38:347-52.
- Solomon RE, Gebhart GF. Mechanisms of effects of intrathecal serotonin on nociception and blood pressure in rats. *J Pharmacol Exp Ther* 1988;245:905-12.
- Stephen D, Siberstein SD. Serotonin(5-HT) and migraine. *Headache* 1994;34:408-17.
- Sugimura Y, Lee CL, Perl ER. Central projections of identified, unmyelinated (C) afferent fibers innervating mammalian skin. *Science* 1987;234:358-61.
- Swett JE, Woolf CJ. The somatotopic organization of primary afferent terminals

- in the superficial lamina of the dorsal horn of the rat spinal cord. **J Comp Neurol** 1985;231:66-77.
- Szkely AM, Barbaccia ML, Costa E. Activation of specific glutamate receptor subtypes increases c-fos proto-oncogene expression in primary cultures of neonatal rat cerebellar granule cells. **Neuropharmacology** 1987;26:1779-82.
- Tenen SS. Antagonism of the analgesic effect of morphine and other drugs by p-chlorophenylalanine , a Serotonin depleter. **Psychopharmacologia** 1968;12:278-85.
- Titeler M, Lyon RA, Glennon RA. Radioligand binding evidence implicates the brain 5-HT<sub>2</sub> receptors as a site of action for LSD and phenylisopropylamine hallucionogens. **Psychopharmacology** 1988; 94:213-16.
- Tjølsen A, Berge O-G, Hunskaar S, Rosland JH, Hole K. The formalin test : and evaluation of the method. **Pain** 1992;51:5-17.
- Traber J, Glaser T. 5-HT<sub>1A</sub> receptor-related anxiolytics. **Trends Pharmacol Sci** 1987;8:432-7.
- Tjølsen A, Lund A, Eide PK, Berge OG, Hole K. The apparent hyperalgesic effect of a serotonin antagonist in the tail flick test is mainly due to increased tail skin temperature. **Pharmacol Biochem Behav** 1989;32:601-8.
- Trickelband MD, Forler C, Middlemiss DN, Fozard JR. Subtypes of the 5-HT receptor mediating the behavioral responses to 5-methoxy-N,N-dimethyltryptamine in the rat. **Eur J pharmacol** 1985;117:15-24.
- Trickelband MD. The behavior response to 5-HT receptor agonists and subtypes of the central 5-HT receptor. **Trends Pharmacol Sci** 1985;6:403-7.
- Vane JR. The relative activities of some tryptamine analogues on the isolated rat stomach strip preparation. **Br J Pharmacol Chemother** 1959;14:87-98.

- Veirck CJ Jr, Hamilton DM, Thornby JI. Pain reactivity of monkeys after lesions to the dorsal and lateral columns of the spinal cord. *Exp Brain Res* 1971;13:140-58.
- Verge D, Daval G, Marcinkiewicz M, Patey A, El Mestikaway S, Gozlan H, Hamon M. Quantitative autoradiography of multiple 5-HT<sub>1</sub> receptor subtypes in the brain of control or 5,7-dihydroxytryptamine treated rats. *J Neurosci* 1986;6:3473-82.
- Vierck CJ Jr, Cooper BY. Guidelines for assessing pain reaction and pain modulation in laboratory animal subjects. In : Kruger L, Leibeskind JC, Cooper BY, editors. *Advances in pain research and therapy*. New York : Raven Press; 1984:305-22.
- Vogt M. The effects of lowering the 5-hydroxytryptamine content of the rat spinal cord on analgesia produced by morphine. *J Physiol* 1974;483-98.
- Voigt MM, Laurie DJ, Seeburg PH, Bach A. Molecular cloning and characterization of a rat brain cDNA encoding 5-hydroxytryptamine<sub>1B</sub> receptor. *EMBO J* 1991;10:4017-23.
- Waeber C, Dietl MM, Hoyer D, Palacios JM. 5-HT<sub>1</sub> receptors in the vertebrate brain : regional distribution examined by autoradiography. *Naunyn Schmiedebergs Arch Pharmacol* 1989a;340:486-94.
- Waeber C, Dietl MM, Hoyer D, Probst A, Palacios JM. Visualization of a novel serotonin recognition site 5-HT<sub>1D</sub> in the human brain by autoradiography. *Neurosci Lett* 1988a;88:11-6.
- Waeber C, Hoyer D, Palacios JM. GR 43175 : A preferential 5-HT<sub>1D</sub> agent in monkey and human brains as shown by autoradiography. *Synapse* 1989a;4:168-70.
- Waeber C, Schoeffter P, Hoyer D, Palacios JM. The Serotonin 5-HT<sub>1D</sub> receptor : a progress review. *Neurochem Res* 1990;15:567-82.
- Waeber C, Schoeffter P, Palacios JM, Hoyer D. Molecular Pharmacology of 5-

- HT<sub>1D</sub> recognition sites : radioligand binding studies in human, pig and calf brain membranes. *Naunyn Schmiedebergs Arch Pharmacol* 1988c;337:595-601.
- Waeber C, Schoeffter P, Palacios JM, Hoyer D. 5-HT<sub>1D</sub> receptors in the guinea-pig and pigeon brain: radioligand binding and biochemical studies. *Naunyn Schmiedebergs Arch Pharmacol* 1989d;340:479-85.
- Wang HY, Eberle-Wang K, Simansky KJ, Friedman E. Serotonin-induced muscle contraction in rat stomach fundus is mediated by a G<sub>αz</sub>-like guanine nucleotide binding protein. *J Pharmacol Exp Ther* 1993;267:1002-11.
- Wang JK. Antinociceptive effect of intrathecally administered serotonin. *Anaesthesiology* 1977;47:269-71.
- Ware JA, Heistad DD. Platelet-endothelium interaction. *N Eng J Med* 1993, 328:628-35.
- Weinshank RL, Zgombick JM, Macchi MJ, Branchek TA, Hartig PR. Human serotonin<sub>1D</sub> receptor is encoded by subfamily of two distinct genes : 5-HT<sub>1Dα</sub> and 5-HT<sub>1Dβ</sub>. *Proc Natl Acad Sci* 1992;89:3630-4.
- Weishank RL, Zgombick JM, Macchi MJ, Branchek TA, Hartig PR. Human serotonin<sub>1D</sub> receptor is encoded by a subfamily of two distinct genes : 5-HT<sub>1Dα</sub> and 5-HT<sub>1Dβ</sub>. *Proc Natl Acad Sci* 1992;89:3630-4.
- Weiss B, Laties VG. Characteristics of aversive thresholds measured by a titration schedule. *J Exp Anal Behav* 1963;6:563-72.
- Weiss S, Sebben M, Kemp DE, Bockaert J. Serotonin 5-HT<sub>1</sub> receptors mediate inhibition of cyclic AMP production in neurons. *Eur J Pharmacol* 1986;120:227-30.
- Wieland S, Lucki I. Antidepressant-like activity of 5-HT<sub>1A</sub> agonists measured with the forced swim test. *Psychopharmacology* 1990;101:497-504.
- Wilcox GL, Alhaider AA. Nociceptive and antinociceptive action of serotonergic agonists administered intrathecally. In: Besson J-M, editor. *Serotonin and pain*. Amsterdam: Elsevier, 1990:205-19.

- Willer JC, Boureau F, Albe-Fessard D. Supraspinal influences on nociceptive flexion reflex and pain sensation in man. *Brain Res* 1979;179:61-8.
- Wisden W, Hunt SP. Induction of early onset genes in spinal cord neurons following sensory stimulation. *Neurosci Lett Suppl* 1989;36:S75.
- Wong-Riley MT, Kageyama GH. Localization of cytochrome oxidase in the mammalian spinal cord and dorsal root ganglia with quantitative analysis of ventral horn cells in marmosets. *J Comp Neurol* 1986;245:41-61.
- Xu W, Qiu XC, Han JS. Serotonin receptor subtypes in spinal antinociception in the rat. *J Pharmacol Exp Ther* 1994;264:1182-9.
- Yagaloff KA, Hartig PR. [ $^{125}\text{I}$ ] Lysergic acid diethylamide binds to a novel serotonin site on rat choroid plexus epithelial cells. *J Neurosci* 1985;5:3178-83.
- Yaksh TL. Direct evidence that spinal serotonin and noradrenaline terminals mediate the spinal antinociceptive effects of morphine in the periaqueductal gray. *Brain Res* 1979;160:180-5.
- Yaksh TL. Pharmacology of spinal adrenergic system which modulates spinal nociceptive processing. *Pharmacol Biochem Behav* 1988;22:845-58.
- Yaksh TL, Wilson PR. Spinal serotonin terminal system mediates antinociception. *J Pharmacol Exp Ther* 1979;208:446-53.
- Yu L, Nguyen H, Le H, Bloem LJ, Kozak CA, Hoffman BJ, et al. The mouse 5-HT<sub>1C</sub> receptor contains eight hydrophobic domains and is X-linked. *Mol Brain Res* 1991;11:143-9.
- Zemlan FP, Corrigan SA, Pfaff DW. Noradrenergic and serotonergic mediation of spinal analgesia mechanisms. *Eur J Pharmacol* 1980;61:111-24.
- Zemlan FP, Kow LM, Pfaff DW. Spinal serotonin (5-HT) receptor subtypes and nociception. *J Pharmacol Exp Ther* 1983;226:477-85.
- Zemlan FP, Behbehani MM, Murphy RM. Serotonin receptor subtypes and the modulation of pain transmission. *Prog Brain Res* 1988;77:349-55.
- Zgombick JM, Borden LA, Cochran TL, Kucharewicz SA, Weinshank RL, Branchek TA. Dual coupling of cloned human 5-hydroxytryptamine<sub>1Dα</sub>

and 5-hydroxytryptamine<sub>1Dβ</sub> receptors stable expressed in murined fibroblasts : inhibition of adenylyl cyclase and elevation of intracellular calcium concentration via pertussis toxin-sensitive g-protein(s). **Mol Pharmacol** 1993;44:575-82.

Zgombick JM, Weinshank RL, Macchi M, Schechter LE, Branchek TA, Hartig PR. Expression and pharmacological characterization of a canine 5-hydroxytryptamine<sub>1D</sub> receptor subtype. **Mol Pharmacol** 1991;40:1036-42.

สถาบันวิทยบริการ  
จุฬาลงกรณ์มหาวิทยาลัย

## BIOGRAPHY

Miss Unchalee Ruangpattanatawee was born on September, 5<sup>th</sup> 1967. She graduated Bachelor of Science in Physical Therapy from school of Physical Therapy, Faculty of Medicine, Siriraj Hospital, Mahidol University in 1990.



สถาบันวิทยบริการ  
จุฬาลงกรณ์มหาวิทยาลัย