

## REFERENCES

- Abu-Arish, A., Frenkiel-Krispin, D., Fricke, T., Tzfira, T., Citovsky, V., Grayer Wolf, S and Elbaum, M. 2004. Three-dimensional reconstruction of *Agrobacterium* VirE2 protein with single-stranded DNA. **J Biol Chem.** 279:25359-25363.
- Acton, N. and Klayman, D.L. 1985. Artemisitene, a new sesquiterpene lactone endoperoxide from *Artemisia annua*. **Planta Med.** 51:441-442.
- Acton, N. and Klayman, D.L. 1986. Conversion of artemisinin (Qinghaosu) to iso-artemisitenene and to 9-epi-artemisinin. **Planta Med.** 53:266-268.
- Acton, N., Klayman, D., and Rollman, I. 1985. Reductive electrochemical HPLC assay for artemisinin (Qinghaosu). **Planta Med.** 51:445-445.
- Acton, N., Klayman, D.L., Rollmann, I.J., and Novotny J.F. 1986. Isolation of artemisinin (qinghaosu) and its separation from artemisitene using the multi-layer coil separator extraction and isolation of arteannuin B. **J. Chromatogr.** 355: 448-450.
- Acton, N. and Roth, R., 1989. A sample conversion of artemisinic acid into artemisinin. **J. Nat. Prod.** 52:1183-1185.
- Adam, K.-P. and Zapp, J. 1998. Biosynthesis of the isoprene units of chamomile sesquiterpenes. **Phytochemistry** 48:953-959.
- Ahmad, A. and Misra, L.N. 1994. Terpenoids from *Artemisia annua* and constituents of its essential oil. **Phytochem.** 37:183-186.
- Akhila, A., Rani, K. and Thakur, R.S. 1990. Biosynthesis of artemisinic acid in *Artemisia annua*. **Phytochem.** 29:2129-2132.
- Akhila, A., Thakur, R.S. and Popoli, S.P. 1987. Biosynthesis of artemisinin in *Artemisia annua*. **Phytochem.** 26(7): 1927-1930.

- Alberts, B., Johnson, A., Lewis, J., Raff M., Roberts, K., and Walter, P. 2002. *Molecular Biology of The Cell*. 4th ed. New York: Garland Science.
- Allen, K. G., Banthorpe, D.V., Charlwood, B.V. and Voller, C.M. 1977. Biosynthesis of artemisia ketone in higher plants. **Phytochem.** 16: 79-83.
- Atchara T. 1996. Determination of artemisinin contents in *Artemisia annua* Linn. cultured in Thailand. Master's Thesis. Faculty of graduate studies. Mahidol University.
- Avery, M.A., Mehrotra, S., Johnson, T.L., Bank, J.D. Vroman, J.A. and Miller, R. 1996. Structure-activity relationships of the antimalarial agent artemisinin analogs of 10-deoxyartemisinin substituted at C-3 and C-9. **J. Med. Chem.** 36:4149-4155.
- Avery, B.A., Venkatesh K.K., and Avery M.A. 1999. Rapid determination of artemisinin and related analogues using high performance liquidchromatography an evaporative light scattering detector. **J. Chromatogr. B.** 730:71-80.
- Bertea CM, Freije JR, van der Woude H, Verstappen FW, Perk L, Marquez V, de Kraker JW, Posthumus MA, Jansen BJ, de Groot, A., Franssen, M.C., Bouwmeester, H.J. 2005. Identification of intermediates and enzymes involved in the early steps of artemisinin biosynthesis in *Artemisia annua*. **Planta Med.** 71:40-47.
- Birnboim, H.C. and Doly, J. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. **Nucleic Acids Res.** 7:1513-1523.
- Bohlmann, F. and Zdero, C. 1972. Constituents of *Artemisia afra*. **Phytochem.** 11:2329-2330.

- Boucher, Y. and Doolittle, W. F. 2000. The role of lateral gene transfer in the evolution of isoprenoid biosynthesis pathways. **Mol. Microbiol.** 37:703–716.
- Bouwmeester, H.J., Wallaart, T.E., Janssen, M.H.A., Loo, B.V., Jansen, B.J.M., Posthumus, M.A., Schmidt, C.O., Kraker, J.W.D., Konig, W.A. and Franssen, M.C.R. 1999. Amorpha-4,11-diene synthase catalyses the first probable step in artemisinin biosynthesis. **Phytochem.** 52:843-854.
- Bown, D. 2001. The herb society of America: new encyclopedia of herbs & their uses. New York: Dorling Kindersly Limited.
- Brossi, A., Venugopalan, B., Gerpe, D., Yeh, H.J.C., Flippen-Anderson, J.L., Buchs, P., Luo, X.D., Mithous, W. and Peter, W. 1988. Arteether, anew antimalarial drug: synthesis and antimalarial properties. **J. Med. Chem.** 31:645-650.
- Brown, G.D. 1994. Cadinanones from *Artemisia annua* that may be intermediates in the biosynthesis of artemisinin. **Phytochem.** 36:637-641.
- Brown, G.D., Liang, G.Y. and Sy, L.K. 2003. Terpenoids from the seeds of *Artemisia annua*. **Phytochem.** 64:303–323.
- Chan, K.L., Yuen, K.H., Jinadasa, S., Peh, K.K. and Toh, W.T. 1997. A high performance liquid chromatography analysis of plasma artemisinin using a glassy electrode for reductive electrochemical detection. **Plant Med.** 63:66-69.
- Chang, Y.J., Song, S.H., Park, S.H. and Kim, S.U. 2000. Amorpha-4,11-diene synthase of *Artemisia annua*: cDNA isolation and bacterial expression of a terpene synthase involved in artemisinin biosynthesis. **Arch. Biochem. Biophys.** 383:178-184
- Charles, D.J., Simon, J.E., Wood, K.V. and Heinstein, P. 1990. Germplasm variation in artemisinin content of *Artemisia annua* using an alternative method of artemisinin analysis from crude plant extracts. **J. Nat. Prod.** 53:157-160.

- Chen, H.L., Wang K.T., Pu, Q.S., Chen, X.G. and Hu, Z.D. 2002. On-line conversion and determination of artemisinin using a flow-injection capillary electrophoresis system. **Electrophoresis**. 23(17):2865-2871
- China Cooperative Research Group on Qinghaosu and Its Derivatives as Antimalarials. 1982. The chemistry and synthesis of qinghaosu derivatives. **J. Trad. Chin. Med.** 2:9-16.
- Christie, P.J. 2004. Type IV secretion: the *Agrobacterium* VirB/D4 and related conjugation systems. **Biochim Biophys Acta**. 1694:219-234.
- Covello, P.S., Teoh, K.H., Polichuk, D.R., Reed, D.W., Nowak, G. 2007. Functional genomics and the biosynthesis of artemisinin. **Phytochemistry**. 68:1864-1871.
- Citovsky, V., Wong, M.L. and Zambryski, P.C. 1989. Cooperative interaction of *Agrobacterium* VirE2 protein with single stranded DNA: implications for the T-DNA transfer process. **Proc Natl Acad Sci USA**. 86:1193-1197.
- Clyde, D.F. 1981. Clinical problems associated with the use of primaquine as a tissue schizontocidal and gametocytocidal drug. **Bull World Health Organ**. 59(3):391-395.
- Croteau, R., Kutchan, T. and Lewis, N. 2000. Natural products (secondary metabolites). In: Buchanan, B.: Grissem, W.: Jones, R., eds. Biochemistry and molecular biology of plants. Rockville, MD: American Society of Plant Physiologists:1250-1318.
- Draper, J., Scott, R., Armitage, P. and Walden, R: 1988. Plant Genetic Transformation and Gene Expression, A Laboratory Manual. London: Blackwell Scientific Publications Ltd.
- Edlund, P.O., Westerlund, D., Carlqvist, J., Bo-liang, W. and Yunhua, J. 1984. Determination of artesunate and dihydroartemisinin in plasma by liquid

- chromatography with post-column derivatization and UV-detection. **Acta Pharm. Succ.** 21: 223-234.
- El-Feraly, S. F., Al-Meshal, I.A., and Khalifa, I. 1989. Epi-deoxyarteannuin B and 6,7-dehydroartemisinic acid from *Artemisia annua*. **J. Nat. Prod.** 52(1):196-189.
- El-Feraly, S.F., Al-Meshal, I.A., Al-Yahya, M.A. and Hifnawy, M.S. 1986. On the possible role of qinghao acid in the biosynthesis of artemisinin. **Phytochem.** 25(12):2777-2778.
- El-Sohly, H.N., Croom, E.M. and El-Sohly, M.A. 1987. Analysis of the antimalarial sesquiterpene artemisinin in *Artemisia annua* by high performance liquid chromatography (HPLC) with post-column derivatization and ultraviolet detection. **Pharmacol Res.** 4:258-260.
- El-Sohly, H.N., Croom, E.M., El-Feraly, F.S. and El-Sherei, M.M. 1990. A large-scale extraction technique of artemisinin from *Artemisia annua*. **J. Nat. Prod.** 53(6):1560-1564.
- Farooqi, A.H.A., Shukla, A., Sharma, S. and Khan, A. 1996. Effect of plant age and GA3 on artemisinin and essential oil yield in *Artemisia annua* L. **J. herd, spices & Medicinal Plnats.** 4(1):73-80.
- Faure, O., Diemer, F., Moja, S. and Jullien, F. 1998. Mannitol and thidiazuron improve in vitro shoot regeneration from spearmint and peppermint leaf disks. **Plant Cell Tiss. Org. Cult.** 52:209-212.
- Ferreira, J. F. S. and Janick, J. 1996. Distribution of artemisinin in *Artemisinin annua*, p. 579-584. In: J. Janick (ed) Progress in new crops, ASHS Press, Arlington, V.A. Available from [http://www.hort.purdue.edu/new\\_crop/proceedings/1966/v3-578.html](http://www.hort.purdue.edu/new_crop/proceedings/1966/v3-578.html)



- Ferreira, J. F. S., Simon, J. E. and Janick, J. 1994a. Relationship of artemisinin content of tissue-cultured, greenhouse-grown, and field-grown plants of *Artemisia annua*. **Planta Med.** 61:351-355.
- Ferreira, J. F. S., Simon, J. E. and Janick, J. 1995. Developmental studies of *Artemisia annua*: flowering and artemisinin production under greenhouse and field conditions. **Planta Med.** 61:167-170.
- Ferreira, J.F.S. and Janick, J. 1996. Immunoquantitative analysis of artemisinin from *Artemisia annua* using polyclonal antibodies. **Phytochem.** 41(1):97-104.
- Ferreira, J.F.S., Charles, D.J., Wood, Karl, Janick, J. and Simon, J.E. 1994b. A comparison of gas chromatography and high performance liquid chromatography for artemisinin analysis. **Phytochem. Anal.** 5:116-120.
- Filichkin, S.A. and Gelvin, S.B. 1993. Formation of a putative relaxation intermediate during T-DNA processing directed by the *Agrobacterium tumefaciens* VirD1, D2 endonuclease. **Mol Microbiol.** 8:915-926.
- Gaudin, V., Vrain T. and Jouanin L: 1994. Bacterial genes modifying hormonal balances in plants. **Plant Physiol Biochem.** 32:11-29.
- Geldre, E.V., Vergacuwe, A. and Van den Eeckout, E. 1997. State of art of the production of the antimalarial compound artemisinin in plants. **Plant Mol. Biol.** 33:199-209.
- Gelvin, S.B. 1998. The introduction and expression of transgenes in plants. **Curr Opin Biotechnol.** 9:227-232.
- Gelvin, S.B. 2003. *Agrobacterium*-mediated plant transformation: the biology behind the 'gene-jockeying' tool. **Microbiol Mol Biol Rev.** 67:16-37.

- Green, M.D., Mont, D.L., Todd, G.D. and Capmacchia, A.C. 1999. Chemiluminescent detection of artemisinin-novel endoperoxide analysis using luminol with out hydrogen peroxide. **J. Chromtogr. A.** 695(2):237-242.
- Gupta, M.M., Jain, D.C., Mathur, A.K., Singh, A.K., Verma, R.K. and Kumar S. 1996. Isoltion of a high artemisinic acid containing plant of *Artemisia annua*. **Planta Med.** 62:280-281.
- Hasegawa, H., Takashima, S. and Nakamura, A. 1995. Effects of gamma-ray irradiation on cultured anthers of tobacco (*Nicotiana tobacum* L.)- radiosensitivity and morphological variants appearing in the haploid plants. **Plnat Tiss Cult Lett.** 12(3):281-287.
- Haynes, R.K. and Vonwiller, S.C. 1996. The behaviors of artemisinin in presence of non-heme iron (II) and (III). **Tetrahedrol Lett.** 37(2):257-260.
- Hien, T.T. and White, N.J. 1993. Qinghaosu. **Lancet** 341:603-608.
- Hwang, H.H. and Gelvin, S.B. 2004 Plant proteins that interact with VirB2, the *Agrobacterium tumefaciens* pilin protein, mediate plant transformation. **Plant Cell.** 16:3148-3167.
- Jaziri, M., Diallo, B., Vanhaelen, M., Homes, J., Yoshimatsu, K. and Shimomura, K. 1993. Immunodetection of artemisinin in *Artemisia annua* cultivated in hydroponic conditions. **Phytochem.** 33(4):821-826.
- Johnson, D.A. and Cumming, T.F. 2000. Evaluation of mint mutants, hybrids and fertile clones for resistance to *Verticillium dahliae*. **Plant Disease.** 84(3):235-238.
- Jung, M., El-Sohly, H.N. and Mc-Chesney, J.D. 1990. Artemisinic acid: a versatile chiral synthon and bioprecursor to natural products. **Planta Med.** 56:624.

- Kamchonwongpaisan, S. and Meshnick, S.R. 1996. The mode of action of the antimalarial artemisinin and its derivatives. **Gen. Pharmacol.** 27:587-592.
- Kawamoto, H., Sekine, H. and Furuya, T. 1999. Production of artemisinin and related sesquiterpenes in Japanese *Artemisia annua* during a vegetation period. **Planta Med.** 65:88-89.
- Kim, Y., Wyslouzil, B.E., Weathers, P.J. 2001. A comparative study of mist and bubble column reactors in the *in vitro* production of artemisinin. **Plant Cell Rep.** 20:451-455.
- Kirdmanee, C., Kitiya, Y. and Kozai, T. 1995. Effects of CO<sub>2</sub> enrichment and supporting material *in-vitro* on photoautotrophic growth of eucalyptus plantlets *in-vitro* and *ex-vitro*. **In-vitro Cell. Devel. Biol.-plant.** 31(3):144-149.
- Klayman, D.J. 1993. *Artemisia annua*: From weed to respectable antimalarial plant. In, Human Medicinal Agents from Plant, pp. 242-255. Washington: (Eds. A.D.Kinghorn and M.F. Balandrin). American Chemical Society.
- Klayman, D.L., 1985. Qinghausu (Artemisinin): An antimalarial drug from China. **Science.** 228:1048-1055.
- Kohler, M., Haerdi, W., Christen, P. and Veuthey, J.L. 1997. Extraction of artemisinin and artemisinic acid from *Artemisia annua* L. using supercritical carbon dioxide, **J. Chromatogr. A.** 785:353-360.
- Koobkokkrud, T., Chochai, A., Kerdmanee, C. and De-Eknamkul, W. 2007. TLC-densitometric analysis of artemisinin for rapid screening of high producing plantlets of *Artemisia annua*. **Phytochemical Anal.** 18:229-234.
- Koobkokkrud, T. 2002. Potential of artemisinin production in *Artemisia annua* mutants. Thesis of Master's Degree. Chulalongkorn University.



- Kudakasseril, G.J., Lam, L. and Staba, E.J. 1986. Effect of sterol inhibitors on the incorporation of  $^{14}\text{C}$ -isopenyl pyrophosphate into artemisinin by a cell-free system from *Artemisia annua* tissue cultures and plants. **Planta Med.** 53:280-284.
- Li, Z-Q., Liu, Y., Liu, B-Y., Wang H., Ye H-C. and Li, G-F. 2006. Cloning, *E. coli* expression and molecular analysis of amorpha-4,11-diene synthase from a high-yield Strain of *Artemisia annua* L. *J. Integrative Plant Biol.* 48 (12): 1486-1492.
- Lichtenthaler, H. 1999. The 1-deoxy-D-xylulose-5-phosphate pathway of isoprenoid biosynthesis in plants. **Ann. Rev. Plant Physiol Plant Mol. Biol.** 50:47-65.
- Liu, C-Z., Zhou, H-Y. and Zhao, Y. 2007. An effective method for fast determination of artemisinin in *Artemisia annua* L. by high performance liquid chromatography with evaporative light scattering detection. **Analytica Chimica Acta.** 581:298-302.
- Liu, C.-Z., Guo, C., Wang, Y-C., Ouyang, F. 2003. Comparison of various bioreactors on growth and artemisinin biosynthesis of *Artemisia annua* L. shoot cultures. **Process Biochem.** 39:45-49.
- Luo, X., Yeh, H.J.C., Brossi, A., Flippen-Anderson, J.L. and Gilardi, R. 1984. Configurations of antimalarial derived from quinghaosu: Dihydroqinghaosu, artemether, and artesunic acid. **Helvetica Chimica Acta** 67: 1515-1522.
- Melendez, V., Peggins, J.O., Brewer, T.G. and Theoharides, A.D. 1991. Determination of the antimalarial artemether and its deethylated metabolite dihydroartemisinin in plasma by high performance liquid chromatography with reductive electrochemical detection. **J. Pharm. Sci.** 80(2):132-138.

- Merck, P., Beng, M., Bouwmeester, H.J., Posthumus M.A. and Brodelius, P.E. 2000. Molecular cloning, expression, and characterization of amorpha-4,11-diene synthase, a key enzyme of artemisinin biosynthesis in *Artemisia annua* L. **Arch. Biochem. Biophys.** 381(2):173-180.
- Mercke, P., Crock J., Croteau, R. and Brodelius, P.E. 1999. Cloning, expression, and characterization of epi-cedrol synthase, a sesquiterpene cyclase from *Artemisia annua*L. **Arch. Biochem. Biophys.** 369(2):213-222.
- Meshnick, S.R.1996. Is haemmozoin a target for antimalarial drugs. **Annu. Trop. Med. Parasito.** 90(4):367-372.
- Murashige, T. and Skoog, F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. **Physiologia Plantarum.** 15:473-496.
- Nair, M.S.R. and Basile, D.V. 1993. Bioconversion of arteannuin B to artemisinin. **J. Nat. Prod.** 56(9):1559-1566.
- Nelson, D.L. and Cox, M.M. 2000. Lehninger Principles of Biochemistry. 3th ed. New York: Worth Publishers.
- Paniego, N.B. and Giulietti 1994. *Artemisia annua* L.: dedifferentiated and differentiated cultures. **Plant Call, Tiss. Org. Cult.** 36:163-168
- Peng, C.A., Ferreira, J.F.S. and Wood, A. J. 2006. Direct analysis of artemisinin from *Artemisia annua* L. Using high-performance liquid chromatography with evaporative light scattering detector, and gas chromatography with flame ionization detector. **J.Chromatogr. A.** 1133:254-258
- Phillips, R.S. 2001. Current status of malaria and potential for control. **Clin. Microb. Rev.** 14(1):208-226.

- Pornnaree Cheewasakulyong. 2001. Biotechnological application for improvement of antimalarial agent production in qing-hao (*Artemisia annua*). Master's Thesis. Faculty of graduate studies. Mahidol University.
- Praphassorn Sutakavtin. 1999. Studies on the production of the antimalarial drug, artemisinin, in tissue and cell cultures of *Artemisia annua*. Doctoral dissertation. School of Bioscience. Cardiff University.
- Prathanturarug, S., Soonthornchareonnon, N., Chuakul, W., Phaidee, Y. and Saralamp, P. 2003. High-frequency shoot multiplication in *Curcuma longa* L. using thidiazuron. **Plant Cell Rep.** 21:1054–1059.
- Quispe-Condori, S., Sanchez, D., Foglio, M.A., Rosa, P.T.V., Zetzl, C., Brunner, G., Angela, M. and Meireles, A. 2005. Global yield isotherms and kinetic of artemisinin extraction from *Artemisia annua* L. Leaves using supercritical carbon dioxide, **J. Supercrit. Fluids.** 36:40–48.
- Ranasing, A., Sweatlock, J.D. and Cook, R.G. 1993. A rapid screening method for artemisinin and its congeners using MS/MS: search for new analogues in *Artemisia annua*. **J. Nat. Prod.** 56(4):552-563.
- Roth, R.J. and Acton, N. 1987. Isolation of arteannuic acid from *Artemisia annua*. **Planta Med.** 53:501-502.
- Roth, R.J. and Acton, N. 1987. Isolation of epi-deoxyarteannuin B from *Artemisia annua*. **Planta Med.** 576.
- Rucker, G., Mayer, R. and Manns, D. 1986. Isolation of qinghaosen from *Artemisia annua* of European source. **Planta Med.** 245.
- Rucker, G., Mayer, R. and Manns, D. 1987. Alpha-and beta-myrcene hydroperoxide from *Artemisia annua*. **J. Nat. Prod.** 50(2):287-289.

- Sanswan, R.S., Agarwal, K., Luthra, R., Thakur, R.S. and Singh-Sangwan, N. 1993. Biosyntrasformation of arteanuic acid into arteannuin B and artemisinin in *Artemisia annua*. **Phytochem.** 34(5):1301-1302.
- Sipahimalani, A.T., Fulzele, D.P., and Heble M.R. 1991. Rapid method for the detection of artemisinin by gas chromatography. **J. Chromtogr.** 538:452-455.
- Sy, L.K. and Brown, G.D. 2002. The mechanism of the spontaneous autoxidation of dihydroartemisinic acid. **Tetrahedron** 58:897-908.
- Tawfig, N.K., Anderson, L.A., Roberts, M.F., Phillipson, J.D. and Warhurst, D.C. 1989. Antiplasmodial activity of *Artemisia annua*. **Plant Cell Rep.** 8:425-428.
- Tefera, W. and Wannakraioj, S. 2004. A Micropropagation Method for Korarima (*Aframomum corrorima* (Braun) Jansen). **Science Asia** 30:1-7.
- Tefera, W. and Wannakraioj, S. 2004. Micropropagation of Krawan (*Amomum krervanh* Pierre ex Gagnep). **Science Asia** 30:9-15.
- Tu, Y. Y., Ni, M.Y., Zhong, Y.R., Li, L.N., Cui, S.L., Zhang. M.G., Wang, X.Z., Ji. Z., and Liang, X.T. 1982. Studies on the constituents of *Artemisia annua* part II. **Planta Med.** 44:143-145.
- Tzfira, T. and Citovsky, V. 2002. Partners-in-infection: host proteins involved in the transformation of plant cells by *Agrobacterium*. **Trends Cell Biol.** 12:121-129.
- Tzfira, T., Li, J., Lacroix, B. and Citovsky, V. 2004. *Agrobacterium* T-DNA integration: molecules and models. **Trends Genet.** 20:375-383.
- Van Nijlen, T., Brennan, K., Van den Mooter, G., Blaton, N., Kinget, R. and Augustijns, P. 2003. Improvement of the dissolution rate of artemisinin by means of supercritical fluid technology and solid dispersions. **Int. J. Pharm.** 254:173-181.

- Vandenbergh, D.R., Vergauwe, A.N., Montagu, M.V., and Van den Eeckhout, E.G. 1995. Simultaneous determination of artemisinin and its bioprecursors in *Artemisia annua*. **J. Nat. Prod.** 58(5):798-803.
- Vanwiller, S.C., Haynes, R.K., King, G., and Wang, H.J. 1993. An improved method for the isolation of Qinghao. (Artemisinic) acid from *Artemisia annua*. **Planta Med.** 59:562-563.
- Vergunst, A.C., Schrammeijer, B., den Dulk-Ras, A., de Vlaam C.M.T., Regensburg-Tuink, T.J. and Hooykaas, P.J.J. 2000. VirB/D4-dependent protein translocation from *Agrobacterium* into plant cells. **Science** 290:979-982.
- Wallaart, T.E., Pras, N. and Quax W.J. 1999. Seasonal variations of artemisinin and its biosynthetic precursors in tetraploid *Artemisia annua* plants compared with the diploid wild-type. **Planta Med.** 65(8):723-728
- Wallaart, T.E., Bouwmeester, H.J., Hille, J., Popping, L. and Maijers, N.C.V. 2001. Amorpha-4,11-diene synthase: cloning and functional expression of a key enzyme in the biosynthetic pathway of the novel antimalarial drug artemisinin. **Planta.** 212:460-465.
- Wallaart, T.E., Pras, N. and Quax, W.J. 1999a. Isolation and identification of dihydroartemisinic acid hydroperoxide from *Artemisia annua*: a novel biosynthetic precursor of artemisinin. **J. Nat. Prod.** 62:1160-1162.
- Wallaart, T.E., Uden, W.V., Lubberink, G.M., Woerdenbag, H.J., Pras, N. and Quax W.J. 1999b. Isolation and identification of dihydroartemisinic acid from *Artemisia annua* and its possible role in the biosynthesis of artemisinin. **J. Nat. Prod.** 62:430-433.



- Wang, K., Wang, Y., Bao, Y.L., Meng, X.Y., Wu, Y., Li, Y.X. 2006. A method for high-quality RNA extraction from *Panax ginseng* plant tissues. **Progr Biochem. Biophys.** 33(1):95-99.
- Wanwimon Luealon. 2006. Effect of hormones for regeneration of *Artemisia annua* L. and production of artemisinin by hairy root cultures. **Master's Degree Thesis**, Khon Kaen University, Thailand.
- Ward, E. and Barnes, W. 1988. VirD2 protein of *Agrobacterium tumefaciens* very tightly linked to the 50 end of T-strand DNA. **Science** 242:927-930.
- Weathers P.J., Elkholy S. and Wobbe K.K. 2006. Artemisinin: the biosynthesis pathway and its regulation in *Artemisia annua*, a terpenoid-rich species. **In Vitro Cell. Dev. Biol.-Plant** 42:309-317.
- Wheeler, C. and Berkly, S. 2001. Initial lessons from public-private partnerships in drug and vaccine development. **Bulletin of the World Health Organization** 79(8):728-734.
- WHO 2001. Antimalarial drug combination therapy report of a WHO technical consultation, World Health Organization, Geneva
- WHO 2002. Improving access to antimalarial medicines, report of the RBM partnership meeting, 30 September-2 October 2002, Malaria Control Department, World Health Organization, Geneva
- Woerdenbag, H.J., Lugt, C.B., and Pras, N. 1990. *Artemisia annua* L.: a source of novel antimalarial durgs. *Pharmaceutisch Weekblad Scientific Scientific* edition 12(5): 169-181.
- Yagen, B., Pu, Y.M., Yeh, H.J.C. and Ziffer, H. 1994. Tandem silica gel-catalysed rearrangments and subsequent Baeyer-Villiger reactions of artemisinin derivatives. **J. Chem. Soc. Perkin Trans.** 1:843-846.

- Yamey, G. 2002. WHO in 2002 faltering steps towards partnerships. **Gen. Med. J.** 325:1236-1240.
- Yang, S.L., Roberts, M.F., O'Neill, M.J., Bucar, F. and Phillipson, J.D. 1995. Flavonoid and Chromens from *Artemisia annua*. **Phytochem.** 38(1):255-257.
- Zakeri, S, Najafabadi, S.T., Zare, A and Djadid, N.D. 2002. Detection of malaria parasites by nested PCR in south-eastern, Iran: evidence of highly mixed infections in Chahbahar district. **Malaria Journal.** 1:1-6.
- Zhao, S. and Zeng M.Y. 1986. Application of precolumn reaction to high performance liquid chromatography of Qinghausu in animal plasma. **Anal. Chem.** 58:289-292.

## **APPENDICS**

## TLC-Densitometric Analysis of Artemisinin for the Rapid Screening of High-producing Plantlets of *Artemisia annua* L.

THONGCHAI KOOBKOKKRUAD,<sup>1</sup> ARAYA CHOCHAI,<sup>2</sup> CHALERMPOL KERDMANEE<sup>2</sup> and WANCHAI DE-EKNAMKUL<sup>3\*</sup>

<sup>1</sup> Department of Biochemistry, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok, Thailand

<sup>2</sup> Plant Biochemistry and Plant Physio-Biochemistry Laboratory, National Center for Genetic Engineering and Biotechnology, Pathumthani, Thailand

<sup>3</sup> Department of Pharmacognosy, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok, Thailand

Received 7 February 2006; Revised 8 August 2006; Accepted 13 August 2006

**Abstract:** A simple TLC-densitometric technique has been developed for the rapid and accurate analysis of artemisinin in a large number of *Artemisia annua* plantlets cultured *in vitro*. This new analytical method is based on the structural conversion of artemisinin on a silica gel layer by ammonia vapour to form 10-azadeseoxyartemisinin, a chromophore-containing compound ( $\lambda_{\max}$  320 nm) that can be detected by UV-based TLC densitometry. The TLC system was evaluated quantitatively in terms of product stability, precision, accuracy and calibration. Good linearity was obtained in the range of 0.01–0.12  $\mu$ g artemisinin. The technique appeared to be accurate and sensitive as compared with the complicated pre-column reaction-HPLC technique. Among 90 samples of *A. annua* plantlets, the artemisinin content in the leaves appeared to be highly variable, ranging from 0.02 to 0.67% w/w dry weight. These results demonstrate that densitometric TLC can be a cheap and simple technique for the accurate screening of high-artemisinin-producing plants. Copyright © 2007 John Wiley & Sons, Ltd.

**Keywords:** TLC densitometry; quantitative analysis; artemisinin; 10-azadeseoxyartemisinin; *Artemisia annua* L.; Asteraceae.

### INTRODUCTION

Although artemisinin (**1**) absorbs light in the UV region between 210 and 220 nm, its extinction coefficient is poor. As a result, the methodology of standard UV detection is ineffective in the quantitative analysis of artemisinin. For this reason, the detection of artemisinin has been accomplished using pre-column derivatisation to convert artemisinin into a UV-active compound that absorbs with a large extinction coefficient at longer wavelengths that are applicable to HPLC-UV methods (Zhao and Zeng, 1986; Zhao, 1987; Thomas *et al.*, 1992). In addition, HPLC employing various detection methods has been reported for the analysis of artemisinin, giving different levels of detection, including chemiluminescence (detection limit 2.5 ng; Green *et al.*, 1995), reductive electrochemical detection (detection limit 1–10 mg; Melendez *et al.*, 1991; Navaratnam *et al.*, 1995; Sandrenan *et al.*, 1997), thermospray mass spectrometry (detection limit of 0.2 ng; Chi *et al.*, 1991) and evaporative light scattering detection (detection range of 6–60 ng; Avery *et al.*, 1999). Other analytical methods have also been developed, such as GC-MS (Sipahimalani *et al.*, 1991), capillary electrophoresis (D'Hulst *et al.*, 1996; Yu *et al.*,

2004) immuno-detection (Jaziri *et al.*, 1993) and tandem MS (Ranasing *et al.*, 1993). These techniques usually require either sophisticated equipment or complicated procedures for the analysis of artemisinin.

Recently, a simpler technique of TLC densitometry has been developed in order to analyse artemisinin and its derivatives, both on a normal silica-gel layer (Gabriels and Platzier-Vercammen, 2003) and on a reversed phase-layer (Gabriels and Platzier-Vercammen, 2004). For both TLC systems, spots are visualized by derivatisation with acidified 4-methoxybenzaldehyde reagent in methanol:water. Whilst these methods permit the rapid quantification of artemisinin, their sensitivity is relatively low, with detection in the range from 0.5 to 8  $\mu$ g. In the present paper, we report an improved TLC-densitometric method that is much more sensitive (10 ng detection limit) and accurate. This new technique has enabled us to screen for high-artemisinin-producing plantlets from a large number of samples.

### EXPERIMENTAL

**Materials.** Authentic artemisinin was obtained from the Plant Biochemistry and Plant Physiobiochemistry Laboratory, National Center for Genetic Engineering and Biotechnology (BIOTEC), Thailand. Organic solvents were all of reagent grade or better and were supplied either by LAB-SCAN Analytical Sciences (Dublin, Ireland) or Merck (Darmstadt, Germany). Water

\* Correspondence to: W. De-Eknamkul, Department of Pharmacognosy, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok, Thailand.  
 E-mail: dwanchai@chula.ac.th  
 Contract/grant sponsor: BIOTEC; Contract/grant number: 2/2545.

was triple deionised. Silica gel Polygram® SIL G/UV254 TLC plates were purchased from Macherey-Nagel (Düren, Germany). For plant material, *Artemisia annua* L. seeds originating from Hanoi, Vietnam, were surface sterilised and then germinated on MS medium for 2 months before their shoots were excised and irradiated with gamma rays (500 rad dose). The exposed shoots were then transferred to fresh MS medium for micro-propagation. The shoot cultures were maintained by sub-culture every 6 weeks. Those plantlets that survived four sub-culture passages were harvested and their leaves were used for the determination of artemisinin as described below.

**Sample preparation.** Fresh leaves obtained from various irradiated plantlets were dried at 65°C in a hot air oven. Each dried sample was then ground to a fine powder in a grinder supplied with a cool water circulator, and the resulting powder passed through a no. 40 sieve. Powder samples (100 mg) were extracted under reflux with 10 mL hexane for 1 h. After cooling, the extract was filtered and used for the TLC-densitometric analysis.

**TLC-densitometric analysis.** Crude extracts of various *A. annua* samples (10 µL each) were spotted onto a pre-coated silica gel 60 F<sub>254</sub> layer (0.25 mm thickness). Up to 12 samples were applied onto each standard TLC plate, which was then developed using the solvent system hexane:ethyl acetate:acetone (16:1:1) to produce a solvent front height of 8 cm. The plate was dried and exposed for 2 h to saturated ammonia vapour in a closed TLC tank maintained in a hot air oven at 100°C for complete chromophore development of artemisinin. The TLC plate was then air-dried and scanned at 320 nm using a Shimadzu (Kyoto, Japan) TLC densitometer dual wavelength scanner model CS-930.

**HPLC analysis.** For method validation, the same samples were analysed using the well-established HPLC method with pre-column reaction as described previously (Zhao, 1987). The analysis was performed on a Varian HPLC system (Walnut Creek, CA, USA) equipped with a ternary solvent pump model 9010 and a UV detector model 9050 set at 260 nm. Samples were automatically injected using a Varian model 9095 auto-sampler, and data were recorded with the aid of Varian Star software (version 6.0). The separation was achieved on a reverse-phase C<sub>18</sub> column (150 × 4 mm i.d.; 5 µm) using an isocratic solvent system of 0.01 M disodium hydrogen phosphate-sodium dihydrogen phosphate buffer (pH 6.7):methanol (38:62) at a flow rate of 1.5 mL/min.

**Optimization of structural conversion of artemisinin on the silica gel layer.** A 10 µL aliquot of a solution of

standard artemisinin (0.4 µg/mL) was applied onto a pre-coated silica gel 60 F<sub>254</sub> layer (0.25 mm thickness), and the plate (loaded with 10 spots) was developed with a mobile phase of hexane:ethyl acetate:acetone (16:1:1). After drying, the TLC plate was cut into 10 strips of width 2 cm, each with an artemisinin spot at an R<sub>f</sub> value of 0.75. Each strip was then exposed separately to ammonia vapour in a closed TLC tank in a hot-air oven and maintained either under a different temperature (from 30 to 125°C) for 2 h, or for different time intervals (from 20 to 180 min) at 100°C. After cooling, each TLC strip was scanned at 320 nm and the intensity of the artemisinin spot determined. The resulting areas under the artemisinin peaks were plotted against either temperature or time of exposure to ammonia vapour. In addition, the artemisinin spot with the highest intensity was also scanned at wavelengths from 200 to 370 nm in order to obtain a UV-absorption spectrum of the chromophore-containing product.


**Standard curve.** A stock solution containing 1.0 mg/mL artemisinin was diluted to obtain standard solutions with various concentrations of artemisinin ranging from 0.06 to 12 µg/mL, and these were employed in the construction of the calibration curve for artemisinin.

**NMR analyses of the product of ammonia-treated artemisinin.** An aliquot (1 mL) of artemisinin solution (2 µg/mL in methanol) was streaked onto a silica gel layer and the plate was developed using the solvent system described above. Ten TLC plates were used for product preparation. TLC plates were exposed to saturated ammonia vapour at 100°C for 2 h and the artemisinin band (R<sub>f</sub> 0.75) scraped from the plate and eluted with absolute ethanol. The obtained product was further purified by TLC using the solvent system hexane:acetone:methanol (10:3:1), and the component at R<sub>f</sub> 0.45 was eluted with absolute ethanol and dried with nitrogen gas. The purified compound was dissolved in deuteriochloroform and <sup>1</sup>H- and <sup>13</sup>C-NMR spectra were recorded on a Jeol (Tokyo, Japan) JNM-A500 NMR at 500 and 125 MHz, respectively. Chemical shifts were determined in ppm as δ values relative to TMS as internal standard.

## RESULTS AND DISCUSSION

The structural conversion of artemisinin on a silica gel layer following exposure to ammonia vapour was investigated by NMR. The resulting <sup>1</sup>H-NMR [(CDCl<sub>3</sub>) δ (ppm), 2.97 (1H, ddd, J = 14, 7, 4, CH-8), 6.20 (1H, brs, CH-10), 5.06 (1H, d, J = 3, CH-11), 1.40 (3H, s, CH<sub>3</sub>-2), 0.86 (3H, d, J = 7, CH<sub>3</sub>-5), 1.06 (3H, d, J = 7, CH<sub>3</sub>-8)] and <sup>13</sup>C-NMR [(CDCl<sub>3</sub>) δ (ppm), 107.5 (C-2), 34.6 (C-3).






**Figure 1** Proposed structural conversion of artemisinin to 10-azadesoxyartemisinin on a silica gel layer following exposure to ammonia vapour for 2 h at 100°C.


22.6 (C-4), 45.6 (C-4a), 35.3 (C-5), 33.6 (C-6), 24.3 (C-7) 43.5 (C-7a), 32.9 (C-8), 173.6 (C-9), 81.5 (C-11), 82.2 (C-11a), 22.2 (CH<sub>3</sub>-2), 18.5 (CH<sub>3</sub>-5), 11.9 (CH<sub>3</sub>-8)] were very similar to those of 10-azadesoxyartemisinin (**2**) described previously by Torok and Ziffer (1995a, b). It was clear that the characteristic C-10 ester ( $\delta$  171.9) of artemisinin was converted into the C-9 amide ( $\delta$  173.6) characteristic of **2**, with the signal at C-11 of **2** being at lower field than the original C-12 of artemisinin. These data confirmed the conversion of artemisinin to **2** (Torok and Ziffer, 1995a, b). In these previous reports, the structural rearrangement of artemisinin was observed in a reaction mixture containing methanolic ammonia and sulphuric acid:silica gel, which produced a mixture of 11-azaartemisinin (45%) and 10-azadesoxyartemisinin (9%). In our case, however, the latter compound appeared to be the major product (>90%) of rearrangement of artemisinin. This might be due to the different conditions of the reaction. Our reaction took place on a silica gel layer with a spot of artemisinin reacting with ammonia vapour under a temperature of 100°C. It is likely that the silica gel-catalysed reaction of artemisinin involves an intermediate that can react with the ammonia molecule to form the final product of 10-azadesoxyartemisinin as shown in Fig. 1.

After performing a UV scan at different wavelengths on the compound, the resulting UV absorption spectrum showed clearly a single  $\lambda_{\max}$  at 320 nm (Fig. 2). It can be seen that the UV absorption characteristic of 10-azadesoxyartemisinin was completely different from that of artemisinin itself. Thus, it is clear that natural artemisinin with no UV-chromophore in the molecule is converted into a chromophore-containing compound of 10-azadesoxyartemisinin (**2**) under the ammonia:silica gel-catalysed reaction. This allowed us to use the wavelength of 320 nm for artemisinin analysis with this TLC-densitometric method. In order to maximise the conversion of artemisinin to 10-azadesoxyartemisinin, the optimum conditions of temperature and time used for the ammonia exposure of artemisinin spot on the TLC plate were studied. The results demonstrated that the amount of 10-azadesoxyartemisinin produced by the reaction



**Figure 2** (A) UV-absorption spectra of artemisinin and 10-azadesoxyartemisinin obtained by scanning spots on a silica gel layer. (B) A typical TLC-chromatogram (scanned at 320 nm) of standard artemisinin after being converted into 10-azadesoxyartemisinin using the developed TLC process.

increased from room temperature to reach its highest level at 75°C, and remained at that level up to a temperature of 125°C [Fig. 3(A)]. In the time-course study, it was found that the conversion of artemisinin increased continuously from 20 to 120 min [Fig. 3(B)], but there was no further increase of product after this time. These results clearly suggested that the minimal conditions for the complete conversion of artemisinin to 10-azadesoxyartemisinin would involve treatment of the TLC plate with ammonia vapour at 75°C for 2 h. However, for consistency and reproducibility, a temperature of 100°C was employed in the remainder of the investigation.




**Figure 3** The effects of time (A) and temperature (B) of exposure to ammonia vapour on the conversion of artemisinin to 10-azadesoxyartemisinin on a silica-gel layer.

A study of the TLC separation of artemisinin in hexane extracts of *A. annua* leaves was carried out using a number of solvent systems. By varying the ratio of hexane, ethyl acetate and acetone in the mobile phase, it was found that complete separation of artemisinin could be achieved with the solvent system containing these components in the ratio 16:1:1. Figure 4 shows the TLC-densitometric chromatograms of hexane extracts obtained from some leaf samples. These chromatograms were obtained after the developed silica gel layer had been exposed to ammonia vapour and scanned at 320 nm. It can be seen that the position of artemisinin ( $R_f$  0.75) was well defined and separate from other components present in the crude extracts. The product peak showed a symmetrical shape with very low baseline noise. The chromatograms also exhibited different peak heights of artemisinin with different samples, suggesting that the developed TLC-densitometric method would be relatively sensitive for detecting the artemisinin product.

A calibration curve was constructed for artemisinin using the described conditions for TLC separation and UV detection. The resulting standard curve showed

linearity in the range 0.5 to 12  $\mu\text{g/mL}$  of artemisinin concentration (equivalent to 5–120 ng of analyte). The limit of detection (0.5  $\mu\text{g/mL}$  of artemisinin) was determined using the criteria of the lowest detectable amount that would produce a peak three times the peak-to-peak baseline noise. Linear regression analysis of the area under the curve ( $x$ ) vs the concentration of calibration standards ( $y$ ) gave the equation  $y = 3159 + 15,743x$  with an  $r^2 = 0.9938$ .

In order to determine the accuracy and precision of the developed TLC-densitometric method, a comparison was conducted using a method involving pre-column reaction coupled with HPLC-UV detection (Zhao, 1987). It was found that the values of artemisinin content in various leaf samples determined by TLC densitometry were similar to those determined by the HPLC method (Table 1). The values of the relative standard deviations between replicate analyses of the artemisinin contents were found to be within 1.4% ( $n = 3$ ). These values indicated that the developed TLC-densitometric method provided high accuracy and precision for the determination of artemisinin in *A. annua* leaves. In terms of stability, the rearranged product of 10-azadesoxyartemisinin appeared to be



**Figure 5** Variation of artemisinin content in various plantlets of *A. annua* that had been exposed to a 500 rad dose of gamma rays. The values were obtained using the TLC-densitometric method described.

## REFERENCES

- Avery BA, Venkatesh KK, Avery MA. 1999. Rapid determination of artemisinin and related analogues using high-performance liquid chromatography and an evaporative light scattering detector. *J Chromatogr B Biomed Sci Applic* **730**: 71–80.
- Chi HT, Ramu K, Baker JK, Hufford CD, Lee IS, Zeng YL, McChesney JD. 1991. Identification of the *in vivo* metabolites of the anti-malarial arteether by thermospray high-performance liquid chromatography/mass spectrometry. *Biol Mass Spectrom* **20**: 609–628.
- D'Hulst A, Augustijns P, Arens S, Van Parijs L, Colson S, Verbeke N, Kinge R. 1996. Determination of artesunate by capillary electrophoresis with low UV detection and possible applications to analogues. *J Chromatogr Sci* **34**: 272–275.
- Gabriels M, Plaizier-Vercammen J. 2003. Densitometric thin-layer chromatographic determination of artemisinin and its lipophilic derivatives, artemether and arteether. *J Chromatogr Sci* **41**: 359–366.
- Gabriels M, Plaizier-Vercammen J. 2004. Development of a reversed-phase thin-layer chromatographic method for artemisinin and its derivatives. *J Chromatogr Sci* **42**: 341–347.
- Green MD, Mont DL, Todd GD, Capmacchia AC. 1995. Chemiluminescent detection of artemisinin-novel endoperoxide analysis using luminol with out hydrogen peroxide. *J Chromatogr A* **695**: 237–242.
- Jaziri M, Diallo B, Vanhaelen M, Homes J, Yoshimatsu K, Shimomura K. 1993. Immunodetection of artemisinin in *Artemisia annua* cultivated in hydroponic conditions. *Phytochemistry* **33**: 821–826.
- Melendez V, Peggins JO, Brewer TG, Theoharides AD. 1991. Determination of the antimalarial arteether and its de-ethylated metabolite dihydroartemisinin in plasma by high-performance liquid chromatography with reductive electrochemical detection. *J Pharm Sci* **80**: 132–138.
- Navaratnam V, Mansor SM, Chin LK, Mordi MN, Asokan M, Nair NK. 1995. Determination of artemether and dihydroartemisinin in blood plasma by high-performance liquid chromatography for application in clinical pharmacological studies. *J Chromatogr B Biomed Applic* **669**: 289–294.
- Ranasing A, Sweatlock JD, Cook RG. 1993. A rapid screening method for artemisinin and its congeners using MS/MS: search for new analogues in *Artemisia annua*. *J Nat Prod* **56**: 552–563.
- Sandrenan N, Sioufi A, Godbillon J, Netter C, Donker M, van Valkenburg C. 1997. Determination of artemether and its metabolite, dihydroartemisinin, in plasma by high-performance liquid chromatography and electrochemical detection in the reductive mode. *J Chromatogr B Biomed Sci Applic* **691**: 145–153.
- Sipahimalani AT, Fulzele, DP, Heble MR. 1991. Rapid method for the detection of artemisinin by gas chromatography. *J Chromatogr* **538**: 452–455.
- Thomas CG, Ward SA, Edwards G. 1992. Selective determination, in plasma, of artemether and its major metabolite, dihydroartemisinin, by high-performance liquid chromatography with ultraviolet detection. *J Chromatogr* **583**: 131–136.
- Torok DS, Ziffer H. 1995a. Syntheses and antimalarial activities of N-substituted 11-azaartemisinins. *J Med Chem* **38**: 5045–5050.
- Torok DS, Ziffer H. 1995b. Syntheses and reaction of 11-azaartemisinin and derivatives. *Tetrahedron Lett* **36**: 829–832.
- Yu Qiao Cheng, Hong Li Chen, Liu Yin Fan, Xing Guo Chen, Zhi De Hu. 2004. On-line conversion and determination of artemisinin and its kinetic parameters using orthogonal design by coupling of flow injection with capillary electrophoresis. *Anal Chim Acta* **525**: 239–245.
- Zhao SS. 1987. High-performance liquid chromatographic determination of artemisinin (qinghaosu) in human plasma and saliva. *Analyst* **112**: 661–664.
- Zhao SS, Zeng MY. 1986. Application of precolumn reaction to high performance liquid chromatography of Qinghausu in animal plasma. *Anal Chem* **58**: 289–292.

## CURRICULUM VITAE

### 1. PERSONAL

**Name** : THONGCHAI KOOBKOKKRUAD  
**Born** : July 9, 1977  
**Born Place** : Nakhon Ratchasima    **Nationality** : Thai  
**Home address** : 170/1 Moo 11 Ladbou-kauw, Srikien,  
 Nakorn Rachsema 30340 Thailand  
 Tel: 044-325237, Mobile Phone:+668-9617-6352  
 e-mail: skoopy@gmail.com, skoopy@hotmail.com

### 2. EDUCATION

- 2004** : The Ph.D. Program in Biomedical Chemistry, Department of Biochemistry, Faculty of Pharmaceutical Science, Chulalongkorn University, Bangkok, Thailand  
**Thesis Title:** Cloning and expression of *amorpha-4,11-diene synthase* gene in *Artemisia annua*
- 2003** : The Master of Science in Biomedical Chemistry, Department of Biochemistry, Faculty of Pharmaceutical Science, Chulalongkorn University, Bangkok, Thailand  
**Thesis Title:** Potential of Artemisinin Production in *Artemisia annua* Mutants
- 2000** : The Bachelor of Science in Biochemistry, Department of Biochemistry, Faculty of Science, Chulalongkorn University, Bangkok, Thailand
- 1998** : The Diploma of Analytical Chemistry Training, Department of Science service, Ministry of Science, Technology and Environment, Bangkok, Thailand (Affiliated Institute of Chulalongkorn University)

### 3. SCHOLARSHIP

- 2004-2006:** Ph.D. Student Scholarship from The National Center of Biotechnology and Genetic Engineering (BIOTEC) National Science and Technology Development Agency, National Science and Technology Development Agency (NSTDA), Thailand

### 4. AWARD

- 2008:** The Outstanding Student Participating in Extracurricular Activity of Chulalongkorn University.

### 5. PUBLICATION

- 2007:** Thongchai Koobkokkrud, Araya Chochai, Chalernpol Kerdmanee and Wanchai De-Eknamkul. **TLC-Densitometric Analysis of Artemisinin for Rapid Screening of High Producing Plantlets of *Artemisia annua* L.** *Phytochemical Analysis* 18:229-234.

### 6. RESEARCH HIGHLIGHTS

- Plant tissue culture technique, Transfer of *in vitro* plantlets to *ex vitro* plants technige, *Agrobacterium*-Mediated transformation techniqe
- Gene cloning, over-expression of recombinant protein, SDS-PAGE, Western Blotting and other molecular biology techniques.
- Radio-tracer technique for biosynthetic pathway
- Experience with HPLC, GC, GC-MS, TLC-desitometry, TLC-radioscanner, DNA sequencing, ELISA
- Natural product analysis for xanthophylls, alliin, allicin, artemisinin, capcicine, plunotol