

REFERENCES

- Abbott FV, Franklin KB, and Westbrook RF. (1995). The formalin test: scoring properties of the first and second phases of the pain response in rats. **Pain** 60: 91–102.
- Abbott FV, Ocvirk R, Najafee R, et al., (1999). Improving the efficiency of the formalin test. **Pain** 83: 561–569.
- Aghajanian GK. (1995). **Electrophysiology of serotonin receptor subtypes and signal transduction pathways**. Bloom, F.R., Kupfer D J (Eds.), *Psychopharmacology*: pp. 1451–1459. The Fourth Generation of Progress. Raven, New York.
- Aghajanian GK and Marek GJ. (1997). Serotonin induces excitatory postsynaptic potentials in apical dendrites of neocortical pyramidal cells. **Neuropharmacology** 4/5: 589-600.
- Aloisi AM, Albonetti ME, and Carli G. (1995). Behavioural effects of different intensities of formalin pain in rats. **Physiol. Behav.** 58: 603–610.
- Alreja M, Mutalik P, Nayar U, et al., (1984). The formalin test: a tonic pain model in the primate. **Pain** 20: 97–105.
- Araneda R and Andrade R. (1991). 5-hydroxytryptamine₂ and 5-hydroxytryptamine_{1A} receptors mediate opposing responses on membrane excitability in rat association cortex. **Neuroscience** 40: 399–412.

- Arnold L, Keck P, and Welge J. (2000). Antidepressant treatment of fibromyalgia. A meta-analysis and review. **Psychosomatic** 41: 104-113.
- Arnt J, Hyttel J, and Larsen JJ. (1984). The citalopram: 5-HTP-induced head shake syndrome is correlated to 5-HT₂ receptor affinity and also influenced by other transmitters. **Acta Pharmacol. Toxicol.** 55: 363-372.
- Bardin L, Lavarenne J and Eschalier A. (2000) Serotonin receptor subtypes involved in the spinal antinociceptive effect of 5-HT in rats. *Pain* 86:11-18
- Baxter G, Kennett G, Blaney F, et al.(1995). 5-HT₂ receptor subtypes: a family re-united? *Trends Pharmacol. Sci.* 16: 105-110.
- Baxter G. (1996). Novel discriminatory ligands for 5-HT_{2B} receptors. **Behav. Brain Res.** 73: 149-152.
- Belmonte C. (1996) **Signal transduction in nociceptors: general principles.** *Neurobiology of Nociceptors* (Belmonte C and Cervero F eds) pp 243-257. Oxford Univ Press, Oxford.
- Bessou P and Perl ER. (1969). Response of cutaneous sensory units with unmyelinated fibers to noxious stimuli. **J. Neurophysiol.** 32: 1025-1043.
- Bevan S. (1996). **Intracellular messengers and signal transduction in nociceptors.** *Neurobiology of nociceptors* (Belmonte C and Cervero F eds). pp 258-297. Oxford Univ Press, Oxford.

- Blue ME, Yagaloff KA, Mamounas LA, et al., (1988). Correspondence between 5-HT₂ receptors and serotonergic axons in rat neocortex. **Brain Res.** 453: 315–328.
- Boadle-Biber MC. (1993). Regulation of serotonin synthesis. **Prog. Biophys. Molec. Biol.** 60: 1-15.
- Bobker DH. (1994). A slow excitatory postsynaptic potential mediated by 5-HT₂ receptors in nucleus prepositus hypoglossi. **J. Neurosci.** 14: 2428–2434.
- Boess FG and Martin I. (1994). Molecular biology of 5-HT receptors. **Neuropharmacology** 33: 275-317.
- Bonhaus DW, Weinhardt KK, Taylor M, et al., (1997). RS-102221: a novel high affinity and selective, 5-HT_{2C} receptor antagonist. **Neuropharmacology** 36: 621–629.
- Brennan TJ, Vandermeulen EP, and Gebhart GF. (1996). Characterization of a rat model of incisional pain. **Pain** 64: 493–501.
- Broton JG and Sessle BJ. (1988). Reflex excitation of masticatory muscles induced by algescic chemicals applied to the temporomandibular joint of the cat. **Arch. Oral Biol.** 33: 741–747.
- Busatto GF and Kerwin RW, (1997). Perspectives on the role of serotonergic mechanisms in the pharmacology of schizophrenia. **J. Psychopharmacol.** 11: 3–12.
- Butler SH, Godefroy F, Besson JM, et al., (1992). A limited arthritic model for chronic pain studies in the rat. **Pain** 48: 73–81.

- Bourson A, Wanner D, Wyler R, et al., (1996). Pharmacologic evaluation of the discriminative stimulus of metachlorophenylpiperazine. **Pharmacol. Biochem. Behav.** 53: 107–114.
- Bullitt E. (1989). Induction of *c-fos* like protein within the lumbar spinal cord and thalamus of the rat following peripheral stimulation. **Brain Res.** 493: 391-397.
- Burnet PWJ, Eastwood SL, Lacey K, et al. (1995). The distribution of 5HT_{1A} and 5-HT_{2A} receptor mRNA in human brain. **Brain Res.** 676: 157-168.
- Cervero F. (1994). Sensory innervation of the viscera: peripheral basis of visceral pain. **Physiol. Rev.** 74: 95-138.
- Chen K, Yang W, Grimsby J, et al. (1992). The human 5-HT₂ receptor is encoded by a multiple intron-exon gene. **Mol. Brain Res.** 14: 20-26.
- Clavelou P, Dallel R, Orliaguet T, et al., (1995). The orofacial formalin test in rats: effects of different formalin concentrations. **Pain** 62: 295–301.
- Clavelou P, Pajot J, Dallel R, et al., (1989). Application of the formalin test to the study of orofacial pain in the rat. **Neurosci. Lett.** 103: 349–353.
- Coderre TJ, Fundytus ME, McKenna JE, et al., (1993). The formalin test: a validation of the weighted-scores method of behavioural pain rating. **Pain** 54: 43–50.
- Coderre TJ and Wall PD. (1987). Ankle joint urate arthritis (AJUA) in rats: an alternative animal model of arthritis to that produced by Freund's adjuvant. **Pain** 28: 379–393.

- Coggeshall RE and Carlton SM. (1997). Receptor localization in the mammalian dorsal horn and primary afferent neurons. **Brain Res. Rev.** 24: 28–66.
- Coggeshall RE and Carlton SM. (1997). Receptor localization in the mammalian dorsal horn and primary afferents, mast cells, and sympathetic efferents. **J. Neurophysiol.** 62: 48-58.
- Conn PJ and Sanders-Bush E. (1984). Selective 5-HT₂ antagonists inhibit serotonin stimulated phosphatidyl inositol metabolism in cerebral cortex. **Neuropharmacology** 23: 993–996.
- Conn PJ and Sanders-Bush E. (1986). Regulation of serotonin-stimulated phosphoinositide hydrolysis: relation to the serotonin 5-HT₂ binding site. **J. Neurosci.** 6: 3669–3675.
- Curzon G and Green A. (1970). The effect of D-and L-p-chloro-phenylalanine on the metabolism of 5-hydroxytryptamine in brain. **Br. J. Pharmacol.** 39: 635-655.
- Dallel R, Raboisson P, Clavelou P, et al., (1995). Evidence for a peripheral origin of the tonic nociceptive response to subcutaneous formalin. **Pain** 61: 11–16.
- Deecher DC, Wilcox BD, Dave V, et al. (1993). Detection of 5-hydroxytryptamine₂ receptors by radioligand binding, northern blot analysis, and Ca²⁺ responses in rat primary astrocyte cultures. **J. Neurosci. Res.** 35: 246-256.

- Deffenu G, Pegrassi L, and Lumachi B. (1966). The use of bradykinin-induced effects in rats as an assay for analgesic drugs. **J. Pharm. Pharmacol.** 18: 135.
- Dirig DM, Salami A, Rathbun ML, et al., (1997). Characterization of variables defining hindpaw withdrawal latency evoked by radiant thermal stimuli. **Neurosci. Methods** 76: 183–191.
- Done CJ and Sharp T. (1992). Evidence that 5-HT₂ receptor activation decreases noradrenaline release in rat hippocampus in vivo. **Br. J. Pharmacol.** 107: 240–245.
- Done CJ and Sharp T. (1994). Biochemical evidence for the regulation of central noradrenergic activity by 5-HT_{1A} and 5-HT₂ receptors: microdialysis studies in the awake and anaesthetized rat. **Neuropharmacology** 33: 411–421.
- Dougherty PM and Willis WD. (1991). Enhancement of spinothalamic neuron responses to chemical and mechanical stimuli following combined micro-inotophoretic application of N-methyl-D-aspartic acid and substance P. **Pain** 47: 85-93.
- Dougherty PM, Palecek J, Paleckova V, et al., (1992). The role of NMDA and non-NMDA excitatory amino acid receptors in the excitation of primate spinothalamic tract neurons by mechanical, chemical, thermal, and electrical stimuli. **J. Neurosci.** 12: 3025-3041.

- Dougherty PM, Palecek J, Paleckova V, et al., (1994). Neurokinin 1 and 2 antagonists attenuate the responses and NK 1 antagonists prevent the sensitization of primate spinothalamic tract neurons after intradermal capsaicin. **J. Neurophysiol.** 72: 1464-1475.
- Dubuisson D and Dennis SG. (1977). The formalin test: a quantitative study of the analgesic effects of morphine, meperidine and brain-stem stimulation in rats and cats. **Pain** 4: 161-174.
- Duclaux R and Kenshalo DR. (1980). Response characteristics of cutaneous warm receptors in the monkey. **J. Neurophysiol.** 43: 1–15.
- Duman RS, Heninger GR, and Nestler EJ. (1997). A molecular and cellular theory of depression. **Arch. Gen. Psychiatry** 54: 597–606.
- Erspamer V. (1956). Pharmacology of indolealdylamines. **Pharmacol Rev** 6: 425-487.
- Ferrari MD, Odink J, Tapparelli C, et al., (1989). Serotonin metabolism in migraine. **Neurology** 39: 1239-1242.
- Fields HL. (1987). Pain. New York: McGraw-Hill.
- Fiorella D, Helsley S, Rabin RA, et al., (1995). 5-HT_{2C} receptor-mediated phosphoinositide turnover and the stimulus effects of m-chlorophenylpiperazine. **Psychopharmacology** 122: 237–243.
- Fishbain D. (2000). Evidence-based data on pain relief with antidepressants. **Ann. Med.** 32: 305-316.

- Foong FW, Satoh M, and Takagi H. (1982). A newly devised reliable method for evaluating analgesic potencies of drugs on trigeminal pain. **J. Pharmacol. Methods** 7: 271–278.
- Francis PT, Pangalos MN, Pearson RC, et al. (1992). 5-Hydroxytryptamine_{1A} but not 5-hydroxytryptamine₂ receptors are enriched on neocortical pyramidal neurones destroyed by intrastriatal volkensin. **J. Pharmacol. Exp. Ther.** 261: 1237-1281.
- Fuller RW. (1996). Serotonin receptors involved in the regulation of pituitary-adrenocortical function in rats. **Behav. Brain Res.** 73: 215–219.
- Gaddum JH and Picarelli Z. (1957). Two kinds of tryptamine receptor. **Br. J. Pharmacol.** 12: 323-328.
- Galbraith JA, Mrosko BJ, and Myers RR. (1993). A system to measure thermal nociception. **J. Neurosci. Methods** 49: 63–68.
- Geppetti P and Holzer P. (1996). Neurogenic Inflammation. CRC Press, Boca Raton.
- Gilchrist HD, Allard BL, and Simone DA. (1996). Enhanced withdrawal responses to heat and mechanical stimuli following intraplantar injection of capsaicin in rats. **Pain** 67: 179–188.
- Glennon RA and Lucki I. (1988). **Behavioural models of serotonin receptor activation.** Sanders-Bush E. (Ed.), The Serotonin Receptors. pp. 253–293. The Humana, Clifton, NJ.

- Glennon RA. (1990). Do classical hallucinogens act as 5-HT₂ agonists or antagonists? **Neuropsychopharmacology** 3: 509–517.
- Godbout R, Mantz J, Glowinski J, et al., (1991). The novel 5-HT₂ receptor antagonist, RP 62203, selectively blocks serotonergic but not dopaminergic induced inhibition in the rat prefrontal cortex. **Eur. J. Pharmacol.** 204: 97–100.
- Godfrey P, McClue SJ, Young MM, et al., (1988). 5-Hydroxy-tryptamine-stimulated inositol phospholipid hydrolysis in the mouse cortex has pharmacological characteristics compatible with mediation via 5-HT₂ receptors but this response does not reflect altered 5-HT₂ function after 5,7-dihydroxytryptamine lesioning or repeated antidepressant treatments. **J. Neurochem.** 50: 730–738.
- Gold MS, Reichling DB, Shuster MJ, et al., (1996) Hyperalgesic agents increase a tetrodotoxin-resistance Na⁺ current in nociceptors. **Proc. Natl. Acad. Sci. USA** 93: 1108-1112.
- Gorea E, Davenne D, Lanfumey L, et al., (1991). Regulation of noradrenergic coerulean neuronal firing mediated by 5-HT₂ receptors: involvement of the prepositus hypoglossal nucleus. **Neuropharmacology** 30: 1309–1318.
- Green AR and Heal DJ. (1985). The effects of drugs on serotonin-mediated behavioural models. In: Green, A.R. (Ed.), **Neuropharmacology of Serotonin** Oxford University, Oxford, pp. 326–365.

- Grotewiel MS and Sanders-Bush E. (1994). Regulation of serotonin_{2A} receptors in heterologous expression systems. **J. Neurochem.** 63: 1255–1260.
- Grotewiel MS, Chu H, and Sanders-Bush E. (1994). m-Chlorophenylpiperazine and m-trifluoromethylphenylpiperazine are partial agonists at cloned 5-HT_{2A} receptors expressed in fibroblasts. **J. Pharm. Exp. Ther.** 271: 1122–1126.
- Gudelsky GA, Koenig JJ, and Meltzer HY. (1986). Thermoregulatory responses to serotonin (5-HT) receptor stimulation in the rat: evidence for opposing roles of 5-HT₂ and 5-HT_{1A} receptors. **Neuropharmacology** 25: 1307–1313.
- Guzman F, Braun C, Lim RKS, et al., (1964). Narcotic and non-narcotic analgesics which block visceral pain evoked by intra-arterial injection of bradykinin and other algescic agents. **Arch. Int. Pharmacodyn.** 149: 571–588.
- Hargreaves K, Dubner R, Brown F, et al., (1988). A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia. **Pain** 32: 77–88.
- Harris A. (1998). Using c-fos as a Neural Marker of Pain. **Brain Res. Bull** 45: pp1-8.
- Hensel A. (1973). **Cutaneous thermoreceptors.** Handbook of Sensory Physiology: So-matosensory System (A Iggo ed) vol 2, pp 79–110, Springer Verlag, Berlin.

- Hoyer D, Clarke DE, Fozard JR, et al. (1994). International union of pharmacological classification of receptors for 5-hydroxytryptamine (serotonin). **Pharmacol. Rev.** 46: 157-193.
- Huidobro-Toro JP and Harris RA. (1996). Brain lipids that induce sleep are novel modulators of 5-hydroxytryptamine receptors. **Proc. Natl. Acad. Sci. U S A.** 93(15):8078-82.
- Humphrey PPA, Hartig P and Hoyer D. (1993). A proposed new nomenclature for 5-HT receptors. **Trends Pharmacol. Sci.** 14: 233-236.
- Hunskar HS, Fasmer OB, and Hole K. (1985). Formalin test in mice, a useful technique for evaluating mild analgesics. **J. Neurosci. Methods** 14 :69–76.
- Hunskar S and Hole K. (1987). The formalin test in mice: dissociation between inflammatory and non-inflammatory pain. **Pain** 30: 103–114.
- Hunt, S, Pini A and Evan G. (1987). Induction of c-fos-like protein in spinal cord neurons following sensory stimulation. **Nature** 328: 632-634.
- Hwang AS and Wilcox GL. (1986). Intradermal hypertonic saline-induced behavior as a nociceptive test in mice. **Life Sci.** 38: 2389–2396.
- Iadarola MJ, Brady LS, Draisci G, et al., (1988). Enhancement of dynorphin gene expression in spinal cord following experimental inflammation: stimulus specificity, behavioral parameters and opioid receptor binding. **Pain** 35: 313–326.

- Iggo A. (1959). Cutaneous heat and cold receptors with slowly conducting (C) afferent fibres. **Q. J. Exp. Physiol.** 44: 362-370.
- Iriuchijima J and Zotterman Y. (1960). The specificity of afferent cutaneous C fibers in mammals. **Acta. physiol.** 49: 267-278.
- Ito H, Nyberg S, Halldin C, et al. (1998). PET imaging of central 5-HT_{2A} receptors with carbon-11-MDL 100 907. **J. Nucl. Med.** 39: 208-214.
- Jett MF and Michelson S. (1996). The formalin test in rat: validation of an automated system. **Pain** 64: 19–25.
- Jourdan D, Ardid D, Bardin L, et al., (1997). A new automated method of pain scoring in the formalin test in rats. **Pain** 71: 265–270.
- Julius D, Huang KN, Livelli TJ, et al. (1990). The 5-HT₂ receptor defines a family of structurally distinct but functionally conserved serotonin receptors. **Proc. Natl. Acad. Sci. USA** 87: 928-932.
- Kehne JH, Baron BM, Carr AA, et al., (1996). Preclinical characterization of the potential of the putative atypical antipsy-chotic MDL 100907 as a potent 5-HT_{2A} antagonist with a favor-able CNS safety profile. **J. Pharmacol. Exp. Ther.** 277; 968–981.
- Kennett GA, Bright F, Trail B, et al., (1996a). Effects of the 5-HT_{2B} receptor agonist, BW 723C86, on three rat models of anxiety. **Br. J. Pharmacol.** 117: 1443–1448.
- Kennett GA, Wood MD, Bright F, et al., (1996b). In vitro and in vivo profile of SB 206553, a potent 5-HT_{2C}: 5-HT_{2B} receptor antagonist with anxiolytic-like properties. **Br. J. Pharmacol.** 117: 427–434.

- Kennett GA, Wood MD, Bright F, et al., (1997a). SB 242084, a selective and brain penetrant 5-HT_{2C} receptor antagonist. **Neuropharmacology** 36: 609–620.
- Kennett GA, Bright F, Trail B, et al., (1997b). Anxiolytic-like actions of the selective 5-HT₄ receptor antagonists, SB204070A and SB207266A in rats. **Neuropharmacology** 4(5): 707–712.
- Kennett GA, Wood MD, Grewal GS, et al., (1994). In vivo properties of SB 200646A, a 5-HT_{2C:2B} receptor antagonist. **Br. J. Pharmacol.** 111: 797–802.
- Kerr FWL. (1975). Neuroanatomical substract of nociception in the spinal cord. **Pain** 1: 325-356.
- Kjørsvik A, Tjølsen A and Hole K. (2001). Activation of spinal serotonin_{2A/2C} receptors augments nociceptive responses in the rat. **Brain Res.** 910(1-2): 179-181.
- Koek W, Jackson A, and Colpaert FC. (1992). Behavioural pharmacology of antagonists at 5-HT₂:5-HT_{1C} receptors. **Neurosci. Biobehav. Rev.** 16: 95–105.
- Kress M and Reeh PW. (1996). **Chemical excitation and sensitization in nociceptors.** Neurobiology of nociceptors (Belmonte C and Cervero F eds). pp 258-297. Oxford Univ Press, Oxford.

- LaMotte RH, Shain CN Simone DA, et al., (1991). Neurogenic hyperalgesia: psychophysical studies of underlying mechanisms. **Exp. Brain Res.** 49: 35-45.
- Lance JW. (1992). **5-hydroxytryptamine mechanisms in primary headaches.** History of involvement of 5-HT primary headaches. (OlesenJ, Saxena PR eds.) pp 19-28 Raven Press. New York.
- Larivie' re WR and Melzack R. (1996). The bee venom test: a new tonic-pain test. **Pain** 66: 271-277.
- LeBars D, Gozariu M and Cadden SW. (2001). Animal Models of Nociception. **Pharmacol. Rev.** 53(4): 597-652.
- Leonard BE. (1992). Subtypes of serotonin receptors: biochemical changes and pharmacological consequences. **Int. Clin. Psychopharmacol.** 7: 23-31.
- Lewis T and Kellgren JH. (1939). Observations relating to referred pain, visceromotor reflexes and other associated phenomena. **Clin. Sci.** 4:47-71.
- Lewis T. (1927). *The Blood Vessels of the Human Skin and Their Responses.* Shaw & Sons, London.
- Lewis T. (1942). *Pain.* The MacMillan Press, London.
- Leysen JE, Niemegeers CJE, Tollenaar JP, et al.(1978). Serotonergic component of neuroleptic receptors. **Nature** 272(168-171).

- Leysen JE, Janssen PMF, Schotte A, et al., (1993). Interaction of anti-psychotic drugs with neurotransmitter receptor sites in vitro and in vivo in relation to pharmacology and clinical effects-role of 5-HT₂ receptors. **Psychopharmacology** 112: S40–S54.
- Lim RK and Guzman F. (1968). **Manifestations of pain in analgesic evaluation in animals and man**, Pain: Proceedings of the International Symposium on Pain Organised by the Laboratory of Psychophysiology (Soulairac A, Cahn J, and Charpentier J eds) pp 119–152, Academic Press, New York.
- Lin Q, Peng YB and Willis WD. (1996). Possible role of protein kinase C in the sensitization of primate spinothalamic tract neurons. **J. Neurosci.** 16: 3026-3034.
- Lin Q, Peng YB, and Willis WD. (1996a). Antinociception and inhibition from the periaqueductal gray are mediated in part by spinal 5-hydroxytryptamine_{1A} receptors. **J. Pharmacol. Exp. Ther.** 276: 958–967.
- Lin Q, Peng YB and Willis WD. (1996b). Inhibition of primate spinothalamic tract neurons. **J. Neurosci.** 16: 3026-3034.
- Lin Q, Peng YB, and Willis WD. (1996c). Inhibition of primate spinothalamic tract neurons by spinal glycine and GABA is reduced during central sensitization. **J. Neurophysiol.** 76: 1005–1014.
- Lin Q, Peng YB and Willis WD. (1997). Involvement of cGMP in nociceptive processing by and sensitization of spinothalamic neurons in primates. **J. Neurosci.** 17: 3293-3302.

- Lo'pez-Gime'nez JF, Mengod G and Palacios JM. (1997). Selective visualisation of rat brain 5-HT_{2A} receptors by autoradiography with [3H]MDL 100 907. **Naunyn-Schmeideberg's Arch. Pharmacol.** 356: 446-454.
- Lu Y, Pirec V, and Yeomans DC. (1997). Differential antinociceptive effects of spinal opioids on foot withdrawal responses evoked by C fibre or A delta nociceptor activation. **Br. J. Pharmacol.** 121: 1210–1216.
- Lundkvist C, Halldin C, Ginovart N, et al. (1996). [¹¹C]MDL 100907, a radioligand for selective imaging of 5-HT_(2A) receptors with positron emission tomography. **Life Sci.** 158: 187-192.
- Ma QP and Woolf C. (1996). Basal and touch-evoked Fos-like immunoreactivity during experimental inflammation in the rat. **Pain** 67: 307-316.
- Malmberg AB and Yaksh TL. (1992). Antinociceptive actions of spinal anti-inflammatory agents on the formalin test in the rat. **J. Pharmacol. Exp. Ther.** 263: 136–146.
- Marek GJ and Aghajanian GK. (1994). Excitation of interneurons in piriform cortex by 5-hydroxytryptamine: blockade by MDL 100 907, a highly selective 5-HT_{2A} receptor antagonist. **Eur. J. Pharmacol.** 259: 137–141.
- Marek GJ and Aghajanian GK. (1996). LSD and the phenethylamine hallucinogen DOI are potent partial agonists at 5-HT_{2A} receptors on interneurons in rat piriform cortex. **J. Pharmacol. Exp. Ther.** 278: 1373–1382.

- McCall WD, Tanner KD, and Levine JD. (1996). Formalin induces biphasic activity in C-fibers in the rat. **Neurosci. Lett.** 208: 45–48.
- McCormack K. (1994). Non-steroidal anti-inflammatory drugs and spinal nociceptive processing. **Pain** 49: 9-43.
- Meller R, Smith S, Harrison PJ, et al. (1997). 5-HT_{2A} receptor induced release of glutamate from C6 glioma cells. **The Pharmacologist** 39: 16.
- Mendell LM. (1966). Physiological properties of unmyelinated fiber projection to the spinal cord. **Exp. Neurol.** 16: 316-332.
- Mengod G, Nguyen H, Le H, et al. (1990a). The distribution and cellular localization of the serotonin_{1C} receptor mRNA in the rodent brain examined by in situ hybridization histochemistry. Comparison with receptor binding distribution. **Neuroscience** 35: 577-591.
- Mengod G, Pompeiano M, Martinez Mir MI, et al. (1990b). Localization of the mRNA for the 5-HT₂ receptor by in situ hybridization histochemistry. Correlation with the distribution of receptor sites. **Brain Res.** 524: 139-143.
- Mense S. (1993). Nociception from skeletal muscle in relation to clinical muscle pain. **Pain** 54: 241-289.
- Meyer RA, Campbell JN, and Raja SN. (1994). **Peripheral neural mechanisms of nociception.** Textbook of Pain (Wall PD and Melzack R eds) pp 13–44, Churchill Livingstone, Edinburgh.

- Meyer RA and Campbell JN. (1981) Myelinated nociceptive afferents account for the hyperalgesia that follows a burn to the hand. **Science** 213: 1527-1529.
- Millan MJ, Dekeyne A, and Gobert A. (1998). Serotonin (5-HT)_{2C} receptors tonically inhibit dopamine (DA) and noradrenaline (NA), but not 5-HT, release in the frontal cortex *in vivo*. **Neuropharmacology** 37: 953–955.
- Millan MJ, Girardon S, and Bervoets K. (1997). 8-OH-DPAT-induced spontaneous tail-flicks in the rat are facilitated by the selective serotonin (5-HT)_{2C} agonist, RO 60-0175: blockade of its actions by the novel 5-HT_{2C} receptor antagonist SB 206 553. **Neuropharmacology** 36: 743–745.
- Millan, M J. (2002). Descending control of pain. **Progress in Neurobiology** 66(6): 355-474.
- Minami K, Minami M and Harris RA. (1997). Inhibition of 5-hydroxytryptamine type 2A receptor-induced currents by n-alcohols and anesthetics. **J. Pharmacol. Exp. Ther.** 281(3):1136-43.
- Moldofsky H. (1982). Rheumatic pain modulation syndrome: The interrelationships between sleep, central nervous system serotonin, and pain. **Adv. Neurol** 33: 51-5.
- Morilak DA, Garlow A, and Ciaranello R.D. (1993). Immunocytochemical localization and description of neurons expressing serotonin₂ receptors in the rat brain. **Neuroscience** 54: 701-717.

- Morilak DA, Somogyi P, Lujan-Miras R, et al., (1994). Neurons expressing 5-HT receptors in the rat brain: neurochemical identification of cell types by immunocytochemistry, **Neuropsychopharmacology** 11:157–166.
- Murray CW, Porreca F, and Cowan A. (1988). Methodological refinements to the mouse paw formalin test. **J. Pharmacol. Methods** 20: 175–186.
- Müller J. (1840-2). **Elements of Physiology**. (Translated from German, with notes by Baly W. 2nd ed., Vol.2), London. Original German: 1838: *Handbuch der Physiologie des Menschen*, 2nd ed Coblenz.
- Ness TJ and Gebhart GF. (1990). Visceral pain: a review of experimental studies. **Pain** 41: 167-234.
- North RA and Uchimura N. (1989). 5-Hydroxytryptamine acts at 5-HT₂ receptors to decrease potassium conductance in rat nucleus accumbens neurones. **J. Physiol.** 417: 1–12.
- Obata H, Saito S, Sakaki M, et al., (2001). Antiallodynic effect of intrathecally administered 5-HT₂ receptor agonist in rats with nerve ligation. **Pain** 90: 173-179.
- Ochao J and Torebjörk E. (1989). Sensations evoked by intraneural microstimulation of C nociceptors fibres in human skin nerve. **J. Physiol.** 415: 583-599.
- Okamoto K, Imbea H, Morikawaa Y, et al., (2002). 5-HT_{2A} receptor subtype in the peripheral branch of sensory fibers is involved in the potentiation of inflammatory pain in rats. **Pain** 99(1-2): 133-143.

- Okuda K, Nakahama H, Miyakawa H, et al., (1984). Arthritis induced in cat by sodium urate: a possible animal model for tonic pain. **Pain** 18: 287–297.
- Otsuki T, Nakahama I, Niizuma H, et al., (1986). Evaluation of the analgesic effects of capsaicin using a new rat model for tonic pain. **Brain Res.** 365: 235–240.
- Pazos A, Palacios J (1985). Quantitative autoradiographic mapping of serotonin receptors in the rat brain. I. Serotonin-1 receptors. **Brain Res.** 346: 205-230.
- Pazos A, Probst A, Palacios JM. (1987). Serotonin receptors in the human brain-IV. Autoradiographic mapping of serotonin-2 receptors. **Neuroscience** 21: 123-139.
- Perkins MN, Campbell E, and Dray A. (1993). Antinociceptive activity of the brady-kinin B1 and B2 receptor antagonists, des-Arg9,(Leu8)-BK and HOE 140, in two models of persistent hyperalgesia in the rat. **Pain** 53: 191–197.
- Perl ER. (1971). Is pain a specific sensation? **J. psychiat. Res.** 8: 273-287.
- Peroutka SJ and Snyder S. (1979). Multiple serotonin receptors: differential binding of [3H]5-hydroxytryptamine, [3H]lysergic acid diethylamide and [3H]spiroperidol. **Mol. Pharmacol.** 16: 687-699.
- Peroutka SJ. (1993). 5-hydroxytryptamine receptor subtypes and the pharmacology of migraine. **Neurology** 43: 34-38.

- Presley RW, Menétrey D, Levine JD, et al., (1990). Systemic morphine suppresses noxious stimulus-evoked Fos protein-like immunoreactivity in the rat spinal cord. **J. Neurosci** 10: 323-335.
- Price DD. (1988). Psychological and neural mechanisms of pain. Raven Press, New York.
- Pritchett DB, Bach AWJ, Wozny M, et al. (1988). Structure and functional expression of a cloned rat serotonin 5-HT₂ receptor. **EMBO J.** 7: 4135-4140.
- Puig S and Sorkin LS. (1996). Formalin-evoked activity in identified primary afferent fibers: systemic lidocaine suppresses phase-2 activity. **Pain** 64: 345–355.
- Randall LO and Selitto JJ. (1957). A method for measurement of analgesic activity on inflamed tissue. **Arch. Int. Pharmacodyn. Ther.** 111:409–419.
- Rapport MM, Green AA and Page IH. (1948). Serum vasoconstrictor (serotonin). IV. Isolation and characterization. **J. Biol. Chem.** 176: 222-225.
- Rosland JH, Tjølsen JH, Mæhle B, et al., (1990). The formalin test in mice: effects of formalin concentration. **Pain** 42: 235–242.
- Roth BL and Ciaranello RD. (1991). Chronic mianserin treatment decreases 5-HT₂ receptor binding without altering 5-HT₂ receptor mRNA levels. **Eur. J. Pharmacol.** 207: 169–172.

- Russell I, Vaeroy H, Javors M, et al., (1992). Cerebrospinal fluid biogenic amine metabolites in fibromyalgia/fibrositis syndrome and rheumatoid arthritis. **Arthritis Rheum** 19: 104-109.
- Russell I. (1996). Neurochemical pathogenesis of fibromyalgia syndrome. *J. Musculoskeletal Pain* 4: 61-69.
- Saika MD, Tokunaga A and Senba E. (1997). Intradermal 5-HT induces Fos expression in rat dorsal horn neurons not via 5-HT₃ but via 5-HT₂ receptors. *Neurosci. Res.* 29:143–149.
- Sakurada T, Katsumata K, Tan-No K, et al., (1992). The capsaicin test in mice for evaluating tachykinin antagonists in the spinal cord. **Neuropharmacology** 31: 1279–1285.
- Saltzman AG, Morse B, Whitman MM. et al. (1991). Cloning of the human serotonin 5-HT₂ and 5-HT_{1C} receptor subtypes. **Biochem. Biophys. Res. Comm.** 181: 1469-1478.
- Sanders-Bush E and Canton H. (1995). **Serotonin receptors: signal transduction pathways.** Bloom, F.E., Kupfer, D.J. (Eds.), *Psychopharmacology The Fourth Generation of Progress.* pp. 431–442. Raven, New York.
- Sanders-Bush E. (1990). Adaptive regulation of central serotonin receptors linked to phosphoinositide hydrolysis. **Neuropsychopharmacology** 3: 411–416.

- Sanders-Bush E, Burris KD, and Knoth K. (1988). Lysergic acid diethylamide and 2,5-dimethoxy-4-methylamphetamine are partial agonists at serotonin receptors linked to phosphoinositide hydrolysis. **J. Pharmacol. Exp. Ther.** 246: 924–928.
- Sasaki M, Ishizaki K, Obata H, et al., (2001). Effects of 5-HT₂ and 5-HT₃ receptors on the modulation of nociceptive transmission in rat spinal cord according to the formalin test. *Eur. J. Pharmacol.* 424:45–52.
- Saudou F and Hen R. (1994). 5-Hydroxytryptamine receptor subtypes in vertebrates and invertebrates. **Neurochem. Int.** 25: 503-532.
- Shiaible HG and Shimidt RF. (1985) Effects of an experimental arthritis on the sensory properties of fine articular afferents units. **J. Neurophysiol.** 54: 1109-1122.
- Schible HG and Grubb BD. (1993). Afferent and spinal mechanisms of joint pain. **Pain** 55: 5-54.
- Schmidt RF, Schable HG, Messlinger K, et al., (1994). **Silent and active nociceptors: structure, functions, and clinical implications.** Progress in Pain Research and Management (Gebhart FG, Hammond DL and Jensen TS eds) Vol 2. pp 213-250. IASP Press, Seattle.
- Schwarz M, Spaeth M, Muller H, Pongratz D, Bondy B, Ackenheil M. et al., (1999). Relationship of substance P, 5-hydroxyindoleacetic acid and tryptophan in serum of fibromyalgia patients. **Neurosci. Lett.** 256: 196-198.

- Sheldon PW and Aghajanian G. (1991). Excitatory responses to serotonin (5-HT) in neurons of the rat piriform cortex: evidence for mediation by 5-HT_{1C} receptors in pyramidal cells and 5-HT₂ receptors in interneurons. **Synapse** 9: 208-218.
- Shibata M, Ohkubo T, Takahashi H, et al., (1989). Modified formalin test: characteristic biphasic pain response. **Pain** 38: 47–352.
- Sicuteri F. (1992). Headache as possible expression of deficiency of brain 5-hydroxytryptamine (cerebral denervation supersensitivity). **Headache** 12: 69-72.
- Sigiyo S, Yonehara N, Appenteng K, et al., (2001). Effects of intrathecal c-fos antisense oligodeoxynucleotide on adjuvant-induced thermal hyperalgesia. **Exp. Brain. Res.** 140(2):198-205.
- Sjoerdsma A, et al., (1970). Serotonin now: clinical implications of inhibiting its synthesis with parachlorophenylalanine. **Ann. Intern Med** 73: 607-29.
- Simone DA, Sorkin LS, Oh U, et al., (1991). Neurogenic hyperalgesia: central neural correlates in responses of spinothalamic tract neurons. **J. Neurophysiol.** 66: 228-246.
- Sluka KA and Willis WD. (1997). The effect of G-protein and protein kinase inhibitors on the behavioral responses of rat to intradermal injection of capsaicin. **Pain** 71: 165-178.

- Sorensen SM, Kehne JH, and Fadayel GM, et al., (1993). Character-ization of the 5-HT₂ receptor antagonist MDL 100 907 as a putative atypical antipsychotic: behavioral, electrophysiological and neurochemical studies. **J. Pharmacol. Exp. Ther.** 266: 684–691.
- Srikiatkachorn A. (2002). Chronic Daily Headache: A Scientist's Perspective. **Headache** 42(6): 532-537.
- Srikiatkachorn A, Anantasetakul T, Maneesri S, Phansuwan-Pujito P (2000). Hypo-serotonin-induced nitric oxide supersensitivity in the cerebral microcirculation. *Headache* 40: 267-275.
- Srikiatkachorn A, Puangniyom S, Govirrapong P. (1998). Plasticity of 5-HT_{2A} serotonin receptor in patients with analgesic-induce transformed migraine. *Headache* 38: 543-539.
- Srikiatkachorn A, Tarasub N and Govitrapong P. (1999). Acetaminophen-induced antinociception via central 5-HT_{2A} receptors. **Neurochem. Int.**34:491-498
- Srikiatkachorn A, Suwattanasophon C, Ruangpattanatawee A, et al., (2002).5-HT_{2A} receptor activation and nitric oxide synthesis: a possible mechanism determining migraine attacks. **Headache** 42:566-74.
- Srikiatkachorn A, Tarasub N and Govitrapong P. (2000). Effect of chronic analgesic exposure on the central serotonin system: a possible mechanism of analgesic abuse headache. *Headache* 40: 343-350.

- Stam NJ, Vanhuizen F, Vanalebeek C, et al. (1992). Genomic organization, coding sequence and functional expression of human 5-HT₂ and 5-HT_{1A} receptor genes. **Eur. J. Pharmacol.** 227: 153-162.
- Stratz T, et al., (1993). Serotonin concentration in serum of patients with generalised tendomyopathy (fibromyalgia) and chronic polyarthritis. **Med.Klin** 88: 458-462.
- Sufka KJ, Watson GS, Nothdurft RE, et al., (1998). Scoring the mouse formalin test: validation study. **Eur. J. Pain** 2: 351–358.
- Swainston Harrison T and Perry CM. (2004). Aripiprazole: a review of its use in schizophrenia and schizoaffective disorder. **Drugs** 64(15):1715-36.
- Taylor BK, Peterson MA, and Basbaum AI. (1995). Persistent cardiovascular and behavioral nociceptive responses to subcutaneous formalin require peripheral nerve input. **J Neurosci** 15: 7575–7584.
- Taylor BK, Peterson MA, and Basbaum AI. (1997). Early nociceptive events influence the temporal profile, but not the magnitude, of the tonic response to subcutaneous formalin: effects with remifentanyl. **J. Pharmacol. Exp. Ther.** 280: 876–883.
- Teiger DG. (1976). A test for antinociceptive activity of narcotic and narcotic antagonist analgesics in the guinea pig. **J. Pharmacol. Exp. Ther.** 197: 311–316.

- Tjølsen A and Hole K. (1992). The effect of morphine on core and skin temperature in rats. **Neuroreport** 3: 512–514.
- Tokunaga A, Saika M and Senba E. (1998). 5-HT_{2A} receptor subtype is involved in the thermal hyperalgesic mechanism of serotonin in the periphery. **Pain** 76:349–355.
- Tominaga M, Caterina MJ, Malmberg AB, et al., (1998) The cloned capsaicin receptor integrates multiple pain-producing stimuli. **Neuron** 21: 531-543.
- Torebjörk HE and Hallin RG. (1973). Perceptual changes accompanying controlled preferential blocking of A and C fibre responses in intact human skin nerves. **Exp. Brain. Res.** 26: 321-332.
- Torebjörk HE. (1974). Afferent C units responding to mechanical, thermal and chemical stimuli in non-glabrous skin. **Acta. Physiol. Scand.** 92: 531-390.
- Tonussi CR and Ferreira SH. (1992). Rat knee-joint carrageenin incapacitation test: an objective screen for central and peripheral analgesics. **Pain** 48: 421–427.
- Treede RD, Meyer RA, Raja SN, et al., (1995) Evidence for two different heat transduction mechanisms in nociceptive primary afferents innervating monkey skin. **J. Physiol. (Lond)** 483: 747–758.
- Van de Kar LD, Rittenhouse PA, Li Q, et al., (1996). Serotonergic regulation of renin and prolactin secretion. **Behav. Brain Res.** 73: 203–208.

- Van Hees J and Gybels JM. (1981). C nociceptors activity in human nerve during painful and non painful skin stimulation. **J. neurol. Neurosurg. Psychiat.** 44: 600-607.
- Watson GS, Sufka KJ, and Coderre TJ. (1997). Optimal scoring strategies and weights for the formalin test in rats. **Pain** 70: 53–58.
- Wheeler-Aceto H and Cowan A. (1991). Standardization of the rat paw formalin test for the evaluation of analgesics. **Psychopharmacology (Berl)** 104: 35–44.
- Wheeler-Aceto H, Porreca F, and Cowan A. (1990). The rat paw formalin test: comparison of noxious agents. **Pain** 40: 229–238.
- Willis WD and Coggeshall RE. (1991). **Sensory Mechanisms of the SpinalCord**. 2nd Edition. pp. 595. Plenum Press, New York.
- Willis W D. (1994). Central plastic responses to pain. In: Gebhart, G.F., Hammond, D.L., Jensen, T.S. (Eds.), *Proceeding of the Seventh World Congress on Pain*, Vol. 2. IASP Press, Seattle, pp. 301–324.
- Willis WD. (1985). **The Pain System**. The neural Basis of Nociceptive Transmission in the Mammalian Nervous System. Karger, Basel.
- Willis WD. (1994). **Central plastic responses to pain**. *Progress in Pain Research and Management* (Gebhart FG, Hammond DL and Jensen TS eds) Vol 2. pp 301-324. IASP Press, Seattle.
- Wright DE, Seroogy K, Lundgren KH, et al. (1995). Comparative localization of serotonin1A, 1C, and 2 receptor subtype mRNAs in rat brain. **J. Comp. Neurol.** 351: 357-373.

- Xie H, Ma F, Zhang Y-Q, Guo X, Ji G-C, Wu G-C. (2002). Expression of 5-HT_{2A} receptor mRNA in some nuclei of brain stem enhanced in monoarthritic rats. **Brain Res.** 954:94-99.
- Yashpal K and Coderre TJ. (1998). Influence of formalin concentration on the antinociceptive effects of anti-inflammatory drugs in the formalin test in rats: separate mechanisms underlying the nociceptive effects of low- and high-concentration formalin. **Eur. J. Pain** 2: 63–68.
- Yeomans DC and Proudfit HK. (1994). Characterization of the foot withdrawal re-sponse to noxious radiant heat in the rat. **Pain** 59: 85–94.
- Yeomans DC and Proudfit HK. (1996). Nociceptive responses to high and low rates of noxious cutaneous heating are mediated by different nociceptors in the rat: electrophysiological evidence. **Pain** 68: 141–150.
- Yu L, Nguyen H, Le H, et al. (1991). The mouse 5-HT_{1C} receptor contains eight hydrophobic domains and is X-linked. **Mol. Brain Res.** 11: 143–149.
- Yu XM, Sessle BJ, Haas DA, et al., (1996). Involvement of NMDA receptor mechanisms in jaw electromyographic activity and plasma extravasation induced by inflammatory irritant application to temporomandibular joint region of rats. **Pain** 68: 169–178.

- Yu XM, Sessle BJ, Vernon H, et al., (1994). Administration of opiate antagonist naloxone induces recurrence of increased jaw muscle activities related to inflammatory irritant application to rat temporomandibular joint region. **J. Neurophysiol.** 72: 1430–1433.
- Yu XM, Sessle BJ, Vernon H, et al., (1995). Effects of inflammatory irritant application to the rat temporomandibular joint on jaw and neck muscle activity **Pain** 60: 143–149.
- Zahn PK, Gysbers D, and Brennan TJ. (1997). Effect of systemic and intrathecal morphine in a rat model of postoperative pain. **Anesthesiology** 86: 1066–1077.
- Zhang Y-Q, Guo X, Ji G-C and Wu G-C. (2001) Expression of 5-HT_{2A} receptor mRNA in rat spinal dorsal horn and some nuclei of brainstem after peripheral tissue inflammation. **Brain Res.** 900:146-151.
- Zimmermann M. Ethical guidelines for investigations of experimental pain in conscious animals. **Pain** 1983;16:109-110

- Xie H, Ma F, Zhang Y-Q, Guo X, Ji G-C, Wu G-C. (2002). Expression of 5-HT_{2A} receptor mRNA in some nuclei of brain stem enhanced in monoarthritic rats. **Brain Res.** 954:94-99.
- Yashpal K and Coderre TJ. (1998). Influence of formalin concentration on the antinociceptive effects of anti-inflammatory drugs in the formalin test in rats: separate mechanisms underlying the nociceptive effects of low- and high-concentration formalin. **Eur. J. Pain** 2: 63–68.
- Yeomans DC and Proudfit HK. (1994). Characterization of the foot withdrawal re-sponse to noxious radiant heat in the rat. **Pain** 59: 85–94.
- Yeomans DC and Proudfit HK. (1996). Nociceptive responses to high and low rates of noxious cutaneous heating are mediated by different nociceptors in the rat: electrophysiological evidence. **Pain** 68: 141–150.
- Yu L, Nguyen H, Le H, et al. (1991). The mouse 5-HT_{1C} receptor contains eight hydrophobic domains and is X-linked. **Mol. Brain Res.** 11: 143–149.
- Yu XM, Sessle BJ, Haas DA, et al., (1996). Involvement of NMDA receptor mechanisms in jaw electromyographic activity and plasma extravasation induced by inflammatory irritant application to temporomandibular joint region of rats. **Pain** 68: 169–178.

- Yu XM, Sessle BJ, Vernon H, et al., (1994). Administration of opiate antagonist naloxone induces recurrence of increased jaw muscle activities related to inflammatory irritant application to rat temporomandibular joint region. **J. Neurophysiol.** 72: 1430–1433.
- Yu XM, Sessle BJ, Vernon H, et al., (1995). Effects of inflammatory irritant application to the rat temporomandibular joint on jaw and neck muscle activity **Pain** 60: 143–149.
- Zahn PK, Gysbers D, and Brennan TJ. (1997). Effect of systemic and intrathecal morphine in a rat model of postoperative pain. **Anesthesiology** 86: 1066–1077.
- Zhang Y-Q, Guo X, Ji G-C and Wu G-C. (2001) Expression of 5-HT_{2A} receptor mRNA in rat spinal dorsal horn and some nuclei of brainstem after peripheral tissue inflammation. **Brain Res.** 900:146-151.
- Zimmermann M. Ethical guidelines for investigations of experimental pain in conscious animals. **Pain** 1983;16:109-110

BIOGRAPHY

Name	Mr. Maroot Kaewwong
Date of Birth	26 th November 1980
Instruction attended	Kasetsart University (1998-2001) Bachelor of Science (Biology)