

## REFERENCES

- [1] Samrejtüksing, N. Preparation of mixed-mode monolithic capillary column with octadecyl and aminopropyl groups for liquid chromatography.
- [2] Kimata, K., et al. Chromatographic characterization of silica C18 packing materials. correlation between a preparation method and retention behavior of stationary phase. Journal of Chromatographic Science 27(12) (1989): 721-728.
- [3] HPLC Columns, Theory, Technology, and Practice. Instrumentation Science & Technology 26(4) (1998): 439-440.
- [4] MacNair, J.E., Lewis, K.C., and Jorgenson, J.W. Ultrahigh-pressure reversed-phase liquid chromatography in packed capillary columns. Analytical Chemistry 69(6) (1997): 983-989.
- [5] Yan, L., et al. Hybrid organic-inorganic monolithic stationary phase for acidic compounds separation by capillary electrochromatography. Journal of Chromatography A 1046(1-2) (2004): 255-261.
- [6] Svec, F. and Huber, C.G. Monolithic materials: promises, challenges, achievements. Analytical Chemistry 78(7) (2006): 2100-2107.
- [7] Tanaka, N., Kobayashi, H., Nakanishi, K., Minakuchi, H., and Ishizuka, N. Peer reviewed: monolithic LC columns. Analytical Chemistry 73(15) (2001): 420 A-429 A.
- [8] Cabrera, K. Applications of silica-based monolithic HPLC columns. Journal of Separation Science 27(10-11) (2004): 843-852.
- [9] Ding, G., Da, Z., Yuan, R., and Bao, J.J. Reversed-phase and weak anion-exchange mixed-mode silica-based monolithic column for capillary electrochromatography. ELECTROPHORESIS 27(17) (2006): 3363-3372.
- [10] Ye, F., Wang, S., and Zhao, S. Preparation and characterization of mixed-mode monolithic silica column for capillary electrochromatography. Journal of Chromatography A 1216(51) (2009): 8845-8850.
- [11] Núñez, O., Nakanishi, K., and Tanaka, N. Preparation of monolithic silica columns for high-performance liquid chromatography. Journal of Chromatography A 1191(1-2) (2008): 231-252.
- [12] Minakuchi, H., Nakanishi, K., Soga, N., Ishizuka, N., and Tanaka, N. Effect of domain size on the performance of octadecylsilylated continuous porous

- silica columns in reversed-phase liquid chromatography. Journal of Chromatography A 797(1–2) (1998): 121-131.
- [13] Tanaka, N., et al. Monolithic silica columns for HPLC, micro-HPLC, and CEC. Journal of High Resolution Chromatography 23(1) (2000): 111-116.
- [14] Ishizuka, N., et al. Performance of a monolithic silica column in a capillary under pressure-driven and electrodriven conditions. Analytical Chemistry 72(6) (2000): 1275-1280.
- [15] Desmet, G. and Eeltink, S. Fundamentals for LC miniaturization. Analytical Chemistry 85(2) (2012): 543-556.
- [16] Hjertén, S., Liao, J.-L., and Zhang, R. High-performance liquid chromatography on continuous polymer beds. Journal of Chromatography A 473(0) (1989): 273-275.
- [17] Wang, Q.C., Svec, F., and Fréchet, J.M.J. Macroporous polymeric stationary-phase rod as continuous separation medium for reversed-phase chromatography. Analytical Chemistry 65(17) (1993): 2243-2248.
- [18] Sýkora, D., Svec, F., and Fréchet, J.M.J. Separation of oligonucleotides on novel monolithic columns with ion-exchange functional surfaces. Journal of Chromatography A 852(1) (1999): 297-304.
- [19] Minakuchi, H., Nakanishi, K., Soga, N., Ishizuka, N., and Tanaka, N. Octadecylsilylated porous silica rods as separation media for reversed-phase liquid chromatography. Analytical Chemistry 68(19) (1996): 3498-3501.
- [20] Allen, D. and El Rassi, Z. Capillary electrochromatography with monolithic silica columns: III. Preparation of hydrophilic silica monoliths having surface-bound cyano groups: chromatographic characterization and application to the separation of carbohydrates, nucleosides, nucleic acid bases and other neutral polar species. Journal of Chromatography A 1029(1–2) (2004): 239-247.
- [21] Ye, F., Xie, Z., Wu, X., Lin, X., and Chen, G. Separation of polar and basic compounds in hydrophilic interaction pressurized CEC using diethylenetriaminopropyl silica monolithic columns. Chromatographia 64(5-6) (2006): 267-272.
- [22] Tanaka, H., Kamada, M., Nyudo, M., and Ohira, M. Development of a novel octadecyl-bonded silica column and evaluation of its reliability in chromatographic analysis. Journal of Chromatography A 762(1–2) (1997): 89-96.
- [23] Roux, R., Abi Jaoudé, M., and Demesmay, C. Improvement of chromatographic performances of in-situ synthesized hybrid C8 silica monoliths by reduction of

- structural radial heterogeneities. Journal of Chromatography A 1216(18) (2009): 3857-3863.
- [24] Scherer, B. and Steiner, F. Application of hydrophobic anion-exchange phases in capillary electrochromatography. Journal of Chromatography A 924(1-2) (2001): 197-209.
- [25] Jaafar, J., Watanabe, Y., Ikegami, T., Miyamoto, K., and Tanaka, N. Anion exchange silica monolith for capillary liquid chromatography. Analytical and Bioanalytical Chemistry 391(7) (2008): 2551-2556.
- [26] Zheng, M.-M., Ruan, G.-D., and Feng, Y.-Q. Hybrid organic-inorganic silica monolith with hydrophobic/strong cation-exchange functional groups as a sorbent for micro-solid phase extraction. Journal of Chromatography A 1216(45) (2009): 7739-7746.
- [27] Preinerstorfer, B., Lubda, D., Lindner, W., and Lämmerhofer, M. Monolithic silica-based capillary column with strong chiral cation-exchange type surface modification for enantioselective non-aqueous capillary electrochromatography. Journal of Chromatography A 1106(1-2) (2006): 94-105.
- [28] Yan, L.-J., et al. Octyl-functionalized hybrid silica monolithic column for reversed-phase capillary electrochromatography. Journal of Chromatography A 1121(1) (2006): 92-98.
- [29] Nogueira, R., et al. Silica-based monolithic columns with mixed-mode reversed-phase/weak anion-exchange selectivity principle for high-performance liquid chromatography. Journal of Separation Science 29(7) (2006): 966-978.
- [30] Ye, F., Li, S., and Zhao, S. Development of a new C14 monolithic silica column containing embedded polar groups for pressurized capillary electrochromatography. Journal of Separation Science 30(17) (2007): 3027-3034.
- [31] O'Gara, J.E., Walsh, D.P., Alden, B.A., Casellini, P., and Walter, T.H. Systematic study of chromatographic behavior vs alkyl chain length for HPLC bonded phases containing an embedded carbamate group. Analytical Chemistry 71(15) (1999): 2992-2997.
- [32] Layne, J. Characterization and comparison of the chromatographic performance of conventional, polar-embedded, and polar-endcapped reversed-phase liquid chromatography stationary phases. Journal of Chromatography A 957(2) (2002): 149-164.



- [33] Snyder, L.R., Kirkland, J.J., and Dolan, J.W. Basic concepts and the control of separation. Introduction to Modern Liquid Chromatography (2010): 19-86.
- [34] Snyder, L.R., Kirkland, J.J., and Dolan, J.W. Equipment. in Introduction to Modern Liquid Chromatography. pp. 87-145: John Wiley & Sons, Inc., 2010.
- [35] Nakanishi, K. Synthesis concepts and preparation of silica monoliths. Monolithic Silicas in Separation Science (2011): 9-33.
- [36] Hench, L.L. and West, J.K. The sol-gel process. Chemical Reviews 90(1) (1990): 33-72.
- [37] Ishizuka, N., Minakuchi, H., Nakanishi, K., Hirao, K., and Tanaka, N. Chromatographic characterization of macroporous monolithic silica prepared via sol-gel process. Colloids and Surfaces A: Physicochemical and Engineering Aspects 187-188(0) (2001): 273-279.
- [38] Zhao, G., Dong, X.-Y., and Sun, Y. Ligands for mixed-mode protein chromatography: Principles, characteristics and design. Journal of Biotechnology 144(1) (2009): 3-11.
- [39] Yang, Y. and Geng, X. Mixed-mode chromatography and its applications to biopolymers. Journal of Chromatography A 1218(49) (2011): 8813-8825.
- [40] Sander, L.C. and Wise, S.A. Shape selectivity in reversed-phase liquid chromatography for the separation of planar and non-planar solutes. Journal of Chromatography A 656(1-2) (1993): 335-351.
- [41] Snyder, L.R., Kirkland, J.J., and Dolan, J.W. The Column. Introduction to Modern Liquid Chromatography (2010): 199-252.
- [42] Kobayashi, H., et al. Chromatographic properties of monolithic silica capillary columns for polar and nonpolar compounds in reversed-phase HPLC. Chromatographia 60(1) (2004): S19-S25.
- [43] Motokawa, M., et al. Monolithic silica columns with various skeleton sizes and through-pore sizes for capillary liquid chromatography. Journal of Chromatography A 961(1) (2002): 53-63.
- [44] Ishizuka, N., et al. Monolithic silica columns for high-efficiency separations by high-performance liquid chromatography. Journal of Chromatography A 960(1-2) (2002): 85-96.
- [45] Snyder, L.R., Kirkland, J.J., and Dolan, J.W. Introduction. Introduction to Modern Liquid Chromatography (2010): 1-17.
- [46] Tanaka, N., et al. Monolithic silica columns for high-efficiency chromatographic separations. Journal of Chromatography A 965(1-2) (2002): 35-49.

- [47] Guiochon, G. Comparison of the performance of particle-packed and monolithic columns in high-performance liquid chromatography. Monolithic Silicas in Separation Science (2011): 127-156.
- [48] Wong, K.-S., Kenseth, J., and Strasburg, R. Validation and long-term assessment of an approach for the high throughput determination of lipophilicity (log POW) values using multiplexed, absorbance-based capillary electrophoresis. Journal of Pharmaceutical Sciences 93(4) (2004): 916-931.
- [49] Michałowicz, J. and Duda, W. Phenols sources and toxicity. Polish Journal of Environmental Studies 16(3) (2007): 347-362.



## VITA

Miss Wanvisa Rattangam was born on Friday 23th May 1986, in Ratchaburi, Thailand. She graduated with a Bachelor's degree of Science in Chemistry from Silpakorn University in 2008. Then she studied for a Master's degree of Science in Analytical Chemistry, the Department of Chemistry, the Faculty of Science, Chulalongkorn University. In 2014, she graduated with a Master's degree of Science.

