Work-related musculoskeletal disorders, psychosocial factors, work productivity, and work ability among Garment factory workers in Myanmar



A Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of Master of Public Health in Public Health Common Course COLLEGE OF PUBLIC HEALTH SCIENCES Chulalongkorn University Academic Year 2021 Copyright of Chulalongkorn University ความผิดปกติทางระบบโครงร่างและกล้ามเนื้อที่เกี่ยวข้องกับการทำงาน ปัจจัยด้านสังคมจิตใจ ประสิทธิภาพและความสามารถในการทำงานของพนักงานในโรงงานผลิตเสื้อผ้า ประเทศเมียนมา



วิทยานิพนธ์นี้เป็นส่วนหนึ่งของการศึกษาตามหลักสูตรปริญญาสาธารณสุขศาสตรมหาบัณฑิต สาขาวิชาสาธารณสุขศาสตร์ ไม่สังกัดภาควิชา/เทียบเท่า วิทยาลัยวิทยาศาสตร์สาธารณสุข จุฬาลงกรณ์มหาวิทยาลัย ปีการศึกษา 2564 ลิขสิทธิ์ของจุฬาลงกรณ์มหาวิทยาลัย

Thesis Title	Work-related musculoskeletal disorders, psychosocial
	factors, work productivity, and work ability among
	Garment factory workers in Myanmar
Ву	Mr. Tun Win Oo
Field of Study	Public Health
Thesis Advisor	Assistant Professor NUTTA TANEEPANICHSKUL, Ph.D.

Accepted by the COLLEGE OF PUBLIC HEALTH SCIENCES, Chulalongkorn University in Partial Fulfillment of the Requirement for the Master of Public Health

Dean of the COLLEGE OF PUBLIC

HEALTH SCIENCES

(Professor SATHIRAKORN PONGPANICH, Ph.D.)

THESIS COMMITTEE

Chairman (Associate Professor WATTASIT SIRIWONG, Ph.D.) Thesis Advisor (Assistant Professor NUTTA TANEEPANICHSKUL, Ph.D.) External Examiner (Assistant Professor Saowanee Norkaew, Ph.D.) ตุน วิน โอ : ความผิดปกติทางระบบโครงร่างและกล้ามเนื้อที่เกี่ยวข้องกับการทำงาน ปัจจัยด้านสังคมจิตใจ ประสิทธิภาพและความสามารถในการทำงานของพนักงานในโรงงานผลิตเสื้อผ้า ประเทศเมียนมา. (Work-related musculoskeletal disorders, psychosocial factors, work productivity, and work ability among Garment factory workers in Myanmar) อ.ที่ปรึกษาหลัก : ณัฏฐา ฐานีพานิชสกุลPh.D.

บทนำ - ความผิดปรกติของทางระบบกระดูกและกล้ามเนื้อที่เกี่ยวข้องกับการทำงาน (Work-related musculoskeletal disorders:WMSDs) เป็นหนึ่งในปัญหาที่สำคัญในกลุ่มคนทำงานส่งผลให้เกิดจำนวนปีที่สูญเสียไปเนื่องจากความเจ็บป่วยและพิการ โดยเฉพาะในกลุ่มอายุระหว่าง 20 – 54 ปี ซึ่งความผิดปรกตินี้คิดเป็นสัดส่วนสูงถึงร้อยละ 45 ของกลุ่มคนที่เกิดความผิดปกตินี้ทั้งหมด โดย ปี จ จัย เสี่ ย ง ที่ ก่ อ ใ ห้ เกิ ด ค ว า ม ผิ ด ป ร ก ติ นี้ ป ร ะ ก อ บ ไ ป ด้ ว ย ปั จ จั ย ด้ า น ร่ า ง ก า ย แ ล ะ ด้ า น จิ ต ใ จ และยังมีความสัมพันธ์กับความสามารถในการทำงาน และประสิทธิภาพของงาน เพื่อการพัฒนาคุณภาพชีวิต และประสิทธิภาพในการทำงาน การศึกษาปัจจัยที่มีความสัมพันธ์กับความผิดปกติทางระบบกระดูกและกล้ามเนื้อนั้นจึงมีส่วนสำคัญในการส่งเสริมประสิทธิภาพในการทำงาน แ ต่ อ ย่ า งไร ก็ ต า ม ก า ร ศึ ก ษ า ใ น ก ลุ่ ม พ นั ก ง า น ใ น โร ง า น ผ ลิ ต เสี้ อ ผ้ า ข อ ง ป ร ะ เท ศ เมี ย น ม า ยั งมี อ ยู่ อ ย่ า ง จำ กั ด การศึกษาในครั้งนี้จึงมีวัตถุประสงค์เพื่อศึกษาความสัมพันธ์ระหว่าง ปัจจัยด้านสังคมจิตใจ ประสิทธิภาพและความสามารถในการทำงาน ต่อความผิดปกติทางระบบโครงร่างและกล้ามเนื้อที่เกี่ยวข้องกับการทำงานของพนักงานในโรงงานผลิตเสื้อผ้า ประเทศเมียนมา

วิธีการศึกษา – การศึกษาภาคตัดขวางครั้งนี้ดำเนินการเก็บข้อมูลในเดือนกันยายนปี 2564 จากพนักงานในโรงงานผลิตเสื้อผ้าจำนวน 370 คน ในพื้นที่อุตสาหกรรม Shwepaukkan โดยใช้การสุ่มตัวอย่างแบบหลายขั้นตอนในการเลือกกลุ่มตัวอย่างในการศึกษาครั้งนี้ใช้แบบสอบถามข้อมูลทั่วไปของพนักงาน และแบบสอบถามมาตรฐานเพื่อประเมินปัจจัยด้านสังคมจิตใจประสิทธิภาพ ความสามารถในการทำงาน และความผิดปกติทางระบบโครงร่างและกล้ามเนื้อที่เกี่ยวข้องกับการทำงาน

ผลการศึกษา - จากพนักงานในโรงงานผลิตเสื้อผ้าจำนวน 370 คน พบว่าความชุกของความผิดปรกติของทางระบบกระดูกและกล้ามเนื้อที่เกี่ยวข้องกับการทำงานอย่างน้อย 1 ตำแหน่งคิดเป็นร้อยละ 90 ของกลุ่มตัวอย่างทั้งหมด การศึกษาความสัมพันธ์พบว่า อายุ ระกับการศึกษามีความสัมพันธ์กับความผิดปรกติฯ ในส่วนของปัจจัยด้านจิตใจพบว่าพนักงานส่วนใหญ่มีความคาดหวังจากการทำงานในระดับต่ำ ความสามารถในการควบคุมการทำงานในระดับสูง การสนับสนุนทางสังคมในระดับสูง และมีความต้องการด้านร่างกายระดับสูง แต่อย่างไรก็ตามความผิดปรกติฯ ไม่มีความสัมพันธ์กับการขาดงานของพนักงานในช่วง 1 สัปดาห์ที่ผ่านมา ความสามารถในการทำงานของพนักงานส่วนใหญ่อยู่ในระดับปานกลาง คิดเป็นร้อยละ 66.5 และอยู่ในระดับยากจนร้อยละ 18.4 จากการวิเคราะห์ความสัมพันธ์โดยการวิเคราะห์การถดถอยโลจิสติค พบว่าปัจจัยที่มีความสัมพันธ์กับความผิดปรกติคือ เพศ (AOR=0.130; 95%CI 0.020-0.969) ความเครียดจากการทำงาน (AOR=8.257; 95%CI 1.465-46.550) ความต้องการทางด้านร่างกาย (AOR=4.702; 95%CI 1.172-18.862) และประสิทธิภาพในการทำงาน (AOR=5.893; 95%CI 1.393-24.920) จากผลการศึกษาในครั้งนี้จะสามารถนำไปสนับสนุนการพัฒนาการทำงานของพนักงานในโรงงานผลิตเสื้อผ้า เพื่อส่งเสริมคุณภาพชีวิตที่ดีของพนักงานต่อไป

สาขาวิชา ปีการศึกษา สาธารณสุขศาสตร์ 2564 ลายมือชื่อนิสิต ลายมือชื่อ อ.ที่ปรึกษาหลัก

6374502953 : MAJOR PUBLIC HEALTH

KEYWORD:

Work-related musculoskeletal disorders, Work productivity, work ability, Psychosocial factors, Garment factory workers

Tun Win Oo : Work-related musculoskeletal disorders, psychosocial factors, work productivity, and work ability among Garment factory workers in Myanmar. Advisor: Asst. Prof. NUTTA TANEEPANICHSKUL, Ph.D.

Background - Work-related musculoskeletal disorders (WRMSD) are the most common occupational problem worldwide. The magnitude of WRMSD is worsened by the physical and psychological factors in different occupations. The global burden of WRMSD comprises the second most common cause of disability; most frequent back pain, measured by Years lived with disability (YLDs) worldwide. Workability also plays a crucial role that links with productivity and WRMSD. Musculoskeletal disorders are also one of the most common causes of YLDS in the working-age group (20-54Years) take in more than 45% of the proportion. Sewing machine operators are one of the common occupational sectors that may encounter a high prevalence of MSDs. For the quality of life of workers and national productivity, we should understand the occupational hazards and their preventive approach in this working sector. Indeed, there is very little research on the occurrence of WRMSDs related to psychosocial, work productivity, and workability among the Myanmar sewing machine operators. This study aims to find the associations among working ability, productivity status, psychosocial factors, and percentage of work-related Musculoskeletal Disorders of sewing machine operators in the garment factory.

Methodology – This is a cross-sectional study that contains personal factors questionnaire, standard questionnaires, and data is collected by self-administered type which was conducted from September to October 2021. 370 Participants were selected from Shwepaukkan Industrial Zone, Myanmar. A multistage sampling method was used and eligible participants were invited to this study. University Ethical Approved was also obtained.

Result - The study enrolled 370 sewing operators who are more than 40 years of age. The prevalence of WRMSD in at least one part of the body was reported around 90 percent of the study population. The study found a significant association between WRMSD and age, education. As regards psychosocial risk, and most of the reported cases are low psychological job demand, high job control, and high social support. In addition, sewing operators responded that their job has high physical demand. There is no significant reported absenteeism and 35% presenteeism rate within one week. The moderate workability rate is about 66.5 % followed by poor workability with 18.4%. The logistic regression model showed that gender (AOR=0.130; 95%CI 0.020-0.969), job stress (AOR=8.257; 95%CI 1.465-46.550), physical demand (AOR=4.702; 95%CI 1.172-18.862) and productivity (AOR=5.893; 95%CI 1.393-24.920) are a significant predictor of WRMSD. This findings can be helped to guide the development of the working situations of sewing operators in a garment factory and also help in developing regulations for the well-being of workers in occupational health sectors

Field of Study: Academic Year: Public Health 2021 Student's Signature Advisor's Signature

ACKNOWLEDGEMENTS

This thesis became completed with the kindly guidance, support, and motivation of many individuals. I would like to express my heartfelt appreciation and gratitude to all of them. First and foremost, I express my deep sincere gratitude to my advisor, Assistant Professor Dr. Nutta Taneepanichskul, for her encouragement, guidance, suggestion, and inspiration for the completion of my work. Without her excellent suggestions and guidance, it is impossible to accomplish my MPH thesis.

I also would like to express my thankful and deep gratitude to my exam thesis committee; the chairman Associate Professor Wattasit Siriwong, Ph.D., and the external examiner Assistant Professor Saowanee Norkaew, Ph.D. who gave their precious time, valuable guidance, and remark for my thesis.

For my data collection, I do thanks sewing machine operators from the Shwepaukkan Industrial area who are participated in this study. The work will not be complete without the cooperation and support from garment factory authorities from the industrial zone.

I would like to express my gratitude to Prof. Sathirakorn Pongpanich, Ph.D., and all teachers and staff of the College of Public Health Sciences, Chulalongkorn University for providing great academic experiences in the MPH journey. I would like to thank my senior brothers and sisters, colleagues, and friends for the kindly moral support, kindness, and happiness we enjoyed together.

Last but not least, I would like to express gratitude to Professor Karasek and the professional team to give permission to the use of the Job Content Questionnaire in my thesis.

CHULALONGKORN UNIVERSITY

Tun Win Oo

TABLE OF CONTENTS

Pa	ge
ABSTRACT (THAI)iii	i
ABSTRACT (ENGLISH)iv	/
ACKNOWLEDGEMENTSv	/
TABLE OF CONTENTS	i
List of tablesx	(
List of figures	i
CHAPTER 1	
INTRODUCTION	
1.1 Background information1	
1.2 Research Gap)
1.3 Research Question	,
1.4 Statistical Hypothesis	,
1.5 Research Objective	}
1.6 Conceptual framework)
1.7 Operational definition9)
CHAPTER 213)
LITERATURE REVIEW	b
2.1 Definition of Work related Musculoskeletal Disorder13	5
2.1.1 Lower limb and non-specific MSD15)
2.2 Association between Personal factors and WRMSD16)
2.3 Work-related stress and psychosocial hazards25	

2.3.1 Prevalence of psychosocial risk factors and stress among workers	26
2.3.2 Workplace stress Theory	29
2.3.3 Karasek Demand Control Support Model	30
2.4 Work Ability and WRMDS	36
2.5 Work productivity	37
2.5.1 Conceptual model of work productivity	38
2.5.2 Relationship between psychosocial factors and work productivity	40
2.5.3 Sustainable Development Goal (SDGs) and productivity	42
2.6 Methodology Explanation	43
2.6.1 Cornell musculoskeletal Discomfort Questionnaire for WRMSD	43
2.6.2 Work productivity and Activity Impairment Questionnaire (WPAIQ) for	
work productivity	44
2.6.3 Job Content Questionnaire (JCQ) for psychosocial factors	44
2.6.4 Work Ability Index (WAI) for work ability	45
CHAPTER 3	46
RESEARCH METHODOLOGY	46
3.1 Study Design	46
3.2 Study Area	46
	40
3.3 Study population and inclusion, exclusion criteria	40
3.3 Study population and inclusion, exclusion criteria3.4 Sample Size Calculation	40 49
3.3 Study population and inclusion, exclusion criteria3.4 Sample Size Calculation3.5 Sampling Method	40 49 50
 3.3 Study population and inclusion, exclusion criteria 3.4 Sample Size Calculation 3.5 Sampling Method 3.6 Measurement Tools 	40 49 50 52
 3.3 Study population and inclusion, exclusion criteria 3.4 Sample Size Calculation 3.5 Sampling Method 3.6 Measurement Tools 3.6.1. Personal factors questionnaire 	40 50 52 52

3.6.3 Cornell Musculoskeletal Discomfort Questionnaire (CMDQ	
3.6.4. Work productivity and Activity impairment questionnaire-genera	l health
Version (WPAI-GH)	
3.6.5. Job content questionnaires (JCQ)	55
3.7 Data Collection Process	56
3.8 Data Entry and Analysis	56
3.9 Ethical Consideration	60
CHAPTER 4	61
RESULT	61
4.1 General characteristics of the participants	62
4.1.1 Association of Personal factors and WRMSD	64
4.2 Percentage of WRMSD in sewing operators	67
4.3 Percentages of Psychosocial risk factors	69
4.3.1 Association of Psychosocial risk factors and WRMSD	73
4.4 Work Ability Index score of sewing machine operators	74
4.4.1 Association of Work ability and WRMSD	76
4.5 Work productivity of sewing machine operators	77
4.5.1 Association of Work productivity and with WRMSD	79
4.6 Binary logistic regression analysis of personal factors, psychosocial fact	ors, Work
ability, and work productivity with WRMSD	79
CHAPTER 5	82
DISCUSSION	82
5.1 Prevalence of WRMSD	83
5.2 Personal factors of sewing machine operators related to WRMSD	86

5.3 Association of Psychosocial risk factors and WRMSD	89
5.4 Association of work ability and WRMSD	92
5.5 Association of work productivity and WRMSD	95
CHAPTER 6	99
CONCLUSION AND RECOMMENDATION	99
6.1. Strengths and Limitations	100
6.1.1 Strengths of the study	100
6.1.2. Limitations of the study	101
6.2. Recommendation	102
6.2.1 Recommendation for program implementation	102
6.2.2. Recommendation for further research	103
REFERENCES	106
APPENDICES	129
APPENDIX A: Certificate of ethical approval by Chulalongkorn University	129
APPENDIX B: Questionnaires (English)	130
APPENDIX C: Participant information sheet and consent form (English)	141
APPENDIX C: Questionnaire (Myanmar)	146
APPENDIX D: Participant information sheet and consent form (Myanmar)	155
VITA	158

List of tables

	Page
Table 1.Data analysis	57
Table 2. General characteristic of participants (N=370)	63
Table 3.Personal factors of sewing machine operators related to WRMSD ($N = 370$)	.65
Table 4.Association of Personal factors and WRMSD	66
Table 5. Musculoskeletal disorders' prevalence at the recent week in our subjects	
with musculoskeletal disorders (N = 370)	67
Table 6. Psychosocial risk factors (N = 370)	70
Table 7.Psychosocial risk factors and WRMSD according to category $\%$ (N = 370)	72
Table 8. Psychosocial risk factors and association with WRMSD	73
Table 9. WAI individual item score (N = 370)	75
Table 10.WAI category score (N= 370)	76
Table 11.WAI score and association with WRMSD	76
Table 12. Work productivity score (n = 370)	77
Table 13.Association of Work productivity and WRMSD	79
Table 14.Binary logistic regression analysis of personal factors, psychosocial factors	,
Work ability, and work productivity with WRMSD	80

List of figures

	Page
Figure 1.Conceptual Framework	9
Figure 2. Association between poor social support and musculoskeletal problems	
Source: Hauke et al., 2011 Burnout	.33
Figure 3. Karasek Job Strain Model	.35
Figure 4. Conceptual model of work productivity	.38
Figure 5.Industrial zone map of Yangon	.48
Figure 6. Sampling frame	.51



CHULALONGKORN UNIVERSITY

CHAPTER 1

INTRODUCTION

1.1 Background information

Work-related musculoskeletal disorders (WRMSD) are the most common occupational problem worldwide. (Niu, 2010; Urwin et al., 1998). It refers to "conditions that involve muscles, tendons, nerves, and other soft tissue pain that could be associated with or influenced by work-related factors."(Bernard & Putz-Anderson, 1997) The magnitude of WRMSD is worsened by the physical and psychological factors in different occupations. WRMSD is also the most economic cost by disease and the individual has to carry the cost, thereby loss of income leads to increasing poverty (Niu, 2010; Urwin et al., 1998). The most painful areas are in low back, shoulder, and neck. Prolong sitting, standing and repetitive jobs are the major risk factors for WRMSD (Ghasemkhani, Mahmudi, & Jabbari, 2008). Loss of productivity at work and sickness absence are usually derived from WRMSD pain

and one of the significant work-related issues (C. Phillips et al., 2008).

The global burden of WRMSD comprises the second most common cause of disability; most frequent back pain, measured by Years lived with disability (YLDs) worldwide (Storheim & Zwart, 2014). In the study of the global burden of disease between 2007 and 2017, low back pain is the common cause of the increase in a ll-age years lived with disability (YLDs) with 17.5 percentages. Musculoskeletal

disorders are also one of the most common causes of YLDS in the working-age group (20-54Years) take in more than 45% of the proportion (Safiri et al., 2020).In the review study of 2007 to 2017, total disability-adjusted life years (DALYs) were also influenced by neck and back pain about 17%. Besides, as regards developing countries, the disease burden also ranked as one of the most common three causes of disability and non-communicable diseases overburden (Hoy et al., 2015; Hoy et al., 2014).

WHO data certifies that over 1000 million workers are comprised in smallscale industry. Physical, physiological risks are the reinforcing factors for the impairment of overall well-being. Musculoskeletal disorders and other cardiovascular, respiratory problems are also influenced by those factors. Garment factory workers are one of the common vulnerable groups that suffer several health risk factors all over the world. Low wages, poor working environment, presence of certain chemicals such as dust, mist, smoke are the common physical factors, and job stress, depression is the prominent psychological factors.(Kabir, Maple, Usher, & Islam, 2019; Steinisch et al., 2013)

For the development of economic growth, productivity plays a crucial role. The setting of productivity is broad and it can be defined as the traditional way of the interaction between input (working hours) and output (numbers of productions) (Escorpizo, 2008). A study suggested an indirect relationship between health risks, productivity, and worker disability. Productivity loss issue can be stated that it is not only for the laborer but can be for the employer. The cost for workers' impaired well-being as a result of health risks can be regarded as the indirect cost that may impact the economic evaluation. (Burton, Conti Dj Fau - Chen, Chen Cy Fau - Schultz, Schultz Ab Fau - Edington, & Edington)

Work ability also plays a crucial role that links with productivity and WRMSD. It can be defined as the capacity of workers in performing jobs that are influenced by job demand, health, and mental resources (Sorić, Golubić, Milosević, Juras, & Mustajbegović, 2013). It is also influenced by a sum of factors that the workers can enable to control their job demand successfully. Impaired work ability is believed to be the outcome of adverse health outcomes, individual resources, and working conditions. (Airila, Hakanen, Punakallio, Lusa, & Luukkonen, 2012; Alavinia, van Duivenbooden, & Burdorf, 2007) WRMSD is one of the common health problems in most occupations. However, few data associate with work ability and WRMSD in sewing operators. (Pacheco Ferreira, 2015)

The conceptual framework of O'Donnell also verified that there is an alliance between human productivity and work performance and is also influenced by physical and emotional factors. That means that our ability to produce rely on how healthy we are (O'Donnell, 2000). Another interesting fact is that the word 'sickness presenteeism'. It means the despite the health problem, the worker turns up in a job that should be absent in the job (Meerding, Ijzelenberg, Koopmanschap, Severens, & Burdorf, 2005). The Swedish workforce study illustrated that 'sickness presenteeism' during 12 months is suffered by about 37% of all workers (G. Aronsson, Gustafsson, & Dallner, 2000). There is also evidences that occupational stress and psychological factors influence neck and shoulder pain (Skov, Borg, & Orhede, 1996).

Sewing machine operators are one of the common occupational sectors that may encounter a high prevalence of MSDs that is often lead to repetitive arm and hand conditions, poor working postures during the whole working period. They can be overlooked by the reason of the high incidence of garment workers. (Li, Haslegrave, & Corlett, 1995)Needle prick injuries, burn, and cuts, and also due to the highly competitive nature of the world garment market, illness and risk of occupational diseases are reported in the previous study (Gupta, 2015). Usually, sewing operators need to perform fine work activities, setting up the parts for the final cloth product. The detailed nature is that the job is sometimes a complex task and highly repetitive comprising the synergism of both hands, usually over a whole working period. The functioning usually requires one to lean forward (head and trunk are forwardly inclined) to have better visual control and focus their attention on the task. The adverse outcome can lead to the progress of MSDs, mainly in the back, neck, and upper limbs (Saha, Dasgupta, Butt, & Chattopadhyay, 2010; Sealetsa & Thatcher, 2011).

Sewing operators are the major workers in that population. Studies showed that sewing operators are the common victim of MSD. Follow-up neck and shoulder pain are the most common symptoms in sewing operators in follow-up studies (Schibye et al., 1995; Wang, Harrison, Yu, Rempel, & Ritz, 2010; P.-C. Wang, D. M. Rempel, R. J. Harrison, J. Chan, & B. R. Ritz, 2007). Physical factor studies confirm that musculoskeletal incidence is also higher in garment workers of the sewing section. Increase concentration on fine works can also increase the risk of diseases (Jaffar & Rahman, 2017; Sadeghi Yarandi, Ghasemi, & Ghanjal, 2020).

Myanmar is now in the transition time to democracy and encourages the development of the economic process. Foreign investment in industrial sectors can give job opportunities for people and there is can be an economic benefit. However, rapid developments of the industrial sector need strong labor regulation, workers protection, and occupational health care. The weak labor laws in Myanmar in these sectors result in the underestimate of the health burden of workers (Burnley, December 2015).

Garment factories in Myanmar are also comprised of the country's economic growth and one of the major sectors of employment. In the recent decade, the Myanmar garment sector has been grown and approximately between 350,000 and 450,000 workers, comprising 905 are women and more than 60 percent of the industry are foreign owners. One of the Oxfam research, Low wages, and lack of proper safety issue, long working hour, and their financial debt are the most concern problems in garment sectors. To solve their financial problem, at least 1 in 4 reported that overtime sometimes becomes their additional income. But due to poor health and labor regulation, the sad story is forced doing overtime and unpaid overtime. Their cycle of poverty and health problems cannot end (Burnley, December 2015; Htay, 2019).

For the quality of life of workers and national productivity, we should understand the occupational hazards and their preventive approach in this working sector. Indeed, there is very little research on the occurrence of WRMSDs related to psychosocial, work productivity, and work ability among the Myanmar sewing machine operators. The research findings in this area will guide the development of the working situations of those jobs regarding sewing operation in a garment factory and also help in developing regulation in preventive measures. It also has the prospective to make proper work practices in similar industrial sectors Nationwide.

1.2 Research Gap

Although there are many kinds of research about WRMSD and psychological factors. However, there is little literature linked work productivity, work ability with WRMSD. Few articles were conducted on sewing operators in garment workers by using those factors. Besides, there is little known about the situation of WRMSD in Myanmar. Therefore, this research will identify a relationship between psychological factors, work productivity, work ability, and WRMSD.

1.3 Research Question

What are the working ability, productivity status, psychosocial factors, and percentage of work-related Musculoskeletal Disorders of sewing machine operators in the garment factory?

What is the association between psychosocial factors and WRMSD?

What is the association between work productivity and WRMSD?

What is the association between work ability and WRMSD?

1.4 Statistical Hypothesis

Statistical Hypothesis 1

Null hypothesis = there is no association between psychosocial factors and workrelated musculoskeletal disorders **KORN UNIVERSITY**

Alternative hypothesis = there is an association between psychosocial factors and work-related musculoskeletal disorders

Statistical Hypothesis 2

Null hypothesis = there is no association between work productivity and workrelated musculoskeletal disorders

Alternative hypothesis = there is an association between work productivity and work-related musculoskeletal disorder

Statistical Hypothesis 3

Null hypothesis = there is no association between work-related musculoskeletal disorders and work ability

Alternative hypothesis = there is an association between work-related musculoskeletal disorders and work ability

1.5 Research Objective

- To find the working ability, productivity, psychosocial factors, and percentage of work-related Musculoskeletal Disorders of sewing machine operators in the garment factory.

- To find an association between psychosocial factors and work-related musculoskeletal disorders

- To find an association between work productivity and work-related musculoskeletal disorders

- To find an association between work ability and work-related musculoskeletal disorders

1.6 Conceptual framework



Figure 1.Conceptual Framework

1.7 Operational definition

Sewing Machine Workers = A sewing machine operator is a textile worker who sews fabric with an electronic sewing machine in the garment factory.

Age = Age is referred to the age of sewing operators in years at the time of study

Gender = Gender is preferred to male and female

Marital status = the marital status referred to single, married, separated, and

widowed

BMI = It is measured by dividing body weight (kilogram) and the square of the weight

(meter) at the time of data collection

Education = it is referred to the primary school, secondary school, high school, and above

Monthly income = monthly salary by the workers, measured in kyats

Working hour per day = it is referred to the number of working hours per day as sewing operators

Work experience = it is referred to the number of working years as sewing operators Work-related musculoskeletal disorders = this referred to the pain all over the body related to work. In this study, WRMSD will be measured by Cornell Musculoskeletal Discomfort Questionnaire. It measures the frequency of the last working week by five categories(Never, 1-2 times last week, 3-4 times last week, Once every day, Several times every day), severity by three categories (slightly uncomfortable, moderately uncomfortable, very uncomfortable), interference by three categories (Not at all, slightly interfered, substantially interfered)

Work productivity = is referred to as the assessment of efficiency of workers.in this study, it will be measured by Work Productivity and Activity Impairment Questionnaire (WPAIQ). The prevalence of presenteeism, absenteeism, overall work impairment is regarded as decrease work productivity.

Presenteeism = the worker is in work but decreased production due to his illness. It is expressed in percentages by WPAIQ.

Absenteeism = lost productivity due to workers absent from the job. .It is expressed in percentages by WPAIQ.

Overall work impairment = impact on productivity due to the combination of absenteeism and presenteeism. It is expressed in percentages by WPAIQ.

Job control =" it is referred to person's ability to influence what happens in his or her work environment, in particular, to influence matters that are relevant to his or her job condition.". Scoring will be dichotomized according to the median cut-off point, High (>median score), and Low (< median score) according to the Job Content Questionnaire (JCQ).

Psychological demand= it is referred to the amount of mental strain that workers get from his/her job. Scoring will be dichotomized according to the median cut-off point, High (>median score), and Low (<median score) according to JCQ.

จุฬาลงกรณ์มหาวิทยาลัย

Physical job demand = it is referred to as the physical exertion effort by the workers. Scoring will be dichotomized according to the median cut-off point, High (>median score), and Low (<median score) according to JCQ.

Job insecurity= "it is referred perceived and an undesired possibility to lose the present job in the future, as well as the fear or worries related to the possibility of job loss." Scoring will be dichotomized according to the median cut-off point, High (>median score), and Low (<median score) according to JCQ.

Social support= is referred to as the support from supervisors and coworkers in the working environment. Scoring will be dichotomized according to the median cut-off point, High (>median score), and Low (<median score) according to JCQ.

Work ability = it is referred to the capability of workers both physically and mentally following the job demand. In this study, it is measured by Work Ability Index. It consisted of 7 items, including(item, score range), current work ability compared with the lifetime best (item 1, 0-10), work ability concerning the demands of the job (item 2, 2-10), number of current disease groups diagnosed by a physician (item 3, 1-7), - / / / htold / / / estimated work impairment due to diseases (item 4, 1-6), sick leave during the past V / / MANA year (item 5, 1-5), the personal prognosis of work ability for 2 years from now (item V Streeter Dooroot N 6, 1,4 or 7) and mental resources, referring to the worker's life in general, both at work and during leisure time(item 7, 1-4). The number in parentheses for each item indicates the scoring range. The total WAI score is calculated by summing up the scores of all items and is ranges from 7 to 49. The total WAI scores are categorized into 4 levels: poor (7-27), moderate (28-36), good (37-43), and excellent (44-49). (Adel, Akbar, & Ehsan, 2019)

CHAPTER 2

LITERATURE REVIEW

2.1 Definition of Work related Musculoskeletal Disorder

MSDs include pain in all musculoskeletal systems, bone, tissue, and muscle, when there is an event or exposure to the injury or illness, and body changes such as bending, climbing, crawling, reaching, twisting; overexertion; or repetitive job (Birschel, 2006). Physical, individual characteristics, mechanical and psychosocial conditions are the major contributor to the development of MSD.

The existence of awkward or fixed postures for prolonged periods during the job performance, extreme temperature, and vibration may worsen the MSD.(Bernard & Putz-Anderson, 1997) Epidemiological research showed that the onset and severity of hand and wrist in forceful and repetitive hand task overuse are the main cause of MSD. (A. E. Barr, Barbe, & Clark, 2004). Both systemic and local inflammatory responses are associated with the performance of highly repetitive tasks that may result in a decline in job speed and effectiveness of the movement. A chronic inflammatory response may develop when there is an elevated repetition rate.(Ann E. Barr, Amin, & Barbe, 2002)

Fibrosis and inflammation can result from chronic contraction of a muscle (A. E. Barr et al., 2004). Using a rat model, repeated muscle strains at fast velocities can damage the muscle and result in myopathy with the development of macrophage and cellular protein. Another intervention in this paper also described that the low speed (25 mm/s) with persistent strains leads to fibrosis. Regarding the adaptive and compensatory response; muscle tonic repetition should be in 10mm/s velocity (A. E. Barr et al., 2004).

One study reports that the participants with upper MSD pain showed a significantly reduced pressure pain threshold in automobile production workers(Gold, Punnett, & Katz, 2006). He also reported that the depressive symptoms, higher work stress, anxiety, conflict with supervisors, higher pain threshold, enhanced sensitivity to pain feeling, and low self-confidence in handling complications have been described in epidemiological and clinical studies of patients with MSDs (Gold et al., 2006).

MSD has influenced mainly three factors (Nunes, 2009)

จหาลงกรณ์มหาวิทยาลัย

1. Individual factors – age, gender, physical activity, working hour, duration of

working, underlying comorbidities, previous MSD, alcohol, smoking, etc.

2. Physical factors –awkward posture, repetitive movement, longer duration, vibration, forceful exertion, etc.

3. Psychosocial factors – job demand, job pace, physical workload, social support from supervisors and coworkers, job insecurity, etc.

WRMSD can be classified according to anatomical site. (Kuorinka et al., 1995)

1. Tendon - comprises inflammation of the tendon and/ or synovial sheet. They are called tendonitis and if it involved synovial sheet, it is tenosynovitis which is the result of the injuries of tendon and synovial sheath

2. Bursa - It is the inflammation of the burs space called bursitis.

3. Nerve - It is involved when compression of the nerve has occurred, e.g. carpal tunnel syndrome

4. Muscle - fatigue muscle e.g. Tension Neck Syndrome

5. Vessels - result from the impairment of vessel and blood flow due to repetitive work e.g. vibration syndrome

2.1.1 Lower limb and non-specific MSD

These are currently a problem in many job activities. They are underestimated even there are recognized activities (e.g., kneeling/ squatting, climbing stairs or ladders, heavy lifting, walking/standing) are risk factors for the lower limb(Bush, 2012). Moreover, they are worth significant concern because these are the major factors for disability and impair the quality of life (Okunribido & Excutive, 2009). The most common disorders are

Hip/thigh – Osteoarthritis (frequent), Piriformis Syndrome, and Hamstring strains, Sacroiliac Joint Pain; Knee / lower leg – Osteoarthritis, Bursitis, Beat Knee/Hyperkeratosis, Meniscal Lesions, Patellofemoral Pain Syndrome, Pre-patellar Tendonitis, Shin Splints, Infrapatellar Tendonitis, Stress Fractures;

Ankle/foot – Achilles Tendonitis, Blisters, Foot Corns, Plantar Fasciitis, Sprained Ankle, Stress fractures, Varicose veins, venous disorders.

Non-specific WRMSD are called the ill-defined site of pain, meaning that symptoms are diffuse all over the body, no specific anatomical site, nerves, muscles, tendon, and reach into the other anatomical location. They presented with pain (worse over-activity), numbness, tingling, cramping without finding out any pathological underlying causes(Ring, 2005).

2.2 Association between Personal factors and WRMSD

Numerous studies are reported that employee personal factors are the predictor of workplace injuries and disorders. Age between 35 to 55 years of age, female, single, receiving medical treatment, lesser education, working as a operators or repair personnel of manufacturing is the risk of work-related injury and more prone to sick leave(Albert, Coutu, & Durand, 2011; Berecki-Gisolf, Clay, Collie, & McClure, 2012; V. Phillips, Hunsaker, & Florence, 2012).

A systematic longitudinal study identified that personal factors such as age, gender, body mass index, working duration, psychosocial factors are influenced to the risk of WRMSD in different parts of the body. Also, they found out that no strong evidence for those risks and only reasonable evidence and insufficient evidence. Most of the WRMSD are related to heavy physical work, awkward positions, and repetitive jobs.(Costa & Vieira, 2009)

Researches by (S. M. Maduagwu et al., 2014) reported that WRMSD is more prevalent in young age between 20-29 years of age than the oldest age group in that study. They explained that these are due to low job experience and low professional skills during that age. Moreover, these age groups exert more workload than the other age group. This result in line with the report of (Afonso, Pinho, & Arezes, 2014) with the younger workers is more prevalent in WRMSD in the sewing sectors of the footwear industry. But some research show contradicts that old age is more vulnerable to WRMSD than young. (Aghili, Asilian, & Poursafa, 2012) showed that older age is more associated with the prevalence of WRMSD. There are also some articles that they found no association between age and WRMSD.(S. Maduagwu et al., 2015; Soe, Laosee, Limsatchapanich, & Rattanapan, 2015)

In the study of (Abaraogu, Okafor, Ezeukwu, & Igwe, 2015), they stated that the prevalence of WRMSD is increasing in old age particularly over 40 years of age. They found that the prevalence of neck, shoulder, lower back, and thigh pain is 100% in over 50 years of age. Over 40 years of age have statically similar prevalence and under 30 years of age are comparable prevalence.

Another influential factor is gender that is related to whole-body pain. (Karlqvist, Tornqvist, Hagberg, Hagman, & Toomingas, 2002) found that if both males and females are working on a similar task, females have different working postures than males. (Punnett, 2000) also, point that being female is also the "risk factor" for musculoskeletal disorders and due to the gender difference in the labor market and demand, it is important to distinguish between gender difference prevalence of S 11 13 -WRMSD. Women are facing strong hormonal changes during menopause and pregnancy that make them suffer more WRMSD due to increased fluid retention and some physiologic conditional changes. Other factors can be muscular strength, anthropometric issues, and hormonal changes, and more risk to carpel tunnel syndrome. (Bush, 2012). Many kinds of research also proof that female is more vulnerable to WRMSD than man even though they counter the same stressor. YA. Moreover, females tend to be different working postures and overall symptoms are more common in females. This finding also supports the research of (Abaraogu et al., 2015) that females are more symptoms in all parts of the body than males except the knee. However, gender and WRMSD association are not always the same. A (Taib, Bahn, Yun, & Taib, 2017) survey of 85 Malaysian dentists showed that age and gender are not associated with WRMSD. (Rocha & Debert-Ribeiro, 2001) study of system analysts also points to the same outcome.

Reported symptoms in all parts of the body are different between male and female. Male are more suffer pain in the upper back and females are more reported in hand/wrist symptoms. Moreover, age, gender, BMI are also significant associations with WRMSD (Dianat, Kord, Yahyazade, Karimi, & Stedmon, 2015). In the study of (Hossain et al., 2018) garment workers in Bangladesh, females more suffered from lower back and males reported the majority in neck pain of 12-month prevalence. To sum up, the association between gender and WRMSD is not clearly stated but females are more prevalent in WRMSD.

Regarding marital status, some researches showed an association and some are not. One of the research of (Soe et al., 2015)Myanmar migrant workers in Thailand fish industry showed that marital status and number of dependent people were significantly associated with WRMSD. They also explained that this can be due to working long hours such as overtime to earn more money to support their family. They did not clearly state the working hour and marital status association. This is also similar to the study in India that married seafood workers are more likely to get upper back MSD(Nag, Vyas, Shah, & Nag, 2012). Similarly, the two studies of sewing operators (Aghili et al., 2012; Jahan et al., 2015)also reported that marital status is a significant predictor of WRMSD.

(Oranye et al., 2016) research showed no association of personal factors such as age, gender, marital status, and education. But when they did the correlation analysis, marital status is the only factor that correlates with the WRMSD. Single workers are more reported of MSD than married/divorced. They explained this association by being positivity of married person that lowers the risk of workplace injuries or disorders. The 12-month prevalence of lower back pain was significantly associated with the marital status of the participant and more married reporting in blue color labor study(Kaka et al., 2016). However, there is also numerous study that showed no association between marital status and WRMSD. (Amin, Nordin, Fatt, Noah, & Oxley, 2014; Koyuncu & Karcioglu, 2018; Tantawy, 2019) Therefore, there is uncertainty about the association between marital status and WRMSD.

Body Mass Index is also another independent factor that influences WRMSD. It is measured by weight and height and is also recognized as a risk factor for MSD(Yang, Matthews, & Chen, 2014).BMI also measured the body adipose tissue and increase BMI had tended to be more MSD than lower BMI people (Viester et al., 2013). Increasing BMI, obesity can be associates with psychological, physical, WRMSD that impair productivity (Lidstone et al., 2006). Many researches related to WRMSD include BMI regarded as one of the risk factors. An (Aghilinejad, Choobineh, Sadeghi, Nouri, & Bahrami Ahmadi, 2012) study reported that higher BMI is significantly associated with WRMSD in Iranian steelworkers. Higher BMI is related to job efficacy and is more prone to work-related trauma. Some researchers found no association between BMI and WRMSD (Amin et al., 2014; Tantawy, 2019) Sewing operators are more likely to increase BMI because their job nature is sitting or standing without proper movement and lead to sedentary habits (Hossain et al., 2018).BMI is also important from an ergonomic point of view that is related to WRMSD. (Sibella, Galli, Romei, Montesano, & Crivellini, 2003) showed that obese workers had remarkably low trunk flexion in sit-to-stand positions and were more vulnerable to low back pain. While (Godde & Taylor, 2011) found that obese people relied on upper limbs than non-obese individuals. This can be the reason that high BMI people are more reported to upper limb WRMSD.

Many researchers found that education is not a risk factor for WRMSD among sewing operators (Rohilla, Hooda, & Mehta, 2020; Sakthi Nagaraj, Jeyapaul, & Mathiyazhagan, 2019). Reversely,(Kaka et al., 2016) found that lower back pain is related to educational attainment. They did not explain in detail about that association. But adults with higher education can lead to better health than lower education. There is much evidence that education strongly influences individual health (Zajacova & Lawrence, 2018). Not many articles were writing about education and occupational health. They only state the general comment about the heath. For the well-being of workers, education is also comprised as an important role. To look at the occupational point of view, education level is related to high income, a better job can handle social, psychological stress, likely to learn about healthy behavior. The relationship is reverse causality. Poor health not only comes from lower education but also impair school performance (Health, 2015). It is also beneficial to economic and social resources. A higher educated individual is less likely to be unemployed and have economic hardship because it involves "access to resources that can be used to avoid risks or to minimize the consequences of the disease once it occurs"(Link, Phelan, & behavior, 1995; Zimmerman, 2014). (Kaufman & Rosenstock, 1991) study also stated that changing the educational level of workers can result in benefits in the prevention of occupational lung diseases.

Income and WRMSD are not directly related. But Myanmar garment workers are paid by day-to-day fees and they will not get income when they can't come to work. WRMSD is one of the leading causes of absenteeism (Burnley, December 2015). MSD also the significant factor that cost workers, employers, family, and the wider economy (Bevan, 2015). Developing countries are facing health inequalities concerning occupational aspects. This is the reason due to unequal income across the population(Scotland, 2021). Although many researches did not found an association between MSD and income, monthly salary still is an important factor for the health and wellbeing of workers (Jahan et al., 2015; Soe et al., 2015). Moreover, economic/financial aspects (e.g. salary, equity, and benefit) are also comprised of psychosocial risk factors that most of the workers worry about, and it may lead to WRMSD problems.(Bernard & Putz-Anderson, 1997). Work schedule is one of the most important factors that contribute to the MSD. The two studies (Krause, Rugulies, Ragland, & Syme, 2004) and (Engkvist, Hjelm, Hagberg, Menckel, & Ekenvall, 2000) showed that the prolonged working workers are related to low back pain in transit operators and nurses respectively. Longer duration of work without a break (e.g. sitting in the same posture without moving for 2 hours) in sewing operators are related to neck and shoulder pain (Dianat et al., 2015; P. C. Wang, D. M. Rempel, R. J. Harrison, J. Chan, & B. R. Ritz, 2007). Also, a study conducted by(Johnston, Souvlis, Jimmieson, & Jull, 2008) revealed that regular rest breaks during work reduced the prevalence of neck and shoulder pain in computer workers. They also advised taking a regular break to reduce exposure.

116610

Sewing operators are regarded as sedentary workers. Prolong-sitting work nature is also associated with the increased risk of MSD (Nourbakhsh, Moussavi, & Salavati, 2001). The hypothesis that explained the association between MSD and long working hour is that increasing in working time also increases to workers physical demand and lead to the higher prevalence of MSD. From an ergonomic point of view, long working hours impair the recovery time of fatigue and not enough leisure time for a refresh. This complex phenomenon is directly involved in disorders of workers' musculoskeletal system and finally induces WRMSDs (Caruso et al., 2006). Many researches were proof that prolongs working hours have been regarded as a risk for WRMSD in different occupational settings (Fogleman & Lewis, 2002; Johnston et al., 2008; Nazari, Mahmoudi, Dianat, & Graveling, 2012). The number of working hours also positively associates with knee joint pain of hand-sewn shoe workers (Dianat & Salimi, 2014).

Long ago studies are showed that years of working as sewing operators are related to upper body MSD(J. H. Andersen & Gaardboe, 1993). (Blåder et al., 1991) the study also reported that the working year as a sewing operator is the predictor of WRMSD. This statement is in line with the researches of sewing operators and computer-telephone interactive tasks workers that neck and shoulder pain are positively associated with working years (Dianat et al., 2015; Ferreira Jr & Saldiva, 2002). However, there is some debate about the researches that some are found an association and some are not. The study of (Soe et al., 2015)showed no association of working years with MSD compared between more than three years and less.

On the other hand, a big study of work organizational and personal factors of **Puril AL AUCODA HAMPORITY** sewing operators reported that distal upper extremities pain is related to year of employment but not to neck, shoulder pain (P.-C. Wang et al., 2007). Job experiences were also associated with MSD(Ahmad et al., 2007; Hossain et al., 2018). The higher percentage of MSD was seen in municipal sewage workers is attributed to a longer duration of employment and other factors (Reddy & Yasobant, 2015). Therefore, many researches are supported that year of employment is also the independent variable that influences the MSD.
2.3 Work-related stress and psychosocial hazards

Work-related stress is defined as the "determination by work organization, work design and labor relations and occurs when the demands of the job do not match or exceed the capabilities, resources, or needs of the worker, or when the knowledge or abilities of an individual worker or group to cope are not matched with the expectations of the organizational culture of an enterprise" (ILO, 2016). The major workplace factors that induce stress at work are called psychosocial hazards.

Psychosocial factors (hazards) were defined by the ILO in 1984 as the "interactions between and among work environment, job content, organizational conditions and workers' capacities, needs, culture, personal extra-job considerations that may, through perceptions and experience, influence health, work performance, and job satisfaction".

The major causes of psychosocial factors are

1. Content of work – working environment and equipment, job design, workload and pace, work schedule

2. Context of work – Organizational and cultural function, social relation and support with supervisors and coworkers, decision latitude and job control, role and personal factors in an organization, career development, home-work interface(ILO, 2016) It is now widely accepted that psychosocial factors and work-related stress affect the workers' health and productivity. It also contributes to the major causes of absenteeism and presenteeism. Stress itself is not a disease, but it is the first sign of a health problem. When the body suffers continuous strain over the mind, it can lead to long-term damage to the body and organs. It is also widely recognized that longer stress can induce memory loss, peptic ulcer, bowel inflammation, and musculoskeletal disorder, hypertension, cardiovascular diseases. It may also impair the immune system and finally may lead to cancer. Taken together, these chronic diseases are leading causes of disability, death, and economic cost and compensation burden all over the world. (ILO, 2021b)

The role of psychosocial factors on WRMSD has been proved in many epidemiologic studies, showing the linkage between them and increased attention in the recent decade. Overall, it is obvious that MSD is associated with high job demand, low job control, poor social support. High job stress level, low job satisfaction. Workplace violence particularly harassment, bullying, effort-reward imbalance, poor social relation with supervisors and coworkers are also proved to be associated with MSD (Forastieri, 2016).

2.3.1 Prevalence of psychosocial risk factors and stress among workers

The data on psychosocial hazards and work-related stress vary according to the extent of the region and countries across the world. Most of the reliable data is found only available in Europe and North America, generally developed countries, and lesser available in Asia –Pacific and Latin America. However, these psychosocial hazards remain the major hazard across the world.

The regional data is mainly collected in Europe by EU agencies. The data of the 4th European working survey reported that work-related stress is suffered by an estimated 400 million people (Eurofound, 2007). The European risk Observatory report in 2009 revealed that 22% of European workers are suffered from stress, which is more prominent in new member states (30%) than older member states (20) %. Moreover, the study also confirmed that 50% to 60% of work loss days are related to psychosocial stress. More prevalent stress is seen in the education and health sector, as well as in the hunting sector, agricultural, forestry, and fishing (28%). The largest group of the occupational sector that is suffered from anxiety are in education and health (12.7%), public administration and defense sectors (11.1%), and agriculture, forestry & fishing, hunting, (9.4%). Irritability was also most common in education and health (15.5%), transport and communication (13.6%), and hotels and restaurants, and public administration and defense (12.6%)(Work, 2009).

Job intensity is also a major problem in work society. High work intensity is also associated with a negative impact on health. Numerous epidemiologic studies were also stated that a high level of job demand is also related to cardiovascular diseases, musculoskeletal diseases, and depression. The 6th European working conditional Survey revised in 2017 reported that 36% of workers are facing working to a tight deadline and 33% are working at a high job pace around three-quarters of overall working time. The study also includes social support from the college and the supervisors. The prevalence of social support from the college in 2015 is 71% of workers in EU 28 and social support from manger is also reported high 58% but less than from the support of the college (Eurofound, 2017).

11113

The Asia, pacific region, (Casey, 2014)showed that 32% of the stressor are related to work and impair well-being. Moreover almost half of the participants said that demanding work is the common burden to achieve a healthy lifestyle. This also supports the previous year's survey that 72% of Australians suffered from at least one of the stress can cause a physical impact on health and 17% only report that current stress may have caused strong physical health impairment. In the first Korean Working Conditions Survey in 2006, 18.4 % of males and 15.1 percent of the female workforce are suffered from work-related stress which is significantly associated with job demand and working hours. In the second time survey in 2010, the overall fatigue level of the working population uprises from 17.8% in 2006 to 26.7% in 2010. However, general depression, anxiety level, insomnia, general sleep difficulties are decreased through four years period (Kim, Park, Rhee, & Kim, 2015). One of the surveys on the Prevention of Industrial Accidents in Japan also revealed that 32.4%

of workers are experienced stress, anxiety, and distress during the previous year.(MHLW, 2011)

2.3.2 Workplace stress Theory

Workplace theory is come in attention to know furthermore. In recent decades, popular theories were tested by recent researches. These theories use different perspective views to achieve the essential parts of stress formation.

The first popular one is the transactional theory of work-related stress. This is the model developed by (R. S. J. S. Lazarus, appraisal, & coping, 1986). They proposed that stress is the result of interaction with the environment and the individual. But later years, the concept is not complete because they only state the simplicity of the interaction and not considering individual ability, perception, experiences, personality, and several factors (Harris, Daniels, Briner, & Counselling, 2004). Yet, Lazarus also stated in his later work that his transaction theory failed to meet the outcome associated with the other social context and interpersonal relationship(R. S. J. J. o. p. Lazarus, 2006). another one is the interactional theory of stress also developed by(R. S. Lazarus & Launier, 1978). It is the oldest personenvironment fit theory of stress emphasis the interaction between an environmental stimulus and associated with the individual response.

2.3.3 Karasek Demand Control Support Model

The leading model to measure workplace stress and well-being is the job Demand control Support model (JDCS). It is first proposed by Robert Karasek, a sociologist at the University of Massachusetts at Lowell in 1979. The original concept mainly focuses on two important aspects of work situation; job demand and job control. In the later decade, a social aspect was added by (Johnson, Hall, Theorell, & health, 1989; Johnson & Hall, 1988) and finally became the Job Demand Control Support model.

This is mainly based on the interaction between job demand controls that results in stress and strain-related outcome. He also explained that working condition that was balanced between job demand and job control is important in a way to gain explain the connection between health and work. High job demand itself is not injurious but when it is combined with low decision latitude, job strain and stress can occur (Karasek Jr, 1979). This model also speculates that a person with high job demand, low job control, low decision latitude, and poor social support from coworkers and supervisors has the greatest risk to suffer physical and psychosocial hazards. (iso-strain hypothesis)(Johnson & Hall, 1988). Therefore, three major aspects of job factors; job demand, job control, social support are a major impact on a worker's well-being(Sargent & Terry, 2000). Bernal et al. (2015) reported that a correlation between the high psychosocial working condition and low levels of job control with pain in the back, neck, shoulder, and knee in the systemic study. (Hauke et al) (Hauke, Flintrop, Brun, & Rugulies, 2011) Review article showed the increased upper extremities pain, low back pains are combined with high job strain and neck, shoulder symptoms are associates with low job control. High job strain and low job control are about three times higher risks to impair health status (Sundquist & Johansson, 2000).

Job demand can be defined as the effort of the person to complete the job. While physical factors are the major contributor to job demand, psychological, organizational, social factors are also determinants of job demand. These include time pressures, the amount of heavy workload, a stressful working environment, role ambiguity, work pace, and poor relationships(Bakker & Demerouti, 2007),

Job control refers that how the worker influences the job task and sometimes **Current Reference Herrory** it is called decision latitude. It comprises two components skill discretion and decision authority. Skill discretion describes that job involves less repetition, level of skill, and creativity required for the job and occasionally arises the opportunity to learn new skills for career development. Decision authority means that the employees' ability to influence the decision about their job and also their work team, including in the decision of their respective job policymaking (Karasek Jr, 1979; Karasek et al., 1998). Job insecurity can be defined as "the perceptive feeling that undesired possibility of losing the job in future and also the fear or possibility related to the loss of job." In other words, it can affect both qualitative aspects such as worrying about loss of job and quantitative aspects related to the job such as health insurance, salary, and social relation, social life (De Witte, 2005; Vander Elst, De Witte, De Cuyper, & Psychology, 2014). Moreover, it has been linked with adverse health outcomes. Psychosocial stress such as anxiety, mental disorders, loss of self-esteem, minor psychiatric symptoms are associated with job insecurity. Besides, it has been found that job insecurity can lead to physical activity restriction due to the occurrence of musculoskeletal disorders such as neck and low back pain (Kivimäki et al., 2001; Lang, Ochsmann, Kraus, Lang, & medicine, 2012).

Social support refers to the support from the supervisors as well as from the college. Poor social support from both coworker and supervisor are also increase risk of psychosocial hazards.(Van der Doef, Maes, & stress, 1999) .Poor social support related to musculoskeletal diseases is already reported in many types of research. (Clays et al., 2007) studied both genders of huge sample size cohort study related to social support. A significant relationship between low back pain and poor supervisor, coworker support is found in men. This includes muscle pain at all sites as well as specifically in the lower back, neck, and shoulders. Moreover, (Hauke, Flintrop, Brun, & Rugulies, 2011) in a meta-analysis of Hauke et al. (2011) also points out that that

- ANN NEW -

there is a remarkable association between poor social support and musculoskeletal problems in the neck/shoulders, upper body parts, and low back.



Figure 2. Association between poor social support and musculoskeletal problems Source: Hauke et al., 2011 Burnout

Chulalongkorn University

The Karasek model also has a huge influence on occupational literature, designing job programs in part because it is quite spare, practical, and testable. The model hypothesis also involved that job stress is a function of how much demand the work and how much control (discretion, authority or decision latitude, etc.) that the person has depended upon responsibilities. This creates four kinds of jobs (F. Jones, Bright, & Clow, 2001).

<u>High strain jobs</u>

They have resulted from the combination of high psychological demand and low decision latitude (high demand, low control). Usually, the workplace situation is rigid and tense, not only in the working environment but also in the policy. This type of job can cause physical and psychological illness among workers and they found difficulty in coping with stress. E.g. garment workers, waiters, and nurse aid.

Passive jobs

A passive job is the combination of low decision latitude and low demand (low demand, low control). This kind of job does not require special skills, easy-going. They also suffer physical and mental illnesses. E.g. janitor, mining, and other manual

labor jobs

Low strain jobs

Few psychological demands and a high level of control (low demand, high control) and usually required creativity and skills. In this kind of job, the employee can benefit from a high level of health and happiness. E.g. architects and natural scientists.

Active job

High psychological demand and high level of control (high demand, high control). In this working environment, the employee can develop new skills and can solve the work problem freely. By doing this, they can maintain their health and reduce the level of job stress. E.g. physicians, teachers, and engineers



Figure 3. Karasek Job Strain Model

To sum up, the Karasek demand control support model is the world know theory to measure psychosocial hazards and job stress. The work nature of sewing operators includes high psychosocial demand and low job control. They are facing with high work pace, have to sew the clothes timely due to factory requirement and many psychosocial hazards., Moreover, we can access the type of job that sewing operators are working on with the help of this model. Therefore, I used the Job content Questionnaire that is based on the job demand-control model by Prof Karasek which contains 27 items. Although there are many research questionnaire, JCQ is fit with the job nature of sewing operators in Myanmar.

2.4 Work Ability and WRMDS

Work ability is a multidimensional concept that shows the balance between workers' capacity and occupational physical demand. (J. J. S. j. o. w. Ilmarinen, environment & health, 2009) Multiple occupational factors affect work ability such as obesity and especially low physical capacity that suffers from WRMSD.(L. L. Andersen, Izquierdo, Sundstrup, & health, 2017; van den Berg, Elders, de Zwart, Burdorf, & medicine, 2008) Furthermore, physical work - typically involving, awkward postures, repetitive work, and heavy lifting - has been associated with increased risk of WRMSD(Da Costa & Vieira, 2010; Søgaard, Sjøgaard, & reviews, 2017) and reduced work ability(El Fassi et al., 2013; T. van den Berg et al., 2008). However, workers with MSD symptoms can still work without showing reducing work ability. (Pensola, Haukka, Kaila-Kangas, Neupane, & Leino-Arjas, 2016). The article of (Silvia Monteiro, Maria Costa Alexandre, Ilmarinen, Mendes Rodrigues, & Ergonomics, 2009) showed that there is a significant association between work ability and WRMSD but the article (Akodu & Ashalejo, 2019) showed no association between them. The study of three variables such as work ability, quality of life (QOL), and WRMSD reported that work ability was positively associated with overall QOL, and with physical and psychological domains as well. The improvement of WRMSD in workers can enhance that work ability and quality of life of workers and they found significant association between them.(Chang et al., 2020). Therefore, to determine the work ability that associated with WRMSD can be important aspect for the sewing operators for the development well-being of workers.

2.5 Work productivity

A basic concept revealed that when a person is physically and mentally sound to be able to work, and the increased desire to work, it can lead to increase performance and overall affect the increased productivity. Health-promoting programs are also involved to target worker's health status to be improved and finally aimed to productivity enhancement and economic. To promote physical and emotional ability, mainly focus on cardiovascular and musculoskeletal problems, job condition, and ergonomic improvement and help them to cope with job stress (O'Donnell, 2000).

Mainly there can be two types of productivity loss. "Absenteeism means that the lost productivity due to workers absent from the job. Presenteeism means that the worker is in work but decreased production due to his illness." (C. Jones, Verstappen, & Payne, 2019). These can affect productivity in altering pathways. Typically, absenteeism is easy to identify and just count the absent day but presenteeism

comprises a more complicated model (1) identifying and measuring the unproductive time and (2) valuing the impact of that unproductive time. One self-reported presenteeism study indicates that a decrease in productivity related to MSD(C. Jones, Payne, Gannon, & Verstappen, 2016).



2.5.1 Conceptual model of work productivity

CHULALONGKORN UNIVERSITY

The above figure is the conceptual model of work productivity that is related to MSD. The leftmost box is the health condition parameter or health risk that is associated with work productivity(WHO, 2001).WMSD is the condition that is associate with pain which is regarded as the common indicator. The inverse association between illness and work productivity is taken as an important aspect to measure absenteeism, presenteeism, and work disability. Moreover, a healthier person can be more productive(Burton, Conti, Chen, Schultz, & Edington, 1999).In other words, the increased risk of ill health in a job is related to decrease productivity and lead to psychosocial stress (Gunnar Aronsson, Gustafsson, & medicine, 2005).

The second box is the concept of ability, capacity, and desire to work that is related to worker performance. A healthy working environment is an important factor for job performance and also a major contributor to absenteeism and presenteeism. WMSD can affect the worker's performance as a mental aspect and in turn later impair work productivity (Burton et al., 1999; Lowe, 2003).

The final box is work-life balance. It means that the non-work factors, social life is also related to productivity. Not only the work factors but also the social life outside the working environment affect the work productivity and these are also regarded as extra work factors that influence productivity (Hagberg, Tornqvist, & Toomingas, 2002). E.g. when a nurse working the night shift and picking a child from a daycare center and equal to non-work factors such as leisure activity.

หาลงกรณ์มหาวิทยาลัย

Observed work productivity (OWP) means that we can count the number of units produced per working day and services that they give to the customers. A study by (Meerding et al., 2005) use OWP by counting output from construction workers. They used the size of the street (''road pavers") and floor surface (''floor layers'') per day was observed as output productivity. However, there are many limitations to use OWP in different occupational sectors. It is only can be used in certain groups of working sectors such as customer service and negotiators that work through telephones, or workers in assemble parts (Escorpizo, 2008). Perceived work productivity (PWP) is regarded as the most useful tool to measure productivity when there is no OWP(Kessler et al., 2004). It is self-reported work activity that comprises presenteeism and absenteeism. Also (Escorpizo et al., 2007) explained the concept of work productivity that is used in musculoskeletal diseases and arthritis.

In the UK, Stress, depression or anxiety, and musculoskeletal disorders accounted for the majority of days lost due to work-related ill health, 17.9 million and 8.9 million respectively (HSE, 2020). Workplace health promotion programs that can positively affect the presenteeism and reduction of other factors are also encouraged by employers to increase productivity and further economic evaluation may need for an employer to decide whether to invest in this program (Cancelliere, Cassidy, Ammendolia, & Côté, 2011).

2.5.2 Relationship between psychosocial factors and work productivity

หาลงกรณมหาวทยาลย

Working conditions, stress, Job conditions, and workers' characteristics all significantly influence work productivity. Productivity loss is influenced by those factors directly or indirectly through health impairment. People with high job demand, good social support, and high intrinsic, extrinsic reward are more likely to perceive their products as healthy men than those who lack these conditions. (Lowe, 2003) The study showed that workers who are satisfied with their working environment including psychosocial conditions are more likely to better work morale, lower absenteeism, rarely quit. Adversely, those who reported that their job is stressful can have a higher level of presenteeism and absenteeism. (Allen, 2008) Psychosocial factors and workers' health is related to production cost. The add up cost related to sick leave, the replacement cost of labor and compensation can go up when the workers suffer turnover of psychosocial conditions. The workplace protective factor is Psychosocial Safety Climate (PSC) which is aimed to prevent and manage workplace psychosocial hazards. PSC concern with how workers value psychological health, commit and support the psychological health problems, prioritize the burden of psychological health of workers as aspect to profit and productivity.(Hall, Dollard, & Coward, 2010) The Australian research reported that depression costs approximately AUD 8 billion per annum due to productivity loss (.5 percent GDP) because of sickness absence and presenteeism (i.e., reduced performance at work) (McTernan, Dollard, LaMontagne, & Stress, 2013). The research paper of (Adib Ibrahim et al., 2019) revealed that psychosocial stressors can explain more than 50% of the variance in health care productivity. The most important factors are job influence, job role clarity, reward, and job satisfaction that is linked with productivity. Also study of private organizations in Nigeria finding reported that psychological factors have a significant contribution to perceived workers 'productivity that is related to workers efficiency in managing resources (Oyebamiji & Akintayo, 2011). Therefore, we need to better understanding of the link between psychosocial factors and productivity and also find the relationship with WRMSD to get better achievement of worker health.

2.5.3 Sustainable Development Goal (SDGs) and productivity

According to ILO, major SDG goals that are linked with workers' productivity are

- SDG 2.3 ("By 2030 double the agricultural productivity and the incomes of small-scale food producers, particularly women, indigenous peoples, family farmers, pastoralists and fishers, including through secure and equal access to land, other productive resources and inputs, knowledge, financial services, markets, and opportunities for value addition and non-farm employment")

- SDG 8.2 ("Achieve higher levels of productivity of economies through diversification, technological upgrading, and innovation, including through a focus on high value-added and labor-intensive sectors")

- SGG 9.5 ("Enhance scientific research, upgrade the technological capabilities of industrial sectors in all countries, in particular developing countries, including, by 2030, encouraging innovation and substantially increasing the number of research and development workers per 1 million people and public and private research and development spending")

Working poor phenomenon such as long working hours, working in informal sectors are the major causes of productivity. Raising productivity increases higher profit in business owners, investors, and then for the workers, the result leads to higher wages and improve the working environment, more decent work, reducing poverty. Therefore, they have circled chains linked with productivity, wage, and poverty. Governments, employers, and workers all unite together to enhance productivity because it is the primary contributor to increase living standards, to fight off working poverty. Moreover, garment sectors include in one of the ILO's five flagship programs cooperation Better Work Organization, which aim to improve working conditions and more compliance with labor standards. (ILO, 2021a)

2.6 Methodology Explanation

2.6.1 Cornell musculoskeletal Discomfort Questionnaire for WRMSD

This is the questionnaire developed by Professor Alan Hedge and Ergonomics students of Cornell University developed a well-designed data collection tool named CMDQ to access the WRMSD. It comprises a 54 item questionnaire for 18 body parts to access the musculoskeletal discomfort, pain, and aches over the last week. Scoring will be done by simply counting the numbers of people that suffer from WRMSD (Hedge, Morimoto, & Mccrobie, 1999). It also developed different questionnaires for the sedentary worker, standing workers, hand symptoms, and different gender. Cornell musculoskeletal questionnaire is more updated, more widely used, and more specific than other tools. Moreover, there are also research articles that CMDQ is fit for sewing operators and also useful in ergonomic evaluation (Nagaraj & Jeyapaul, 2018; Nagaraj, Jeyapaul, & Mathiyazhagan, 2019).

2.6.2 Work productivity and Activity Impairment Questionnaire (WPAIQ) for work productivity

This is the questionnaire developed by Margaret Reilly, President of Reilly Associates since 1989. Until the early 1990s, there was no quantitative measure of health-related work productivity loss for the employed population, i.e., there was no instrument that assessed the amount of both absenteeism (work time missed) and presenteeism (reduced on-the-job effectiveness) due to health problems. Therefore, they decided to develop tools for these measurements (Reilly, Zbrozek, & Dukes, 1993). The above literature mentions that presenteeism and absenteeism are major contributors to productivity that is related to WRMSD.WPAIQ is also mainly based on these two variables also can access the activity impairment that is not related to work. WPAIQ also developed access work productivity that is associated with certain kinds of diseases such as asthma, rheumatoid arthritis. They are also been translated to 140 languages and many validity and reliability test are done in different languages (ASSOCIATES, 2019).

2.6.3 Job Content Questionnaire (JCQ) for psychosocial factors

This is the questionnaire developed by Professor Karasek based on the Demand control support model explained in the literature part. It is based on the world's know stress theory that measures the psychosocial factors in all kinds of jobs. It covers a wide variety of job characteristics such as physical demand, psychological demand, job insecurity, social support. Moreover, the instrument is widely used in academic and University-based research and has focused on the worker's task only. It is a short, efficient, and self-administered questionnaire in about fifteen minutes

2.6.4 Work Ability Index (WAI) for work ability

The WAI questionnaire has been developed, based on the balance model of the stress-strain concept, by the Finnish Institute of Occupational Health (FIOH) in the early 1980s. WAI is widely applied for conceptualizing the work ability by obtaining information related to diseases, functional limitations, sick leave, and mental resources. (Nygård, Savinainen, Kirsi, & Lumme-Sandt, 2011; T. I. van den Berg et al., 2008).WAI is now widely used in an international research setting and translated into 30 different languages. (J. J. S. j. o. w. Ilmarinen, environment & health, 2009) .This is the occupational and clinical research instrument that is used in workplace surveys and access work ability that is also related to health. This is the series of the questionnaire that is used to determine the score on basis of answers and contain the demands of work, the worker's health status, and resources. (J. J. O. m. Ilmarinen, 2007)

CHAPTER 3

RESEARCH METHODOLOGY

3.1 Study Design

This study is aimed to find the association of psychosocial factors, work productivity, and musculoskeletal disease in sewing machine operators in a garment factory in Yangon, Myanmar. A cross-sectional method was used and sewing operators from respective garment factories were recruited. The interesting outcome may be psychosocial risk factors present (or) not, work productivity decrease (or) not, musculoskeletal disorders present (or) not, work ability impaired or not. Data collection was done by self-answered questionnaires.

3.2 Study Area

The study area was in Yangon, Yangon Region, and Myanmar which is one of the most industrialized areas in Myanmar and continues expanding. In the Yangon region, there are a number of industrial zones/areas situated in Hlaingtharyar, Shwepyithar, Shwepaukkan, and Mingaladon townships. Moreover, these areas are also the major employment sector in Yangon. During the semi-lockdown period due to the surge of the third wave of COVID-19 pandemic in the Yangon region, most of the industrial zones/areas were closed. Among them, I had to choose Shwepaukkan industrial zone purposively because it is also recognized as one of the densest purposes. Most of them are foreign owners that were invested in Myanmar. More importantly, it was the only industrial zone/area that allowed me to survey a limited number of garment factories among those which were still open and the managerial bodies were willing to accept my study in their factories. Therefore, I chose the sample factories from that study area because of the above reasons.





Figure 5.Industrial zone map of Yangon

3.3 Study population and inclusion, exclusion criteria

Participants were the person working as sewing machine operators roughly 4000 participants in the Shwepaukkan Industrial area, Yangon. Inclusion and exclusion criteria will be provided to the factory and they will perform these criteria and select research participants. Moreover, during recruitment, before explaining the research process, and questionnaire, confirm again these criteria by the researcher.

Inclusion criteria

- Those who worked at least 1 year as sewing operators
- Those who are equal or more than 18 years of age
- Those who are free from disability
- Those who are willing to participate
- Those who can read and write

Exclusion criteria

- Those who suffered from injury, accident, or undergo surgery during the last 3

months

- Those who were pregnant (female respondent)

- Those who were taking any prescribed medicine by self (or) from a doctor

3.4 Sample Size Calculation

The estimated sample size was calculated from the following formula. The formula would be used for calculating the adequate sample size in the prevalence study. (Charan & Biswas, 2013; Daniel & Cross, 2018; Pourhoseingholi, Vahedi, & Rahimzadeh, 2013)

$$n = \frac{(Z\alpha/2)^2 \times p \times (1-p)}{(d)^2}$$

n= the estimated sample size

 α = 0.05 (the level of statistical significance)

 $(Z\alpha/2)^2$ = the value from normal distribution associated with a confidence interval

(1.96 with 95% confidence interval)

P= 0.5 (maximum number of prevalence decided by a researcher is used in this study

because previous research prevalence is not relevant to this study)

d =0.05 (the value of maximum allowable error or desired precision)

Then, the sample size is calculated as follow:

n =
$$\frac{(1.96)^2 \times 0.5 \times (1-0.5)}{(0.05)^2}$$
 = 384

Therefore, the minimum sample size for this study will be 384. To added to cover incomplete questionnaires or any missing data and withdrawal. Therefore, the total required sample size will be approximately 400.

3.5 Sampling Method

Among the most industrialized region in Myanmar, purposively select the Yangon region. Again in the Yangon region, the Shwepaukkan Industrial zone/area was chosen. There is a total of 15 garment factories. Then using the stratified method, select large garment factories which a population is more than 1000. Due to the limitation of the semi-lockdown period during the third wave of the COVID-19 pandemic in Myanmar, only 2 factories agreed to be participated after approaching all factories in that industrial zone/area. 400 sample size is used according to calculation and then select 200 sample sizes from each factory. The sampling method is multi-stage sampling. In each factory, 200 participants will be selected by systematic random sampling. A list of participants will be provided by the factory. Participants with odd IDs will be included until reaching the 200 sample size.

After getting the list of workers, researcher recruit and explain about the research, benefit, and all the details about the questionnaire. All are free to participate and get informed consent.



Figure 6. Sampling frame

3.6 Measurement Tools

The self-administered questionnaire was used to conduct the study. The questionnaire was prepared in the English language first and then translated into the local language, Myanmar by a language expert and then back-translated to English by the occupational and Environmental Health Specialist. The disputes between the two translators was decided by the main researcher. The questionnaire contains five parts which contain 66 including questions, items, and statements. The personal factors questionnaire contains 7 questions. CMQ contains 20 questionnaire. JCQ contains 27 questionnaire. WAI contain 7 item questionnaire and WPAI contains 5 questionnaire. These are as follow.

3.6.1. Personal factors questionnaire

This includes the baseline characteristic of the sample, including age, sex, education, marital status, height and weight, working hour per day, BMI, working experience, monthly income of the study participant.

3.6.2 Work Ability Index (WAI)

It measures the working ability of the workers using a seven-item questionnaire. It includes current work ability compared with the lifetime best (item 1, 0-10), work ability concerning the demands of the job (item 2, 2-10), number of current disease groups diagnosed by a physician (item 3, 1-7), estimated work impairment due to diseases (item 4, 1-6), sick leave during the past year (item 5, 1-

5), the personal prognosis of work ability for 2 years from now (item 6, 1,4 or 7) and mental resources, referring to the worker's life in general, both at work and during leisure time(item 7, 1-4). The number in parentheses for each item indicates the scoring range. The total WAI score is calculated by summing up the scores of all items and is ranges from 7 to 49. The total WAI scores are categorized into 4 levels: poor (7-27), moderate (28-36), good (37-43), and excellent (44-49). (J. J. O. m. Ilmarinen, 2007). For internal validity, predictive validity, and reliability now documented subjective assessment of work ability using the WAI questionnaire seems to provide a good instrument and test-retest reliability. (Cordeiro, Souza, & Araújo, 2017; de Zwart, Frings-Dresen, & van Duivenbooden, 2002) Moreover, WAI has a high level of cross-national stability and reliability. (Radkiewicz & Widerszal-Bazyl, 2005). Cronbach's alpha in this study was >0.7 for each item.

3.6.3 Cornell Musculoskeletal Discomfort Questionnaire (CMDQ)

CHULALONGKORN UNIVERSITY

It examines the prevalence of musculoskeletal diseases in 20 body parts including left and right that were suffered during the previous week. Questionnaire for both male and female are same as only different is figure. It includes three sections (frequencies, severity, and inference). The validity of the CMDQ has not been formally tested in the US but it has been extensively tested by Dr. Oguzhan Erdinc in Turkey with good results (Erdinc, Hot, & Ozkaya, 2011). Many other crosscultural translation validities and reliability tests were done with a satisfactory outcome. (Afifehzadeh-Kashani et al., 2011; Kreuzfeld et al., 2016) T-CMQ Cronbach's alpha statistic for both three scales was 0.876, 0.895, and 0.875 respectively which indicated that internal consistency was high (Erdinc et al., 2011).

3.6.4. Work productivity and Activity impairment questionnaire-general health version (WPAI-GH)

This is the measurement for productivity and activity impairment in employees in the factory. WPAI-GH revealed the 4 types of scores related to Absenteeism (work time missed), Presenteeism (impairment at work / reduced on-the-job effectiveness), Work productivity loss (overall work impairment/absenteeism plus presenteeism), and Activity Impairment. The WPAI-GH consists of six questions: 1 = current employment; 2 = missed hours due to illness of health; 3 = missed hours due to other reasons; 4 = hours worked; 5 = health status affected productivity while working, 6 = health status affected productivity in regular unpaid activities. The duration for memory recall about the questions past seven days, not including today.

Four main generated outcomes from the WPAI-GH and described in percentages by multiplying the following scores by 100: (1) Absenteeism score = Q2/(Q2 + Q4) (2), Presenteeism score = Q5/10 both for those who were currently employed and worked in the past seven days(3) Overall work productivity loss = $Q2/(Q2 + Q4) + ((1 - Q2/(Q2 + Q4))) \times (Q5/10)$ for those who were currently employed; (4) percent activity impairment due to health = Q6/10 for all respondents. In, this study, we exclude the activity impairment part because all participants are currently working employees. This tool is translated into different languages and validity, reliability test is done and Cronbach's alpha statistic for each scoring is in >0.7. (Reilly et al., 1993). Cronbach's alpha in this study was calculated and it was >0.7 for each item.

3.6.5. Job content questionnaires (JCQ)

The psychosocial factors are measured by Job content questionnaire. In this study, we use 27 item questionnaires which include 5 scale sections, job control, psychological job demand, workplace social support, physical job demand, and job insecurity. Skill discretion, (evaluated by 6 items), and decision authority, (evaluated by 3 items) are sum up to measure the job control scale. The psychological job demand scale comprises 5 items. Supports from supervisors and from co-workers, (each includes 4 items) are combined to evaluate the social support scale. The physical job demand scale and job security scale are evaluated by 3 and 2 items each. The outcome response will be recorded for each questionnaire on a 4-point Likert-type scale, ranging from 1 (strongly disagree), 2 (disagree), 3 (agree), to 4 (strongly agree). The scales and sub-scales of Cronbach's alpha statistics (α) are also validated and range between 0.319-0.894. (Karasek et al., 1998; Sein, Howteerakul, & Jirachewee, 2010) .Cronbach's alpha in this study was calculated and it was >0.7 for each item.

3.7 Data Collection Process

Permission from the local authority and respective factories were asked. First, we select participants according to inclusion and exclusion criteria. The study procedure was explained to the participant and writing consent with signature was requested from each participant. A research team was built with one research assistant to facilitate explaining questionnaires. A research assistant will be a qualified graduate person who will relate to the medical field such as nurse or pharmacist, community health care workers. The appropriate time for data collection is also needed to be considered and requested from the factory authority, in order not to interfere with their production. Explaining about research questionnaire to the participant was done in the open area that will be provided by the factory. To avoid data contamination, collection was done only 50 samples per one day, and CMQ, JCQ, WAI, and WPAI are all explained and self-administered, set up to fill at their home, and collected the next day, allowing time for 24 hours.

3.8 Data Entry and Analysis

Statistical analysis for social science version 26 was used for data analysis. Descriptive statistics such as mean, median, standard deviation, percentage, minimum, maximum values are used for interpretation of the variable. The test of the level of statistical significance in this study is alpha (α) less than 0.05.

Table 1.Data analysis

Objectives	Variable	Statistic
-To find an association	Dependent variable	
between psychosocial	- work-related	
factors and work-related	musculoskeletal Disorders (Binary	
musculoskeletal Disorders	outcomes: "no musculoskeletal	
- To find an association	disorders at least one part of the	
between work productivity	body" / "musculoskeletal disorders	
and work-related	at least one part of the body")	
musculoskeletal Disorders	Independent variable	
- To find an association	- psychosocial factors/work	Independent T-test
between work ability and	productivity/work ability with	(Normally distributed
work-related	continuous data	data)
musculoskeletal Disorders	ลงกรณ์มหาวิทยาลัย เ	Mann - Whitney U
UTULA	- psychosocial factors/work	Test (Non-Normal
	productivity/work ability with	distributed data)
	categorical data	Pearson's Chi-square
-To find out the combined	Dependent variable	
effect of psychosocial	- work-related	
factors and work	musculoskeletal Disorders (Binary	

productivity on work		
productivity on work-	outcomes: no musculoskeletal	
related musculoskeletal	disorders at least one part of the	
Disorders	body" / "musculoskeletal disorders	
	at least one part of the body")	
	Independent variable	Binary logistic
	- combine effect of	regressions (Odd ratio
	psychosocial factors and work	and Adjusted Odd
	productivity	ratio** with 95%CI will
		be reported)
<u>الا</u>	Group 1 : poor psychosocial factors	**Adjusted with some
	x poor work productivity	confounding factors in
Sel.	and a marked and a m	general characteristic
-	Group 2 : good psychosocial factors	
จุหา	x poor work productivity	
Chula	LONGKORN UNIVERSITY	
	Group 3 : poor psychosocial factors	
	x good work productivity	
	Group 4 : good psychosocial factors	
	x good work productivity	
	Psychosocial factors are determined	
	by mean	

	lf > mean, High	
	lf< mean, Low	
	Overall Psychological factors are	
	determined by	
	-High Psychological job demand,	
	low job control, low social support,	
	poor psychosocial factors	
	- Low Psychological job demand,	
all a second sec	high job control, high social support,	
	low job insecurity is regarded as	
1 States	good psychosocial factors	
- 1011	Work productivity is determined by	
จุฬา	ลงกรณมหาวิทยาลัย	
Chula	mean percentage.	
	If > mean = Poor productivity	
	If < mean = Good productivity	
	The total WAI scores are categorized	
	into 4 levels: poor (7 - 27), moderate	
	(28-36), good (37-43), and excellent	

(44-49)	

3.9 Ethical Consideration

The research protocol was submitted and approval obtained by the Ethical Review Committee for Human Research, Faculty of Public Health, Chulalongkorn University before collecting data. Before the data collection, the right to participate, refuse, or withdraw was informed to the respondents. All have voluntarily participated in this study.


CHAPTER 4

RESULT

The objective of the study is aimed to determine the percentage of Workrelated musculoskeletal disorders, psychosocial risk factors, work productivity, work ability and to find the association with WRMSD to other factors. The study population was sewing machine operators who were more than 18 years of age and currently working in Shwepaukkan Industrial Zone, Myanmar.

There are five parts to the result section. Each section contains both descriptive and inferential statistics. The result of the study is presented as follows.

4.1 General characteristics of the participants.

4.1.1 Association of Personal factors and WRMSD

4.2 Percentages of WRMSD in sewing operators

4.3 Percentages of psychosocial risk factors

4.3.1 Association of Psychosocial risk factors and WRMSD

4.4 Work Ability Index Score of sewing machine operators

4.4.1 Association of Work ability and WRMSD

4.5 Work productivity of sewing machine operators

4.5.1 Association of Work productivity and WRMS

4.6 Binary logistic regression analysis of personal factors, psychosocial factors,Work ability, and work productivity with WRMS

4.1 General characteristics of the participants

The personal factors of garment factory workers were shown in Table 2. The table included information related to age, gender, education, marital status, monthly income, working hour per day, duration of work, and BMI.

The result showed that among 370 garment factory workers who participated in this study, the average age is 25.6 ± 4.6 years. The majority of the sewing workers (61.9 %) were in the age group 22-30 years and about 26.5 % were in the age below 22 years. The youngest responding garment workers in the study were 18 years old while the eldest was 51.Regards to gender, the majority of sewing operators were occupied by females about 95.4 % with 353 respondents.

For the educational level, secondary school level comprised majority with 54.6% followed by tertiary level and above 33.8% and primary school level 11.6%. Concerning the marital status of sewing operators, most of the portion was occupied by a single with 83.5% and then married was 13.5%. Regarding the income, the majority earn more than 200000 MMK with 65.9% and less than 200000 MMK is 34.1% with the mean value of 227151±61015. Working hour per day was cut off to 8

hours and most of the operators worked more than 8 hours with 60.8%. The average working hour was 9.25±1.3.

As for the working experiences, it was also divided to more or less than 5 years and the respondents' rate to more than 5 years is only 26.8%. The average means of working experience was $4.2\pm$ 3.2 years. If we looked at the BMI, Healthy BMI comprised with 61.6% majority followed by underweight 30.3%.

		n (%)
Age	<22 years	98 (26.5)
	22 to 30 years	229 (61.9)
จหา	>30 years	43 (11.6)
Gender CHULA	Female	353 (95.4)
	Male	17 (4.6)
Education	Primary school	43 (11.6)
	Secondary school	202 (54.6)
	Tertiary and above	125 (33.8)
Marital Status	Single	309 (83.5)
	Married	50 (13.5)

Table 2. General characteristic of participants (N=370)

	Divorced	5 (1.4)
	Widowed	6 (1.6)
Monthly	≤200000	126 (34.1)
income(Myanmar kyats)	>200000	224 (65.9)
Working hour per day	≤8 hr	145 (39.2)
	>8 hr	225 (60.8)
Duration of work as sewing	≤5 yrs	271 (73.2)
operators	>5 yrs	99 (26.8)
ВМІ	<18.5	112 (30.3)
	18.5 to 24.9	228 (61.6)
	25.0 to 29.9	23 (6.2)
จุหา	>30 ลงกรณ์มหาวิทยาลัย	7 (1.9)

Chulalongkorn University

4.1.1 Association of Personal factors and WRMSD

To find the association with WRMSD, p-values are also presented in Table 3 with chi-square statistical analysis. Most of the personal factors are not significantly associated with WRMSD except for age and education status. Age is significantly associated with a p-value of 0.010 and an educational level of 0.01. According to the results, \leq 30 years of age is 3.7 times more likely to develop WRMSD than >30years

of age with a p-value of 0.010. Similar to educational attainment, tertiary educational level workers are 12 times more likely to suffer WRMSD than primary and secondary education with a p-value of 0.002 as shown in Table 4.

		Work-related		p-value
		musculoske	letal	
		disorders		
		Yes	No	
		n Row%	n Row%	
Age	<22 years	91 (92.9)	7 (7.1)	0.011 * (f)
	22 to 30 years	220 (96.1)	9 (3.9)	
	>30 years	36 (83.7)	7 (16.3)	
Gender	Female	332 (94.1)	21 (5.9)	0.286 ^(f)
	Male_ALONGKO	15 (88.2)	2 (11.8)	
Education	Primary school	42 (97.7)	1 (2.3)	0.001 *(f)
	Secondary school	181 (89.6)	21(10.4)	
	Tertiary and above	124 (99.2)	1(0.8)	
Marital Status	Single	288 (93.2)	21 (6.8)	0.880 ^(f)

Table 3.Personal factors of sewing machine operators related to WRMSD (N = 370)

	Married	48 (96.0)	2 (4.0)	
	Divorced	5 (100)	0 (0)	
	Widowed	6 (100)	0 (0)	
Monthly	≤200000	117 (92.9)	9 (7.1)	0.651
income(Myanmar	>200000	230 (94.3)	14 (5.7)	
kyats)		11/122		
Working hour per	≤8 hr	135 (93.1)	10 (6.9)	0.665 ^(f)
day	>8 hr	212 (94.2)	13 (5.8)	
Duration of work	≤5 yrs	258 (95.2)	13 (4.8)	0.086
as sewing			3	
operators	>5 yrs	89 (89.9)	10 (10.1)	
BMI	<18.5	6 (5.4)	106 (94.6)	0.661 ^(f)
	18.5 to 24.9	17 (7.5) มหาวิทยา	211 (92.5)	
	25.0 to 29.9	ION UNIVE	23 (100)	
	>30	0	7 (100)	

f = fisher exact test

Table 4.Association	of Personal	factors an	id WRMSD
---------------------	-------------	------------	----------

X ²	Crude	95% CI	p-value
	OR		

>30 years	8.451	3.780	1.458 – 9.801	0.010*
≤30years				
Primary and secondary school	9.499	12.233	1.639 – 91.850	0.002*
Tertiary and above school				
v.	14000	122 -		
	Q			

Reference group - ≤30 years of age, tertiary and above school

4.2 Percentage of WRMSD in sewing operators

370 respondents of the Standard Cornell Musculoskeletal Discomfort questionnaire were administered. The following Table 5 represents the percentages of WRMSD. Sewing operators reported that pain in at least one part of the body is 347 (93.8%) and mostly occupied by a female. Most common symptoms areas are in upper back (34.3%), neck (27.6%), right shoulder (24.9%) right lower leg (23.8%) and lower back (21.9 %)

Table 5. Musculoskeletal disorders' prevalence at the recent week in our subjects with musculoskeletal disorders (N = 370)

Body Region	Frequency	Percentage
Neck	102	27.6

Shoulder (Right)	92	24.9
Shoulder (Left)	80	21.6
Upper back	127	34.3
Upper arm (Right)	70	18.9
Upper Arm (Left)	56	15.1
Lower back	81	21.9
Forearm (Right)	52	14.1
Forearm (Left)	39	10.5
Wrist (Right)	49	13.2
Wrist (Left) จุฬา	29 ลงกรณ์มหาวิทยาลัย	7.8
Hip CHULA	68 GKORN UNIVERSIT	18.4
Thigh (Right)	68	18.4
Thigh (Left)	49	13.2
Knee (Right)	73	19.7
Knee (Left)	45	12.2



จุหาลงกรณ์มหาวิทยาลัย

4.3 Percentages of Psychosocial risk factors

Regards to psychosocial factors according to Karasek Demand Control model of Job strain. Job Demand Control Model measures psychological job demands, job control, workplace social support, physical demands, and job insecurity. It is the modification of Karasek job Content Questionnaires and all these items measure psychosocial hazards. Table 6 shows that 41.4% have reported high psychological demand. About 29.5% of the respondents had a low score of supervisor support and 90.3 % had a high score of co-worker support. But overall 77.0 were reported high workplace support. About 75.9% of the respondents had a high score of physical demands, 55.6% has high job control, and 53.8 % response with high job insecurity. To sum up, low psychological demand, high social support, and high job control, high job insecurity are obtained. However, most of the sewing operators reported that their job requires high physical demand.

Table 7 represent the percentage of WRMSD in accordance with psychosocial factors. The total percentage of WRMSD pain in at least one part of the body is higher in low psychological demand (94.5%) than high psychological demand (91.5%). A similar result can be seen in other psychological factors such as workplace social support, coworker support, and job security. However, a nearly similar percentage of WRMSD pain was reported in job control and supervisor support with high (93.2%), low (94.5%), and high (93.9%), low (93.6%). In addition, high physical demand was recorded more percentage of WRMSD with 95.7% than low demand.

The Median cut	High	Low
off point	(n %)	(n %)

Table 6. Psychosocial risk factors (N = 370)

Psychological job demand	31	High (≥31)	Low(<31)
		153 (41.4)	217 (58.6)
Job control	61	High (≥61)	Low(<61)
		205 (55.4)	165 (44.6)
Workplace Social support	24	High (≥24)	Low(<24)
		285 (77.0)	85 (23.0)
Supervisor support	12	High (≥12)	Low(<12)
		261(70.5)	109 (29.5)
Coworker Support	12	High (≥12)	Low(<12)
		334 (90.3)	36 (9.7)
Physical demand	6	High (≥6)	Low(<6)
จุฬาส	งกรณ์มหาวิท	281(75.9)	89 (24.1)
Job insecurity CHULA	⁸ NGKORN UNIV	High (≥8)	Low(<8)
		19 (53.8)	171 (46.2)
Job strain		High	Low
(high psychological			
demand, low job control)		75(20.3%)	295(79.7%)

	High		Low	
	(MSD)Yes	(MSD) (No)	(MSD)Yes	(MSD) (No)
	n %	n %	n %	n %
Psychological job	140 (91.5)	13(8.5)	207 (95.4)	10 (4.6)
demand				
Job control	191 (93.2)	14 (6.8)	156 (94.5)	9 (5.5)
Workplace Social support	265 (93.0)	20 (7.0)	82 (96.5)	3 (3.5)
Supervisor support	245 (93.9)	16 (6.1)	102 (93.6)	7 (6.4)
Coworker Support	311 (93.1)	23 (6.9)	36 (100)	0 (0)
จุห'	เลงกรณ์มห	าวิทยาลัย		
Physical demand CHUL	269(95.7)	12 (4.3) RSN	78 (87.6)	11(12.4)
Job insecurity	158 (92.4)	13 (7.6)	189 (95.0)	10 (5.0)

Table 7.Psychosocial risk factors and WRMSD according to category % (N = 370)

4.3.1 Association of Psychosocial risk factors and WRMSD

The association between psychosocial risk and WRMSD is calculated according to the chi-square test. There is no significant association between psychosocial risk and WRMSD except physical demand a shown in Table 8. Moreover, job strain also found no significant association between high and low. Physical demand is significantly associated with WRMSD with a p-value of 0.010. According to the statistical result, high physical demand is 3.161 times more likely to develop WRMSD than low physical demand, and 95 % confident to say that the range is 1.343 – 7.441.

		X ²	Crude OR	95% CI	p-value
Psychological job	Low	2.327	0.520	0.222 – 1.219	0.188
demand				-	
	High	เงกรถ	เมหาวิทยาส	18	
C	HULAI	.ONGK	DRN UNIVER	SITY	
Job control	Low	0.296	0.787	0.332 – 1.867	0.668
	High				
Workplace Social	Low	0.011	0.485	0.140 - 1.673	0.312
support					
	High				
Supervisor support	Low	0.011	1.051	0.420 - 2.631	1.000

Table 8. Psychosocial risk factors and association with WRMSD

	High				
Coworker Support	Low	2.643	1.116	1.077 – 1.156	0.148
	High				
Physical demand	Low	7.586	3.161	1.343 - 7.441	0.010*
	High		NI/12		
Job insecurity	Low	1.048	0.643	0.275 – 1.506	0.389
	High				
Job Strain	Low	0.513	1.422	0.541 – 3.741	0.432
	High			3	

Reference group - High WIANDSOLWHIDDE

Chulalongkorn University

4.4 Work Ability Index score of sewing machine operators

Table 9 shows the individual item score of the WAI and the range score that was obtained from sewing operators according to scoring criteria. The average mean score is 34.3. Table 10 show the group criteria, categorized into four groups, and moderate work ability occupied in the majority of the respondents with 66.5 % which is needed to be improved followed by poor work ability 18.4 % which is needed to be restored. Moreover, good work ability is 12.7% and only 2.4 % reported excellent ability score.

Table 9. WAI individual item score (N = 370)

<u>ltem</u>	<u>Range</u>	<u>Mean ± SD</u>
1. Current work ability compared with the lifetime best	2 - 10	5.9 ± 1.3
2. Work ability with the demands of the job	4 - 10	6.9 ± 1
3. Number of current diseases diagnosed by a physician	1 - 7	6.4 ± 1.1
4.Estimated work impairment due to diseases	1 - 6	3.4 ± 2.1
5. Sick leave during the past year	2 - 5	4.7 ± 0.6
6. Own prognosis of work ability two years from now	1 - 7 RSITY	4.4 ± 1.6
7.Mental resources	1 - 4	2.6 ± 0.5
Total score	7 - 49	34.3

Table 10.WAI category score (N=	370)
-------------------------------	----	------

	n	Percentage
Poor (7 – 27)	68	18.4
Moderate (28 – 36)	246	66.5
Good (37 – 43)	47	12.7
Excellent (44 – 49)	9	2.4

4.4.1 Association of Work ability and WRMSD

The result of Table 11 explained the association of work ability and WRMSD. Poor work ability respondents reported more WRMSD percentage (95.6%) than poor work ability (91.9%). In addition, there are no reported cases of (No) WRMSD in good and excellent work ability. There is no significant association between WRMSD and work ability at the p-value of 0.166.

Table 11.WAI score and association with WRMSD

	(MSD)(Yes)	(MSD) (No)	X ²	df	p- value
	N %	N %			
Poor (7 – 27)	65(95.6)	3(4.4)	5.637	3	0.116
Moderate (28 – 36)	226(91.9)	20 (8.1)			

Good (37 – 43)	47 (100)	0 (0)		
Excellent (44 – 49)	9 (100)	0 (0)		

4.5 Work productivity of sewing machine operators

There is no significant reported absenteeism score. Only 13 respondents show absenteenism hours and show 0 percent according to WPAI calculation. The presenteeism score is about 34% and which can be considered as less impairment and good productivity. Overall work impairment is the sum of presenteeism and absenteeism. Therefore, The overall work impairment is the same as presenteeism as seen in Table 12.The total working hour during past 7 days 51±7.7. Impairment while working due to health is 3.39 according to WPAI calculation as mentioned in Chapter 2 and convert into percentage, therefore it is 34%.

Overall Activity impairment is excluded because that item only measures in non-working groups and this study, all are working groups.

Items	Unit Expression	WAI score with	p-value
		the	
		percentage	

Table 12. Work productivity score (n = 370)

Hours missed due to	Hr	N (%)	0 percent	0.150
health problem (>0 hr)	2Hr	3 (0.8)		
(Absenteeism)	4Hr	6 (1.6)		
	5Hr	1 (0.3)		
	8Hr	2 (0.5)		
	24Hr 8	1 (0.3)		
Hours actually work in				
nours actually work in	Mean= 51.71	II, SU = 1.1		
	Minimum=40l	nr,		
	Maximum= 70	0 hr		
Impairment while working	Mean= 3.39	, SD = 1.9	34 %	
due to health	Minimum = 0	, Maximum=	ΓY	
(Presenteenism)	8			
Overall work impairment			34 %	
due to health				
(Presenteeism				
+Absenteeism)				

4.5.1 Association of Work productivity and with WRMSD

To find the association with work productivity and WRMSD, productivity score is categorized into two groups (<50% and >50%). After running the chi-square test, there is no significant difference between work productivity and WRMSD with a pvalue of 0.150 as shown in Table 13.

			X ²	Crude OR	95% CI	p-value
		14	111			
Work	productivity	≤50%	2.192	0.382	0.102 - 1.430	0.150
score		>50%				

Table 13.Association of Work productivity and WRMSD

Reference group - >50%

4.6 Binary logistic regression analysis of personal factors, psychosocial factors,

Work ability, and work productivity with WRMSD

A binary logistic regression is carried out to determine the predictors of WRMS with controlling cofounders. Out of 17 variables, only 12 variables were selected as the most important variables to be entered in the logistic model. Hosmer-Lemeshow (Goodness of fit) text was used and assumptions for some variables show linear correlation with WRMSD. Only variables that meet assumptions are put into the model. Table 14 shows the final logistic regression model. The variables that were retained in the final model were age, gender, education status, duration of work, psychological demand, job control, physical demand, job insecurity, work productivity score, and work ability index score.

The result shows that there is a significant difference between gender, physical demand, job stress, and work productivity with WRMSD. Moreover, these are the significant predictors of a regression model. Male is 87% less likely to develop than female with a p-value of 0.046, 95% CI (0.020-0.969). Concerning high job stress is 8.2 times likely to develop WRMSD than low stress with a p-value of 0.017, 95% CI (1.456 - 46.550). Also, high physical demand is 4.7 times more likely to develop WRMSD than low demand with a p-value of 0.029, 95% CI (1.172 – 18.562). A similar situation is seen in work productivity percentage, the \geq 50 % productivity impairment is 5.8 times more likely to develop than the < 50% with a p-value of 0.016, 95% CI (1.393 – 24.920). Other variables are not significantly associated with WRMSD in regression analysis.

Chulalongkorn University

Table 14.Binary logistic regression analysis of personal factors, psychosocial factors, Work ability, and work productivity with WRMSD

Variable	В	S.E	Wald	df	Sig	Exp (B)	959	% CI
							Lower	Upper
Age	-1.534	0.798	3.693	1	0.055	0.216	0.045	1.031
Gender	-1.980	0.994	3.968	1	0.046*	0.130	0.020	0.969

80

Education	-2.218	1.157	3.676	1	0.483	0.588	0.133	2.598
status								
Duration of	-0.611	0.680	0.807	1	0.369	0.543	0.143	2.058
work								
Physical	1.548	0.709	4.769	1	0.029*	4.702	1.172	18.862
demand		1000	11100					
Job insecurity	1.229	0.886	1.922	1	0.166	3.418	0.601	19.425
Total social	-0.451	0.820	0.302	1	0.582	0.637	0.128	3.177
support	نظ			Va				
		AL COL						
Job stress	2.111	0.882	5.724	1	0.017*	8.257	1.465	46.550
				No.				
Work	1.774	0.736	5.813	1	0.016*	5.893	1.393	24.920
productivity %	จุหา	ลงกรณ	มหาวิท	ยาล์	٤			
productivity /0	CHULA	LONGKO	orn Uni	VERS	SITY			
(<50% and								
≥50%)								
WAI	0.599	0.994	0.402	1	0.526	1.820	0.286	11.588

Note – reference groups are \leq 30 years of age, female, primary education, \leq 8 hrs of work per day, low physical demand, low job security, low social support, low job stress, < 50% work productivity impairment, poor work ability score.

CHAPTER 5

DISCUSSION

The prevalence of WRMAD among garment workers is becoming a wider occupational problem. They are the basic workers and garment factories are one of the major sectors that give employment to Myanmar people. Moreover, young age is more involved in industrial work as sewing operators and also facing physical, mental, and behavioral problems but also cause many job-related sickness and injury and make them low performance, poor job satisfaction, and poor quality of life. Much research is studied on WRMSD in developing countries over the recent decade. However, studies on WRMSD, psychosocial factors, work productivity, work ability, published research focusing on sewing machine operators are very limited in Myanmar. The workforce of garment sectors still growing in Myanmar and if major occupational problems are left undiagnosed, there will be an impact on the country's economy.

This study of "Work-related musculoskeletal disorders, psychosocial factors, work productivity, and work ability among Garment factory workers in Myanmar" is conducted to find the percentage of productivity, work ability, psychosocial factors, and percentage of work-related Musculoskeletal Disorders of sewing machine operators in the garment factory. The results can provide information about the situation of productivity, work ability, psychosocial factors, and WRMSD and appropriate measures can be implemented based on the results of this study. In this chapter, a discussion of the result findings and a comparison of the previous research studies will be presented.

5.1 Prevalence of WRMSD

WRMSD has become a major occupational problem in the recent decade due to its growing burden not only its negative impact on the performance of workers but also on the well-being of workers.

Due to the limited number of previous studies on WRMSD in Myanmar, it was not possible to cite the literature that has the same criteria to define "Work-related musculoskeletal disorders" to obtain the appropriate comparison with the results of the present study. However, the prevalence of sewing machine operators and other factors that were studied in different countries are discussed. Moreover, most of the studies are focused on all sectors of Garment factory workers. But in this study, the **Chulcionecon University** discussion will only be done on studies focus only on sewing operators. The prevalence of WRMSD in this study is about 93.8 %. This finding is also similar to the study of the prevalence, of Musculoskeletal Disorders among Sewing Machine Operators in Nigeria about 92 % (Ak, Ba, & Adebisi, 2013). The finding is higher than in the study of sewing machine operators in Sir Lanka which the prevalence of WRMSD is about 81% (Nagaraj et al., 2019) and another study in Nigeria with 69% (Akinpelu, Oyewole, Odole, & Ogunbamowo, 2016). Moreover, the study of WRMSD of sewing operators in India is far lower percentage about 43% and the reason can be due to Indian study contains only 60 participants.(E. Mehta, Mehta, Sharma, & Wellbeing, 2020) In addition, there are very few studies on sewing machine operators using 7 day prevalence of CMDQ. Most of the studies are using Nordic Musculoskeletal Questionnaire with a 12-month prevalence. As compared to the 12-month prevalence, this study is consistent with the findings of Iranian sewing operators with a prevalence rate of 91% (Dianat et al., 2015)and in Estonia study 91.2% (Merisalu, Mannaste, Hiir, & Traumann, 2016).

Most common symptoms areas are in upper back (34.3%), neck (27.6%), right shoulder (24.9%), right lower leg (23.8%) and lower back (21.9%). The top three pain areas are consistent with the finding of sewing machine operators in the textile manufacturing industry in Botswana with the upper back (32.5%), Shoulder (21.7%), and neck (15.4%) (Sealetsa & Thatcher, 2011). These results are consistent with other studies that have highlighted problems in the shoulder and neck regions (J. H. Andersen & O. J. A. j. o. i. m. Gaardboe, 1993; Blåder et al., 1991). A large study of upper extremities WRMSD of sewing machine operators in Los Angeles, California reported that neck and shoulder mostly occupied musculoskeletal pain with 24% which is consistent with these findings (P.-C. Wang et al., 2007). Moreover, the most common pain areas are in accordance with the study of sewing machine operators in turkey reported that upper trunk pain (upper back and lower back) was 62.5.%, neck 50.5% and shoulder 50.2% (Öztürk & Esin, 2011). However, most of the studies are also reported that the low back is one of the most common pain in sewing machine operators and then followed by neck and shoulder. (Dianat et al., 2015; Nagaraj et al., 2019).In addition, a self-reported study of sewing operators in Nigeria also stated that 7 days prevalence of body pains are commonest in the lower back (31.6%), neck (21.3%), and shoulder (19.0%)(S. Maduagwu et al., 2015). Nevertheless, these three body regions- back, neck, and shoulders are frequently affected because the sewing operation is characterized by a static sitting posture for a long time, head and trunk forward inclined posture, and relatively uncomfortable ankle and knee angles (P olajnar, Leber, & Herzog, 2010).

The prevalence of WRMSD and most pain anatomical sites among sewing machine operators can be varied in different countries from different studies. The variation of the research can be due to different working environments, different geographical areas, traditions and culture, and differences, different research methodologies, sample size and population, duration of the study, study period. The instrument CMDQ is also accessed musculoskeletal pain based on their perceptions of workers. As it is the subjective measure of musculoskeletal pain, it can provide different findings in different countries. It is noted that WRMSD can be influenced by other factors such as job stress, increased demand from the competitive economy, changes in the working environment and labor market. Despite being the same nature of work, improvement and sustainability of working environment among factory workers are different in Myanmar compared to other countries. As weak in occupational health sectors and labor law, and Myanmar sill developing countries, therefore most of the workers suffer more occupational problems compared to others.

5.2 Personal factors of sewing machine operators related to WRMSD

Regarding the age of the sewing machine operators, \leq 30 years of age is 88.4% and > 30 years is 11.6%. In an analysis of this study, there is a significant relationship between age and WRMSD, and younger age groups are more reported than older age groups. Perhaps, this can be the younger population age of sewing machine operators. Sewing task requires more physical effort, repetitive job and younger age are more fit in sewing operators. Moreover, Myanmar is still a developing country and the major employment sector is provided by garment factories. Due to the poverty in the country, the younger age group has to work for their family earning (Burnley, December 2015; Marian Boquiren, 2019). This finding is in line with the research paper of sewing operators related to WRMSD in Nigeria which also reported that younger age is more prevalent of WRMSD than older(S. Maduagwu et al., 2015). Moreover, other research findings also support this result regards to young age groups(S. M. Maduagwu et al., 2014; Roquelaure et al., 2012). However, there is also a contradictory report that increasing age is also associated with the prevalence of WRMSD (Aghili et al., 2012). In addition, major contributory factors that high prevalence in the younger age group can be due to lack of skill and knowledge, little experience in the work, and increase workload (Roquelaure et al., 2012).

This study also find that education is another associated factor related to WRMSD. Higher educational attainment is more likely to suffer than low education and very few studies support this relationship. (Kaka et al., 2016) found that low back pain is a significant association between low back pain and high educational attainment. Another study conducted in Japan also revealed that only ankle pain is associated with high educational attainment (Tokuda et al., 2007). But these studies did not state possible explanations on high education and WRMSD. This can be a perhaps huge different educational level of sewing operators in Myanmar and this can lead to a significant association. Most are primary and secondary level education and very few percentages are in high education (Marian Boquiren, 2019). One study suggested that low education was associated with a lower level of active coping skills against musculoskeletal pain events encountered in the workplace (Carroll, Mercado, Cassidy, & Cité, 2002). However, many kinds of research about education and WRMSD found no significant association between them (Dianat et al., 2015; Nagaraj et al., 2019; Öztürk & Esin, 2011). Moreover, in some research findings, lower educational attainment is a risk factor for WRMSD and many articles are contradictory about these associations (J. H. Andersen & O. J. A. j. o. i. m. Gaardboe, 1993; Ming & Zaproudina, 2003). Therefore, further studies are needed to conclude the clear-cut association and mechanism pathway between WRMSD and educational level.

Gender is no significant association with WRMSD. The possible explanation is that the majority of respondents are female in sewing machine operation and cannot conclude clearly that there is no association. Marital status, duration of work, working hours per day, income, and BMI also did not show a significant association. These findings are also consistent with the study of ergonomic risk factors of sewing operators in Turkey (Öztürk & Esin, 2011). As regards marital status, some studies find no association (Amin et al., 2014; Dianat et al., 2015; Koyuncu & Karcioglu, 2018; Tantawy, 2019), and some find a weak significant relationship (Nagaraj et al., 2019). Similarly to BMI and years of working, most studies predict that both factors are significant factors for WRMSD (Dianat et al., 2015; Nagaraj et al., 2019) and some did not(Amin et al., 2014). The study of (Soe et al., 2015) also weight the finding of no association between income, years of work, and WRMSD, but working hours per day is associated with WRMSD in that study. In addition, the study of Nigeria sewing operators found no association of working hours per day, duration of work with WRMSD and support the finding of this study (Akinpelu et al., 2016). The primary reason that these personal factors which do not find an association with WRMSD is due to the finding the overall association of whole-body pain with statistical test. In other words, these variables are sometimes only associate with certain parts of body pain. These factors may be associated with WRMSD if we can find the individual part of the body analysis. Although, this study does not include the finding of individual body part association and therefore weakens the associated measures.

5.3 Association of Psychosocial risk factors and WRMSD

This study determines the psychosocial risk factors using the Job Content Questionnaire. It is based on the Karasek Demand control Support model. However, very few articles using this model to determine the psychosocial risks of sewing operators. Therefore, a comparison is done with studies of other industrial sectors. Although the job natures of sewing operation are the same in the garment factory, it is necessary to determine the high strain job according to the model. High strain job is determined by high psychological demand and low job control. It is more likely to be the perception of sewing operators and not depend on the job nature. In this study, 21% reported high strain jobs and 79% reported non-high strain jobs. This finding is consistent with the finding of occupational stress in Iranian car manufacturing workers about 21.3% (Soori, Rahimi, & Mohseni, 2008) and stress of dental workers in Kelantan 22.2% (Rusli, Edimansyah, & Naing, 2006). Moreover, the study of psychological stress in Korean workers using the job strain model is also 20% which is nearly similar to this study. (Rhee, 1999) . However, the prevalence of job stress in numerous studies is higher than in this study. The study of Malaysian automotive assembly plant with the prevalence of 31% (Edimansyah et al., 2008),

Thai rubber glove manufacturing company in which job strain prevalence reached 28% (Sein et al., 2010), Mohan and collaborators who measured a 25% rate of job strain among workers of a foundry company (Mohan, Elangovan, Prasad, Krishna, & Mokkapati, 2008) and Textile Company in Congo study which was showed that 28% of workers suffer from stress (Kitronza, Mairiaux, & health, 2015).

These psychosocial risk studies can be varied in different industrial sectors in different countries. There is very limited job stress study on sewing machine operators. The different methodology, different study areas, different working environments are the major contributors to obtain different job stress prevalence. Different perceives stress points are also can varied considering cultural differences, economic disparities between industrialized and emerging countries. In the European Union, 22% to 28% suffer at least one factor of psychosocial risk that may impact the well-being of mental health. (Eurofound, 2010)

The result in this study comprises seven items of psychosocial factors, namely high psychological demand(41.4%), low job control(44.6%), poor social support(23.0%), poor supervisor support(29.5%), poor coworker support(9.7%), high physical demand(75.9%), and high job insecurity(53.8%) respectively. To be summarized these factors, workplace social support is high, psychological, job control, job security are nearly equal in both high and low, and physical demand is the significant risk factor. These findings are in line with the result of Job Strain among Rubber-Glove-Factory Workers in Central Thailand except for job security item (Sein et al., 2010). Among the seven items, physical demand is the only factor related to WRMSD.

Very few articles are conducted between psychological risk factors and WRMSD using JCQ items. This finding is coherent with the finding of workorganizational and personal factors associated with upper-body musculoskeletal disorders among sewing machine operators in Los Angeles except for psychological demand. The high job demand in that study is associated with the prevalence of neck/shoulder pain(P.-C. Wang et al., 2007). The study of (Choobineh, Motamedzade, Kazemi, Moghimbeigi, & Heidari Pahlavian, 2011) also found that significant difference means before and after the intervention is physical demand, and none of the other JCQ items show any difference between them. This finding also supports the finding of job stress in Iranian nurses that all the items show no significant relationship to MSD but surprisingly found some association when they analyzed with individual parts of the back, upper extremities, lower extremity with WRMSD (Barzideh, Choobineh, & Tabatabaee, 2014). It can be noted that psychosocial factors can be associated with each of the body parts and this analysis does not include in this study. However, this result is not in line with the study of (Gholami, Rahnavard, Sadeghzadeh, & Tahmtan, 2018) that they found a significant association between social support, job security with WRMSD. Moreover, the study of IT professionals in India reported that psychological demand, decision latitude, job control, social support are significantly related to musculoskeletal pain (R. K. Mehta & Parijat, 2012). These contradictory findings can be due to different sample populations and different research frames.

The result in this study is high physical demand is 3.1 times more likely to develop WRMSD according to chi-square statistics. As regards physical demand, most of the studies found an association between physical demand and WRMSD. The study of (Gholami et al., 2018; Shan, Bin Adon, Rahman, Hassan, & Ismail, 2011; Wang et al., 2010) found that increases in physical exertion are more likely to get WRMSD than low physical demand. Job strain found no association with WRMSD which is not consistent with many studies (Barbieri, Nogueira, Bergamin, & Oliveira, 2012; P.-C. Wang et al., 2007). However, many researchers claimed that job strain is related to certain parts of the body such as the back, and not associated with whole-body pain which is proved by a meta-analysis study (Amiri & Behnezhad, 2020). Therefore, to get association, we need to analyze various parts of body pain and WRMSD.

5.4 Association of work ability and WRMSD

The mean work ability score is 34.3 and it can be regarded as a moderate ability score. There is no research article on the work ability of sewing machine operators using WAI scores. However, comparison can be done with other types of workers. This finding gets nearly the result of (Rostamabadi, Mazloumi, & Foroushan, 2014) with 35. Another study of work ability index in Slovenian hospital nurses also near with a mean score of 36 (Žmauc, Železnik, & Težak, 2019). However, most of the studies get higher WAI scores. The study of work ability of Iranian workers with 38.4 (Gharibi et al., 2016), computers workers in Portugal with 40.5 (A. F. Costa, Puga-Leal, & Nunes, 2011), and petrochemical workers in Iran about 39.1 (Mazloumi, Rostamabadi, Saraji, & Foroushani, 2012).In addition, the WAI score is also well validated for the need for rehabilitation if WAI scores less than 37 (Bethge, Radoschewski, & Gutenbrunner, 2012). The WAI score is also strongly associated with age and the younger age had higher scores than older age in women and should consider in this study (Monteiro, Ilmarinen, Filho, & Ergonomics, 2006).

Most of the percentages of WAI scores are occupied by poor and moderate ability with 84.9 %. It is very difficult to compare individual percentages because most of the articles did not show individual scores. Therefore comparative is done through the majority of the score. This result is consistent with the finding of work ability index (WAI) values in a sample of the working population in Poland with the majority of poor and moderate ability 65% (Juszczyk et al., 2019), and a test-retest study with 61% (Zwart, Frings-Dresen, & Duivenbooden, 2002). Moreover, another study of employees of the hospital's hygiene and cleaning sector in Brazil (90% women and 10% men) similar with age distribution with this study also reported that 68% (Monteiro et al., 2006). However, there are also contradictory findings that the excellent and good group mostly occupied the overall percentage of WAI score. The article of Comparison of the Work Ability Status in Manual and Office Workers, an Occupational Health Survey reported that excellent, good with 90% (Hosseininejad, Mirzamohammadi, Labbafinejad, Mazhari, & Mohammadi, 2017), and a similar report can be seen in work ability of Dutch construction workers with 85% (Alavinia et al., 2007). Another Hong Kong study of construction workers also reported a very high excellent and good score of 97%. It can be noted that work ability score is based on individual perception and it can be varied with different occupations and different working environments (Ng & Chan, 2018).

In this study, workability found no association with WRMSD. This finding is consistent with the finding of a large study among a variety of professionals in the UK, no significant association was found between self-rated work performances, ability with physical health, including musculoskeletal symptoms. It is reasonable to think that physical factors can be strong predictors of work ability. But one possible explanation is that there is a strong relationship between psychological health and physical health, then it covered the priority of impact physical health – in other words, musculoskeletal pain is indirectly impacting on work ability through mental well-being (Donald et al., 2005; SIU, Cooper, & Donald, 1997). However, many studies showed that there is a significant association between work ability and MSD (Fernandes, da Silva Pataro, De Carvalho, & Burdorf, 2016; Monteiro et al., 2006; Skovlund, Bláfoss, Sundstrup, & Andersen, 2020). But these findings are very specifically related to the certain body part and not stated the overall significant result. Therefore, further research is needed on industrial workers to determine the overall association. In addition, (Gharibi et al., 2016) study revealed that job stress is significantly associated with work ability and possible factors that should be considered when we analyze with WRMSD.

5.5 Association of work productivity and WRMSD

There is no significant reported absenteeism and overall work impairment is the same as the presenteeism score of 34%. Very few articles are conducted on garment workers and rarely use WPAI. This outcome is in line with the finding of workers' mental health and productivity of different employees in Japan with 34% (Kuroda, Yamamoto, & Economy, 2018). This result is more than the study of health problems that lead to considerable productivity loss at work among industrial workers with 21%. Moreover, that study also stated that work limitation due to WRMSD is 9% which comprised huge components of productivity impairment (Meerding et al., 2005). One of the Swedish studies demonstrated that during 12 month period, 37% of workers suffer productivity impairment (G. Aronsson et al., 2000). In addition, a large study of US employees also revealed that work impairment is 27% (Boles, Pelletier, & Lynch, 2004). Another study of American workers, a developed nation, also stated that average reported productivity losses when working while sick range around 20% (presenteeism). A significant difference may be due to huge differences in socioeconomic and working environment between countries.

Although there is no association between WRMSD and work productivity, logistic regression analysis showed that decreased productivity is the strong predictor of WRMSD. The possible explanation can be due to the chronic non-specific pain (CMP) nature of sewing operators. The work nature of sewing operation needs long working, static work and heavy pain cannot be suffered. They may be reported that musculoskeletal pain as CMP. Some studies stated that workers can perform well with CMP in work even though they have presenteeism. (Mannion et al., 2009; Van den Heuvel, Geuskens, Hooftman, Koppes, & Van den Bossche, 2010) Presenteeism "should not necessarily be interpreted as a negative thing, either for the individual or the company". Working with pain may be regarded as a healthy coping behavior, which will help to stabilize the workers' participation in work and quality of life(Waddell & Burton, 2006). Moreover, the lack of association can be explained by working hours. This might be explained by the term "extensionism", which has been introduced to describe the situation of working extended hours beyond those expected, to compensate for productivity loss by the employer. This was confirmed in our study, where actual working hours (>8hr) exceeded the expected hours (\leq 8hr). Reduce work productivity and performance can be replaced by extending more
working hours (negative absenteeism)(de Vries, Reneman, Groothoff, Geertzen, & Brouwer, 2013; Hilton, Sheridan, Cleary, & Whiteford, 2009). However many studies and literature are reported that WRMSD is a significant risk factor for work productivity (Boles et al., 2004; Daneshmandi, Choobineh, Ghaem, Alhamd, & Fakherpour, 2017; C. Jones et al., 2019)

Univariate analysis revealed that only three variables, age education, and physical demand are significantly associated with WRMSD. After controlling confounding variables, age, gender, education status, duration of work, physical demand, job insecurity, total social support, job stress, work productivity percentage (<50% and \geq 50%), work ability index scores are put into the regression analysis. The binary logistic regression model showed that gender, job stress, physical demand, and work productivity are significant predictors of WRMSD. Male is 87% less likely to develop than female with a p-value of 0.046, 95% CI (0.020-0.969). Concerning job stress, high strain job is 8.2 times likely to develop WRMSD than the low strain with a p-value of 0.017, 95% CI (1.456 - 46.550). Also, high physical demand is 4.7 times more likely to develop WRMSD than low demand with a p-value of 0.029, 95% CI (1.172 – 18.562). A similar situation is seen in work productivity percentage, the \geq 50 % productivity impairment is 5.8 times more likely to develop than the < 50% with a p-value of 0.016, 95% CI (1.393 – 24.920). Other variables are not significantly associated with WRMSD in regression analysis. 34% of the variables can be explained by all the combinations of independent variables of the regression model.



CHAPTER 6

CONCLUSION AND RECOMMENDATION

This cross-sectional study was conducted to determine work-related musculoskeletal disorders, psychosocial factors, work productivity, and work ability among garment factory workers in Myanmar in Shwepaukkan Industrial Zone, Yangon, Myanmar. The study was designed to be a cross-sectional descriptive study to access the prevalence of WRMSD and its association with personal factors, psychosocial factors, work productivity, and work ability among garment factory workers in Shwepaukkan Industrial Zone, Yangon, Myanmar. This study revealed the personal factors, psychosocial factors, work productivity, work ability, and WRMSD. Data were collected in 2 garment factories which are agreed to participate during the COVID situation from 10th August to 12th September 2020. 2 garment factories were selected out of 15 garment factories by convenience sampling method. The total required sample size was approximately about 400. One research assistant was recruited as a research assistant and rained for one day before data collection. Inclusion, Exclusion criteria were provided to the factory authorities then approached the respondents, introduced, and explained the nature and purpose of the study. Data collection was done through the self-administered method due to the COVID situation.

The study found out that high percentages of WRMSD with 93.8 in sewing machine operators. The prevalence rate is far higher than in other studies. Many

factors and variables are included in this study. This study found an association only with age, education, and physical demand. Moreover, the study found out that gender, job stress, physical demand, and work productivity are the significant predictor of suffering WRMSD in a high odd ratio although other factors are not significantly related.

6.1. Strengths and Limitations

6.1.1 Strengths of the study

- This is the first research that uses these four variables (psychosocial factors, work productivity, workability, and work-related musculoskeletal disorder) in Garment factory workers in Myanmar.
- This study show that the high prevalence of WRMSD, high physical demand high job stress, poor work ability are the common occupational hazards among sewing machine operators in Myanmar.
- This study can provide detailed data of all four main variables which lacks in most of the studies.
- The findings of the study will be useful to the Ministry of Health and Sports, Ministry of labor, other related Social welfare organizations, Ministry of Industry in promotion well-being of workers both physically, mentally, and socially.

- This finding can help in developing occupational health law which is still underdeveloped in Myanmar .e.g. law about working hour
- At the service provider level, public health authorities could be helped to advocate and guide in strengthening the well-being of workers in all aspects

6.1.2. Limitations of the study

Although this study contributes important findings on parenting practices in

Myanmar, there are many limitations.

- Firstly, as the data used in this study are collected from a cross-sectional study, the causality of the associations cannot be elaborated, so that, interpretation is limited as associations.
- Secondly, recall bias could affect the results because the survey relied exclusively on the self-administered type.
- Moreover, recall bias can affect all variables, under-reporting and overreporting of answers should be considered.
- Response bias should be considered because all are validated questionnaires and can lead to misleading and misunderstanding of questionnaires.
- Other workplace physical factors such as heat, noise, light, and environmental factors such as job design are not included in the study and this can be a huge influence on WRMSD prevalence.

 These findings may not represent male because 95.4% are occupied by female and focusing only on sewing machine operators and cannot represent the whole garment factory.

6.2. Recommendation

This study aimed to determine the percentages of work-related musculoskeletal disorder psychosocial factors, work productivity, workability of sewing machine operators. Based on the finding, recommendations are discussed in two components; for program implementation and further research.

6.2.1 Recommendation for program implementation

- The prevalence of WRMSD is very high compared to other studies. To reduce the prevalence of WRMSD, workers should be supported physically and mentally. For example high temperature, poor ventilation, limited working space, unpleasant odors, poor seating, and poor sanitation in toilets.
- Job stress prevalence is 20% which occupied one-fifth of the study population and it can be regarded as high job stress in sewing operators. To reduce this, self-centered stress management ways are recommended. As an example, playing music at the factory, exercise promotion during break time, building playground and creation of happy working environment by factory authorities.

- Working hours should be reduced as prolong working hours are confirmed in this study. Adjust the workload and reducing time pressure according to the action plan which will be designed by the factory authorities.
- Proper job training program to prevent WRMSD. Workers must learn about the risk factors of MSD's and how to make ergonomic changes to prevent them.
- As reported high physical demand, high WRMSD, it is better to provide ergonomic tools and chairs. Using the wrong tool can cause immense discomfort and long-term side effects.
- The work ability score is moderate and needs to be improved. Finding the factors that relate to the decrease work ability should be explored and implementation programs should start as early as possible as it can impact the country's economy.
- It is also important to strengthen the legal framework, policies, and services for the prevention and response to the well-being of workers. Hence, respective authorities should implement the occupational health law that is still underdeveloped in Myanmar.

6.2.2. Recommendation for further research

• As the study was conducted in Yangon Region, Myanmar, the percentages of WRMSD and association are different from other country regions. Therefore, it

is advisable to conduct in other regions in Myanmar to know more about the percentages of WRMSD and association.

- As the study includes only sewing operators, the other sectors of the garment factory workers should include in the further study.
- This study includes the self-administered type and many biases can contain.
 Therefore, other studies should be conducted with interviewed types to reduce biases.
- This study does not include physical factors such as job design, heat, noise light and should conduct in the future study using these factors.
- Both quantitative and qualitative measures should be conducted to identify the associated factors of WRMSD among sewing machine operators.
- Further prospective studies, evidence-based investigation, and research by using longitudinal study designs should be carried out based on the results and findings of this study to address the cause and effect relationship of WRMSD among sewing operators.
- Intervention programs to reduce WRMSD, reduce psychosocial factors, promote work ability and work productivity are also conducted to know the effectiveness of these programs which is valued for the well-being of the workers.

• Further study should use different instruments and different models for these factors to identify the WRMSD and associated factors.



Chulalongkorn University

REFERENCES



Chulalongkorn University

- Abaraogu, U., Okafor, C., Ezeukwu, A. O., & Igwe, S. E. (2015). Prevalence of workrelated musculoskeletal discomfort and its impact on activity: A Survey of Beverage Factory Workers in Eastern Nigeria. *Work (Reading, Mass.), 52.* doi:10.3233/WOR-152100
- Adel, M., Akbar, R., & Ehsan, G. (2019). Validity and reliability of work ability index
 (WAI) questionnaire among Iranian workers; a study in petrochemical and car manufacturing industries. *Journal of occupational health*, *61*(2), 165-174. doi:10.1002/1348-9585.12028
- Adib Ibrahim, M., Abdul Aziz, A., Suhaili, N. A., Zahid Daud, A., Naing, L., & Abdul Rahman, H. (2019). A Study into Psychosocial Work Stressors and Health Care Productivity. *Int J Occup Environ Med*, *10*(4), 185-193. doi:10.15171/ijoem.2019.1610
- Afifehzadeh-Kashani, H., Choobineh, A., Bakand, S., Gohari, M., Abbastabar, H., & Moshtaghi, P. (2011). Validity and reliability of farsi version of Cornell Musculoskeletal Discomfort Questionnaire (CMDQ).
- Afonso, L., Pinho, M., & Arezes, P. (2014). Prevalence of WMSD in the sewing sector of two companies of the footwear industry.
- Aghili, M. M., Asilian, H., & Poursafa, P. (2012). Evaluation of musculoskeletal disorders in sewing machine operators of a shoe manufacturing factory in Iran. *J Pak Med Assoc, 62*(3 Suppl 2), S20-25.
- Aghilinejad, M., Choobineh, A. R., Sadeghi, Z., Nouri, M. K., & Bahrami Ahmadi, A. (2012). Prevalence of Musculoskeletal Disorders among Iranian Steel Workers. *Iranian Red Crescent medical journal, 14*(4), 198-203.
- Ahmad, S. A., Sayed, M., Khan, M. H., Faruquee, M., Yasmin, N., Hossain, Z., . . . medicine, s. (2007). Musculoskeletal Disorders and Ergonomic Factors among the Garment Workers. *26*(2), 97-110.
- Airila, A., Hakanen, J., Punakallio, A., Lusa, S., & Luukkonen, R. (2012). Is work engagement related to work ability beyond working conditions and lifestyle factors? *Int Arch Occup Environ Health, 85*(8), 915-925. doi:10.1007/s00420-012-0732-1

- Ak, A., Ba, T., & Adebisi, O. A. (2013). Prevalence, Pattern and Impact of Musculoskeletal Disorders among Sewing Machine Operators in Surulere Local Government Area of Lagos State, Nigeria. *Indian Journal of Physiotherapy and Occupational Therapy*, 7(2), 15-20.
- Akinpelu, A., Oyewole, O., Odole, A., & Ogunbamowo, F. (2016). Work-related musculoskeletal pain and health-seeking behavior among Nigerian sewing machine operators. *Tropical Journal of Medical Research, 19*, 152.
- Akodu, A. K., & Ashalejo, Z. O. (2019). Work-related musculoskeletal disorders and work ability among hospital nurses. *Journal of Taibah University Medical Sciences, 14*(3), 252-261. doi:https://doi.org/10.1016/j.jtumed.2019.02.009
- Alavinia, S. M., van Duivenbooden, C., & Burdorf, A. (2007). Influence of work-related factors and individual characteristics on work ability among Dutch construction workers. *Scand J Work Environ Health, 33*(5), 351-357. doi:10.5271/sjweh.1151
- Albert, V., Coutu, M.-F., & Durand, M.-J. J. C. J. o. O. T. (2011). Structure factorielle et cohérence interne de l'Échelle de mesure du sentiment d'efficacité personnelle. *78*(5), 311-317.
- Allen, H. (2008). Using routinely collected data to augment the management of health and productivity loss. *J Occup Environ Med, 50*(6), 615-632. doi:10.1097/JOM.0b013e31817b610c
- Amin, N. A., Nordin, R., Fatt, Q. K., Noah, R. M., & Oxley, J. (2014). Relationship between Psychosocial Risk Factors and Work-Related Musculoskeletal Disorders among Public Hospital Nurses in Malaysia. *Annals of occupational* and environmental medicine, 26, 23-23. doi:10.1186/s40557-014-0023-2
- Amiri, S., & Behnezhad, S. (2020). Is job strain a risk factor for musculoskeletal pain? A systematic review and meta-analysis of 21 longitudinal studies. *Public Health,* 181, 158-167. doi:https://doi.org/10.1016/j.puhe.2019.11.023
- Andersen, J. H., & Gaardboe, O. (1993). Musculoskeletal disorders of the neck and upper limb among sewing machine operators: a clinical investigation. *Am J Ind Med, 24*(6), 689-700. doi:10.1002/ajim.4700240605

- Andersen, J. H., & Gaardboe, O. J. A. j. o. i. m. (1993). Prevalence of persistent neck and upper limb pain in a historical cohort of sewing machine operators. *24*(6), 677-687.
- Andersen, L. L., Izquierdo, M., Sundstrup, E. J. I. a. o. o., & health, e. (2017).
 - Overweight and obesity are progressively associated with lower work ability in the general working population: cross-sectional study among 10,000 adults. *90*(8), 779-787.
- Aronsson, G., Gustafsson, K., & Dallner, M. (2000). Sick but yet at work. An empirical study of sickness presenteeism. *Journal of epidemiology and community health, 54*(7), 502-509. doi:10.1136/jech.54.7.502
- Aronsson, G., Gustafsson, K. J. J. o. o., & medicine, e. (2005). Sickness presenteeism: prevalence, attendance-pressure factors, and an outline of a model for research. *47*(9), 958-966.
- ASSOCIATES, R. (2019). WPAI General Information.
- Bakker, A. B., & Demerouti, E. J. J. o. m. p. (2007). The job demands-resources model: State of the art.
- Barbieri, D. F., Nogueira, H. C., Bergamin, L. J., & Oliveira, A. B. (2012). Physical and psychosocial indicators among office workers from public sector with and without musculoskeletal symptoms. *Work, 41*, 2461-2466. doi:10.3233/WOR-2012-0481-2461
- Barr, A. E., Amin, M., & Barbe, M. F. (2002). Dose-Response Relationship between Reach Repetition and Indicators of Inflammation and Movement Dysfunction in a Rat Model of Work-Related Musculoskeletal Disorder. *Proceedings of the Human Factors and Ergonomics Society Annual Meeting, 46*(16), 1486-1490. doi:10.1177/154193120204601617
- Barr, A. E., Barbe, M. F., & Clark, B. D. (2004). Work-related musculoskeletal disorders of the hand and wrist: epidemiology, pathophysiology, and sensorimotor changes. J Orthop Sports Phys Ther, 34(10), 610-627. doi:10.2519/jospt.2004.34.10.610

- Barzideh, M., Choobineh, A., & Tabatabaee, H. J. W. (2014). Job stress dimensions and their relationship to musculoskeletal disorders in Iranian nurses. *47*(4), 423-429.
- Berecki-Gisolf, J., Clay, F. J., Collie, A., & McClure, R. J. J. J. o. o. r. (2012). Predictors of sustained return to work after work-related injury or disease: insights from workers' compensation claims records. *22*(3), 283-291.
- Bernard, B. P., & Putz-Anderson, V. (1997). Musculoskeletal disorders and workplace factors; a critical review of epidemiologic evidence for work-related musculoskeletal disorders of the neck, upper extremity, and low back.
- Bethge, M., Radoschewski, F. M., & Gutenbrunner, C. J. J. o. r. m. (2012). The Work Ability Index as a screening tool to identify the need for rehabilitation: longitudinal findings from the Second German Sociomedical Panel of Employees. 44(11), 980-987.
- Bevan, S. (2015). Economic impact of musculoskeletal disorders (MSDs) on work in Europe. *Best Practice & Research Clinical Rheumatology, 29*(3), 356-373. doi:https://doi.org/10.1016/j.berh.2015.08.002
- Birschel, D. J. B. Q. (2006). Lost-Worktime Injuries and Illnesses: Characteristics and Resulting Time Away From Work, 2004. *22*(2), 61.
- Blåder, S., Barck-Holst, U., Danielsson, S., Ferhm, E., Kalpamaa, M., Leijon, M., . . . Markhede, G. (1991). Neck and shoulder complaints among sewing-machine operators: a study concerning frequency, symptomatology and dysfunction. *Appl Ergon, 22*(4), 251-257. doi:10.1016/0003-6870(91)90228-a
- Boles, M., Pelletier, B., & Lynch, W. (2004). The Relationship Between Health Risks and Work Productivity. *Journal of Occupational and Environmental Medicine*, *46*(7).
- Burnley, D. G. a. J. (December 2015). MADE IN MYANMAR -Entrenched poverty or decent jobs for garment workers? *Oxfam International December 2015*.
- Burton, W. N., Conti, D. J., Chen, C. Y., Schultz, A. B., & Edington, D. W. (1999). The role of health risk factors and disease on worker productivity. J Occup Environ Med, 41(10), 863-877. doi:10.1097/00043764-199910000-00007

- Burton, W. N., Conti Dj Fau Chen, C. Y., Chen Cy Fau Schultz, A. B., Schultz Ab Fau
 Edington, D. W., & Edington, D. W. The role of health risk factors and disease on worker productivity. (1076-2752 (Print)).
- Bush, I. L. N. a. P. M. (2012). Work-Related Musculoskeletal Disorders Assessment and Prevention, Ergonomics - A Systems Approach, Dr. Isabel L. Nunes (Ed.),. doi:978-953-51-0601-2
- Cancelliere, C., Cassidy, J. D., Ammendolia, C., & Côté, P. (2011). Are workplace health promotion programs effective at improving presenteeism in workers? A systematic review and best evidence synthesis of the literature. *BMC Public Health, 11*, 395-395. doi:10.1186/1471-2458-11-395
- Carroll, L., Mercado, A. C., Cassidy, J. D., & CĴté, P. (2002). A population-based study of factors associated with combinations of active and passive coping with neck and low back pain. *J Rehabil Med, 34*(2), 67-72. doi:10.1080/165019702753557854
- Caruso, C. C., Bushnell, T., Eggerth, D., Heitmann, A., Kojola, B., Newman, K., . . . Vila,B. J. A. j. o. i. m. (2006). Long working hours, safety, and health: toward a National Research Agenda. *49*(11), 930-942.
- Casey, L. P.-T. L., R. (2014). Stress and Wellbeing Survey 2014. Australia: APS. Australian Psychological Society (APS).
- Chang, Y.-F., Yeh, C.-M., Huang, S.-L., Ho, C.-C., Li, R.-H., Wang, W.-H., & Tang, F.-C. (2020). Work Ability and Quality of Life in Patients with Work-Related Musculoskeletal Disorders. *17*(9), 3310.
- Charan, J., & Biswas, T. (2013). How to calculate sample size for different study designs in medical research? *Indian journal of psychological medicine, 35*(2), 121-126. doi:10.4103/0253-7176.116232
- Choobineh, A., Motamedzade, M., Kazemi, M., Moghimbeigi, A., & Heidari Pahlavian, A. (2011). The impact of ergonomics intervention on psychosocial factors and musculoskeletal symptoms among office workers. *International Journal of Industrial Ergonomics, 41*(6), 671-676.

doi:https://doi.org/10.1016/j.ergon.2011.08.007

- Cordeiro, T. M. S. C. e., Souza, D. S., & Araújo, T. M. d. (2017). Validity, Reproducibility and Reliability of the Work Ability Index: A Systematic Review. *2017, 7*(1), 10 %J Revista de Epidemiologia e Controle de Infecção. doi:10.17058/reci.v7i1.7788
- Costa, A. F., Puga-Leal, R., & Nunes, I. L. (2011). An exploratory study of the Work Ability Index (WAI) and its components in a group of computer workers. *Work, 39*, 357-367. doi:10.3233/WOR-2011-1186
- Costa, B., & Vieira, E. (2009). Risk Factors for Work-Related Musculoskeletal Disorders: A Systematic Review of Recent Longitudinal Studies. *American Journal of Industrial Medicine, 53*, 285-323. doi:10.1002/ajim.20750
- Da Costa, B. R., & Vieira, E. R. J. A. j. o. i. m. (2010). Risk factors for work-related musculoskeletal disorders: a systematic review of recent longitudinal studies. *53*(3), 285-323.
- Daneshmandi, H., Choobineh, A. R., Ghaem, H., Alhamd, M., & Fakherpour, A. (2017). The effect of musculoskeletal problems on fatigue and productivity of office personnel: a cross-sectional study. *Journal of preventive medicine and hygiene, 58*(3), E252-E258.
- Daniel, W. W., & Cross, C. L. (2018). *Biostatistics: a foundation for analysis in the health sciences*: Wiley.
- de Vries, H. J., Reneman, M. F., Groothoff, J. W., Geertzen, J. H. B., & Brouwer, S. (2013). Self-reported Work Ability and Work Performance in Workers with Chronic Nonspecific Musculoskeletal Pain. *J Occup Rehabil, 23*(1), 1-10. doi:10.1007/s10926-012-9373-1
- De Witte, H. J. S. j. o. I. P. (2005). Job insecurity: Review of the international literature on definitions, prevalence, antecedents and consequences. *31*(4), 1-6.
- de Zwart, B. C. H., Frings-Dresen, M. H. W., & van Duivenbooden, J. C. (2002). Testretest reliability of the Work Ability Index questionnaire. *Occupational Medicine, 52*(4), 177-181. doi:10.1093/occmed/52.4.177 %J Occupational Medicine

- Dianat, I., Kord, M., Yahyazade, P., Karimi, M. A., & Stedmon, A. W. (2015). Association of individual and work-related risk factors with musculoskeletal symptoms among Iranian sewing machine operators. *Applied Ergonomics, 51*, 180-188. doi:https://doi.org/10.1016/j.apergo.2015.04.017
- Dianat, I., & Salimi, A. J. E. (2014). Working conditions of Iranian hand-sewn shoe workers and associations with musculoskeletal symptoms. *57*(4), 602-611.
- Donald, I., Taylor, P., Johnson, S., Cooper, C., Cartwright, S., & Robertson, S. J. I. J. o. S. M. (2005). Work environments, stress, and productivity: An examination using ASSET. *12*(4), 409.
- Edimansyah, B. A., Rusli, B. N., Naing, L., Rusli, B. A. M., Winn, T., & Ariff, B. R. H. T. M. J. I. h. (2008). Self-perceived depression, anxiety, stress and their relationships with psychosocial job factors in male automotive assembly workers. *46*(1), 90-100.
- El Fassi, M., Bocquet, V., Majery, N., Lair, M. L., Couffignal, S., & Mairiaux, P. (2013). Work ability assessment in a worker population: comparison and determinants of Work Ability Index and Work Ability score. *BMC Public Health, 13*(1), 305. doi:10.1186/1471-2458-13-305
- Engkvist, I.-L., Hjelm, E. W., Hagberg, M., Menckel, E., & Ekenvall, L. (2000). Risk indicators for reported over-exertion back injuries among female nursing personnel. In: LWW.
- Erdinc, O., Hot, K., & Ozkaya, M. J. W. (2011). Turkish version of the Cornell Musculoskeletal Discomfort Questionnaire: cross-cultural adaptation and validation. *39*(3), 251-260.
- Escorpizo, R. (2008). Understanding work productivity and its application to workrelated musculoskeletal disorders. *International Journal of Industrial Ergonomics, 38*(3), 291-297. doi:https://doi.org/10.1016/j.ergon.2007.10.018
- Escorpizo, R., Bombardier, C., Boonen, A., Hazes, J. M., Lacaille, D., Strand, V., & Beaton, D. J. T. J. o. R. (2007). Worker productivity outcome measures in arthritis. *34*(6), 1372-1380.

Eurofound. (2007). 4th European Working Survey.

Eurofound. (2010). Fifth European Working Conditions survey 2010.

Eurofound. (2017). Sixth European Working Conditions Survey – Overview report (2017 update). *Publications Office of the European Union, Luxembourg.*

- Fernandes, R. d. C. P., da Silva Pataro, S. M., De Carvalho, R. B., & Burdorf, A. J. B. P. H. (2016). The concurrence of musculoskeletal pain and associated work-related factors: a cross sectional study. *16*(1), 1-9.
- Ferreira Jr, M., & Saldiva, P. H. J. A. E. (2002). Computer–telephone interactive tasks: predictors of musculoskeletal disorders according to work analysis and workers' perception. *33*(2), 147-153.
- Fogleman, M., & Lewis, R. J. J. I. J. o. I. E. (2002). Factors associated with self-reported musculoskeletal discomfort in video display terminal (VDT) users. *29*(6), 311-318.
- Forastieri, V. (2016). Psychosocial risks, stress and violence in the world of work. International Journal of Labour Research, 8(1-2).
- Gharibi, V., Mokarami, H., Taban, A., Yazdani Aval, M., Samimi, K., & Salesi, M. (2016).
 Effects of Work-Related Stress on Work Ability Index among Iranian Workers.
 Safety and health at work, 7(1), 43-48.
 doi:https://doi.org/10.1016/j.shaw.2015.10.005
- Ghasemkhani, M., Mahmudi, E., & Jabbari, H. (2008). Musculoskeletal Symptoms In Workers. *International Journal of Occupational Safety and Ergonomics, 14*(4), 455-462. doi:10.1080/10803548.2008.11076784
- Gholami, T., Rahnavard, N., Sadeghzadeh, M., & Tahmtan, S. (2018). Application of MAPO Index and JCQ in Assessing the Risk of Musculoskeletal Disorders and Psychosocial Stressors in Nurse Aids at Fasa Hospital. *Iran-J-Ergon, 6*(2), 24-33. doi:10.30699/jergon.6.2.24
- Godde, K., & Taylor, R. W. J. F. s. i. (2011). Musculoskeletal stress marker (MSM) differences in the modern American upper limb and pectoral girdle in relation to activity level and body mass index (BMI). *210*(1-3), 237-242.
- Gold, J. E., Punnett, L., & Katz, J. N. (2006). Pressure pain thresholds and musculoskeletal morbidity in automobile manufacturing workers.
 International Archives of Occupational and Environmental Health, 79(2), 128-134. doi:10.1007/s00420-005-0005-3

- Gupta, R. D. (2015). Occupational Health Hazards among Workers in Garment Factories in Bangladesh: A Cross-Sectional Study *IISTE, Vol.5, No.5*.
- Hagberg, M., Tornqvist, E. W., & Toomingas, A. J. J. o. o. r. (2002). Self-reported reduced productivity due to musculoskeletal symptoms: associations with workplace and individual factors among white-collar computer users. *12*(3), 151-162.
- Hall, G. B., Dollard, M. F., & Coward, J. J. I. J. o. S. M. (2010). Psychosocial safety climate: Development of the PSC-12. *17*(4), 353.
- Harris, C., Daniels, K., Briner, R. B. J. B. J. o. G., & Counselling. (2004). How do work stress and coping work? Toward a fundamental theoretical reappraisal. *32*(2), 223-234.
- Hauke, A., Flintrop, J., Brun, E., & Rugulies, R. (2011). The impact of work-related psychosocial stressors on the onset of musculoskeletal disorders in specific body regions: A review and meta-analysis of 54 longitudinal studies. *Work & Stress, 25*(3), 243-256. doi:10.1080/02678373.2011.614069
- Health, C. o. S. a. (2015). Why Education Matters to Health: Exploring the Causes.
- Hedge, A., Morimoto, S., & Mccrobie, D. J. E. (1999). Effects of keyboard tray geometry on upper body posture and comfort. *42*(10), 1333-1349.
- Hilton, M. F., Sheridan, J., Cleary, C. M., & Whiteford, H. A. J. I. J. o. M. i. P. R. (2009). Employee absenteeism measures reflecting current work practices may be instrumental in a re-evaluation of the relationship between psychological distress/mental health and absenteeism. *18*(1), 37-47.
- Hossain, M. D., Aftab, A., Al Imam, M. H., Mahmud, I., Chowdhury, I. A., Kabir, R. I., & Sarker, M. J. P. o. (2018). Prevalence of work related musculoskeletal disorders (WMSDs) and ergonomic risk assessment among readymade garment workers of Bangladesh: A cross sectional study. *13*(7), e0200122.
- Hosseininejad, M., Mirzamohammadi, E., Labbafinejad, Y., Mazhari, A. M., & Mohammadi, S. J. J. P. H. (2017). Comparison of the Work Ability Status in Manual and Office Workers–an Occupational Health Survey. *2*(3), 555590.
- Hoy, D. G., Smith, E., Cross, M., Sanchez-Riera, L., Blyth, F. M., Buchbinder, R., . . . March, L. M. (2015). Reflecting on the global burden of musculoskeletal

conditions: lessons learnt from the global burden of disease 2010 study and the next steps forward. *Ann Rheum Dis, 74*(1), 4-7. doi:10.1136/annrheumdis-2014-205393

Hoy, D. G., Smith, E., Cross, M., Sanchez-Riera, L., Buchbinder, R., Blyth, F. M., . . .
March, L. M. (2014). The global burden of musculoskeletal conditions for 2010: an overview of methods. *Ann Rheum Dis, 73*(6), 982-989. doi:10.1136/annrheumdis-2013-204344

HSE. (2020). Working days lost in Great Britain.

- Ilmarinen, J. J. O. m. (2007). The work ability index (WAI). 57(2), 160-160.
- Ilmarinen, J. J. S. j. o. w., environment, & health. (2009). Work ability—a comprehensive concept for occupational health research and prevention. *35*(1), 1-5.
- ILO. (2016). Workplace stress: a collective challange.
- ILO. (2021a). Decent work for sustainable development (DW4SD) Resource Platform
- ILO. (2021b). Psychosocial risks and work-related stress.
- Jaffar, N. A. T., & Rahman, M. N. A. (2017). Review on risk factors related to lower back disorders at workplace. *IOP Conference Series: Materials Science and Engineering, 226*, 012035. doi:10.1088/1757-899x/226/1/012035
- Jahan, N., Das, M., Mondal, R., Paul, S., Saha, T., Akhtar, R., . . . Banik, P. C. J. S. m. j. (2015). Prevalence of musculoskeletal disorders among the Bangladeshi garments workers. *2*(1), 102-113.
- Johnson, J. V., Hall, E. M., Theorell, T. J. S. j. o. w., environment, & health. (1989). Combined effects of job strain and social isolation on cardiovascular disease morbidity and mortality in a random sample of the Swedish male working population. 271-279.
- Johnson, J. V., & Hall, E. M. J. A. j. o. p. h. (1988). Job strain, work place social support, and cardiovascular disease: a cross-sectional study of a random sample of the Swedish working population. *78*(10), 1336-1342.
- Johnston, V., Souvlis, T., Jimmieson, N. L., & Jull, G. J. A. e. (2008). Associations between individual and workplace risk factors for self-reported neck pain and disability among female office workers. *39*(2), 171-182.

- Jones, C., Payne, K., Gannon, B., & Verstappen, S. (2016). Economic Theory and Self-Reported Measures of Presenteeism in Musculoskeletal Disease. *Current rheumatology reports, 18*(8), 53-53. doi:10.1007/s11926-016-0600-1
- Jones, C., Verstappen, S. M. M., & Payne, K. (2019). A Systematic Review of Productivity in Economic Evaluations of Workplace Interventions: A Need for Reporting Criteria? *Applied Health Economics and Health Policy, 17*(5), 591-613. doi:10.1007/s40258-019-00473-8
- Jones, F., Bright, J., & Clow, A. (2001). *Stress : myth, theory and research*. Harlow, England; New York: Prentice Hall.
- Juszczyk, G., Czerw, A. I., Religioni, U., Olejniczak, D., Walusiak-Skorupa, J., Banas, T., . . . Staniszewska, A. (2019). Work Ability Index (WAI) values in a sample of the working population in Poland. *26*(1), 78-84. doi:10.26444/aaem/91471
- Kabir, H., Maple, M., Usher, K., & Islam, M. S. (2019). Health vulnerabilities of readymade garment (RMG) workers: a systematic review. *BMC Public Health*, *19*(1), 70. doi:10.1186/s12889-019-6388-y
- Kaka, B., Idowu, O. A., Fawole, H. O., Adeniyi, A. F., Ogwumike, O. O., & Toryila, M. T. (2016). An Analysis of Work-Related Musculoskeletal Disorders Among Butchers in Kano Metropolis, Nigeria. *Safety and health at work, 7*(3), 218-224. doi:https://doi.org/10.1016/j.shaw.2016.01.001
- Karasek Jr, R. A. J. A. s. q. (1979). Job demands, job decision latitude, and mental strain: Implications for job redesign. 285-308.
- Karasek, R., Brisson, C., Kawakami, N., Houtman, I., Bongers, P., & Amick, B. (1998). The Job Content Questionnaire (JCQ): an instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol, 3*(4), 322-355. doi:10.1037//1076-8998.3.4.322
- Karlqvist, L., Tornqvist, E. W., Hagberg, M., Hagman, M., & Toomingas, A. (2002). Selfreported working conditions of VDU operators and associations with musculoskeletal symptoms: a cross-sectional study focussing on gender differences. *International Journal of Industrial Ergonomics, 30*(4), 277-294. doi:https://doi.org/10.1016/S0169-8141(02)00131-2

- Kaufman, J. D., & Rosenstock, L. (1991). The role of worker education in preventing occupational lung disease. *Occup Med, 6*(1), 1-9.
- Kessler, R. C., Ames, M., Hymel, P. A., Loeppke, R., McKenas, D. K., Richling, D. E., . . . Medicine, E. (2004). Using the World Health Organization Health and Work Performance Questionnaire (HPQ) to evaluate the indirect workplace costs of illness. *46*(6), S23-S37.
- Kim, Y., Park, J., Rhee, K. Y., & Kim, H. (2015). Second Korean Working Conditions Survey: A Comparison between the Second KWCS and the First Korean Working Conditions Survey. *Safety and health at work, 1.* doi:10.1016/j.shaw.2015.02.005
- Kitronza, P. L., Mairiaux, P. J. T. m., & health. (2015). Occupational stress among textile workers in the Democratic Republic of Congo.
- Kivimäki, M., Vahtera, J., Ferrie, J. E., Hemingway, H., Pentti, J. J. O., & medicine, e. (2001). Organisational downsizing and musculoskeletal problems in employees: a prospective study. *58*(12), 811-817.
- Koyuncu, N., & Karcioglu, Ö. (2018). Musculoskeletal complaints in healthcare personnel in hospital: An interdepartmental, cross-sectional comparison. *Medicine*, *97*(40).
- Krause, N., Rugulies, R., Ragland, D. R., & Syme, S. L. J. A. j. o. i. m. (2004). Physical workload, ergonomic problems, and incidence of low back injury: A 7.5-year prospective study of San Francisco transit operators. *46*(6), 570-585.
- Kreuzfeld, S., Seibt, R., Kumar, M., Rieger, A., Stoll, R. J. J. O. O. M., & Toxicology.(2016). German version of the Cornell Musculoskeletal DiscomfortQuestionnaire (CMDQ): translation and validation. *11*(1), 1-12.
- Kuorinka, I., Forcier, L., Hagberg, M., Silverstein, B., Wells, R., Smith, M., . . . Pérusse, M. (1995). Work related musculoskeletal disorders (WMSDs): a reference book for prevention: Taylor & Francis London.
- Kuroda, S., Yamamoto, I. J. J., & Economy, t. W. (2018). Good boss, bad boss, workers' mental health and productivity: Evidence from Japan. *48*, 106-118.
- Lang, J., Ochsmann, E., Kraus, T., Lang, J. W. J. S. s., & medicine. (2012). Psychosocial work stressors as antecedents of musculoskeletal problems: a systematic

review and meta-analysis of stability-adjusted longitudinal studies. *75*(7), 1163-1174.

- Lazarus, R. S., & Launier, R. (1978). Stress-related transactions between person and environment. In *Perspectives in interactional psychology* (pp. 287-327): Springer.
- Lazarus, R. S. J. J. o. p. (2006). Emotions and interpersonal relationships: Toward a person-centered conceptualization of emotions and coping. *74*(1), 9-46.
- Lazarus, R. S. J. S., appraisal, & coping. (1986). Folkman. S.(1984). 117-180.
- Li, G., Haslegrave, C. M., & Corlett, E. N. J. A. e. (1995). Factors affecting posture for machine sewing tasks: the need for changes in sewing machine design. *26*(1), 35-46.
- Lidstone, J., Ells, L., Finn, P., Whittaker, V., Wilkinson, J., & Summerbell, C. J. P. h. (2006). Independent associations between weight status and disability in adults: results from the Health Survey for England. *120*(5), 412-417.
- Link, B. G., Phelan, J. J. J. o. h., & behavior, s. (1995). Social conditions as fundamental causes of disease. 80-94.
- Lowe, G. S. (2003). *Healthy workplaces and productivity: a discussion paper*: Minister of Public Works and Government Services Canada.
- Maduagwu, S., Sokunbi, G., Bwala, M., Akanbi, O., Jajere, A., Jaiyeola, O., . . . Affairs, H. (2015). Work-related musculoskeletal disorders among self employed sewing machine operators in maiduguri, Nigeria.
- Maduagwu, S. M., Maijindadi, R. D., Duniya, K. I., Oyeyemi, A. A., Saidu, I. A., Aremu, B. J. J. O. M., & Affairs, H. (2014). Prevalence and patterns of work-related musculoskeletal disorders among bankers in Maiduguri, Northeast Nigeria. 1-6.
- Mannion, A. F., Horisberger, B., Eisenring, C., Tamcan, O., Elfering, A., Müller, U. J. J. o. O., & Medicine, E. (2009). The association between beliefs about low back pain and work presenteeism. *51*(11), 1256-1266.
- Marian Boquiren, M. I. V. I., Thein Than Htay and Aung Myaing Htay. (2019). OccupationalSafety and Health in the MyanmarGarment Sector;EXECUTIVE SUMMARY. *ILO*.

- Mazloumi, A., Rostamabadi, A., Saraji, J. N., & Foroushani, A. R. J. J. o. o. h. (2012). Work ability index (WAI) and its association with psychosocial factors in one of the petrochemical industries in Iran. 1202130234-1202130234.
- McTernan, W. P., Dollard, M. F., LaMontagne, A. D. J. W., & Stress. (2013). Depression in the workplace: An economic cost analysis of depression-related productivity loss attributable to job strain and bullying. *27*(4), 321-338.
- Meerding, W. J., Ijzelenberg, W., Koopmanschap, M. A., Severens, J. L., & Burdorf, A.
 (2005). Health problems lead to considerable productivity loss at work among workers with high physical load jobs. *Journal of Clinical Epidemiology, 58*(5), 517-523. doi:https://doi.org/10.1016/j.jclinepi.2004.06.016
- Mehta, E., Mehta, M., Sharma, P. K. J. I. J. o. H., & Wellbeing. (2020). A study on workrelated musculoskeletal disorders among sewing machine operators. *11*.
- Mehta, R. K., & Parijat, P. J. W. (2012). Associations between psychosocial risk factors and musculoskeletal disorders: application to the IT profession in India. *41*(Supplement 1), 2438-2444.
- Merisalu, E., Mannaste, M., Hiir, K., & Traumann, A. J. A. R. (2016). Predictors and prevalence of musculoskeletal disorders among sewing machine operators. *14*(4), 1417-1426.
- MHLW. (2011). Survey on the Prevention of Industrial Accidents(Tokyo, MHLW).
- Ming, Z., & Zaproudina, N. J. P. (2003). Computer use related upper limb musculoskeletal (ComRULM) disorders. *9*(3), 155-160.
- Mohan, G. M., Elangovan, S., Prasad, P., Krishna, P. R., & Mokkapati, A. K. J. W. (2008). Prevalence of job strain among Indian foundry shop floor workers. *30*(4), 353-357.
- Monteiro, M. S., Ilmarinen, J., Filho, H. R. C. J. I. J. o. O. S., & Ergonomics. (2006). Work ability of workers in different age groups in a public health institution in Brazil. *12*(4), 417-427.
- Nag, A., Vyas, H., Shah, P., & Nag, P. K. (2012). Risk factors and musculoskeletal disorders among women workers performing fish processing. *55*(9), 833-843. doi:https://doi.org/10.1002/ajim.22075

- Nagaraj, T. S., & Jeyapaul, R. (2018). Ergonomic Study on Work Postures of Sewing Machine Operators in Government Industry: A Case in Lean Environment Garment Industry. In *Ergonomic Design of Products and Worksystems-21st Century Perspectives of Asia* (pp. 83-101): Springer.
- Nagaraj, T. S., Jeyapaul, R., & Mathiyazhagan, K. J. I. J. o. I. E. (2019). Evaluation of ergonomic working conditions among standing sewing machine operators in Sri Lanka. *70*, 70-83.
- Nazari, J., Mahmoudi, N., Dianat, I., & Graveling, R. J. H. p. p. (2012). Working Conditions in Carpet Weaving Workshops and Muscu-loskeletal Complaints among Workers in Tabriz-Iran. *2*(2), 265.
- Ng, J. Y. K., & Chan, A. H. S. (2018). The Work Ability of Hong Kong Construction Workers in Relation to Individual and Work-Related Factors. *Int J Environ Res Public Health, 15*(5). doi:10.3390/ijerph15050990
- Niu, S. (2010). Ergonomics and occupational safety and health: An ILO perspective. *Applied Ergonomics, 41*(6), 744-753. doi:https://doi.org/10.1016/j.apergo.2010.03.004
- Nourbakhsh, M. R., Moussavi, S. J., & Salavati, M. J. C. S. S. (2001). Effects of lifestyle and work-related physical activity on the degree of lumbar lordosis and chronic low back pain in a Middle East population. *14*(4), 283-292.
- Nunes, I. L. J. W. (2009). FAST ERGO_X–a tool for ergonomic auditing and work-related musculoskeletal disorders prevention. *34*(2), 133-148.
- Nygård, C.-H., Savinainen, M., Kirsi, T., & Lumme-Sandt, K. (2011). *Age Management during the Life Course-Proceedings of the 4th Symposium on Work Ability*: Tampere University Press.
- O'Donnell, M. P. (2000). Health and Productivity Management: The Concept, Impact, and Opportunity: Commentary to Goetzel and Ozminkowski. *American Journal of Health Promotion, 14*(4), 215-217. doi:10.4278/0890-1171-14.4.215
- Okunribido, O. J. H., & Excutive, S. (2009). Lower Limb MSD-Scoping work to help inform advice and research planning.

- Oranye, N. O., Wallis, B., Roer, K., Archer-Heese, G., Aguilar, Z. J. W. h., & safety. (2016). Do personal factors or types of physical tasks predict workplace injury? *, 64*(4), 141-151.
- Oyebamiji, M., & Akintayo, D. J. W. J. o. E. (2011). Some Psychological Factors as Predictors of Perceived Workers' Productivity in Private Organizations in Nigeria. 1(2), 30-38.
- Öztürk, N., & Esin, M. N. (2011). Investigation of musculoskeletal symptoms and ergonomic risk factors among female sewing machine operators in Turkey. *International Journal of Industrial Ergonomics, 41*(6), 585-591. doi:https://doi.org/10.1016/j.ergon.2011.07.001
- Pacheco Ferreira, A. J. C. T. (2015). Work ability and psychosocial factors among hairdressers workers, Rio de Janeiro, Brazil. 83-88.
- Pensola, T., Haukka, E., Kaila-Kangas, L., Neupane, S., & Leino-Arjas, P. J. S. j. o. p. h. (2016). Good work ability despite multisite musculoskeletal pain? A study among occupationally active Finns. 44(3), 300-310.
- Phillips, C., Main, C., Buck, R., Aylward, M., Wynne-Jones, G., & Farr, A. (2008).
 Prioritising pain in policy making: The need for a whole systems perspective. *Health Policy, 88*(2), 166-175.
 doi:https://doi.org/10.1016/j.healthpol.2008.03.008
- Phillips, V., Hunsaker, A., & Florence, C. S. J. S. c. (2012). Return to work and productive activities following a spinal cord injury: the role of income and insurance. *50*(8), 623-626.
- Polajnar, A., Leber, M., & Herzog, N. V. J. S. V. J. o. M. E. (2010). Muscular-Skeletal Diseases Require Scientifically Designed Sewing Workstations. *56*(1).
- Pourhoseingholi, M. A., Vahedi, M., & Rahimzadeh, M. (2013). Sample size calculation in medical studies. *Gastroenterology and hepatology from bed to bench, 6*(1), 14-17.
- Punnett, L., Herbert, R.,. (2000). Work-related musculoskeletal disorders: is there a gender differential, and so, what does it mean? In: Goldman, M., Hatch, M. (Eds.), Women & Health. Academic Press, USA, pp. 474–492.

- Radkiewicz, P., & Widerszal-Bazyl, M. (2005). Psychometric properties of Work Ability Index in the light of comparative survey study. *International Congress Series, 1280*, 304-309. doi:https://doi.org/10.1016/j.ics.2005.02.089
- Reddy, E. M., & Yasobant, S. (2015). Musculoskeletal disorders among municipal solid waste workers in India: A cross-sectional risk assessment. *Journal of family medicine and primary care, 4*(4), 519-524. doi:10.4103/2249-4863.174270
- Reilly, M. C., Zbrozek, A. S., & Dukes, E. M. (1993). The Validity and Reproducibility of a Work Productivity and Activity Impairment Instrument. *PharmacoEconomics*, 4(5), 353-365. doi:10.2165/00019053-199304050-00006
- Rhee, K. J. A.-P. N. (1999). Psychological stress in the Korean work life: a study based on the job strain model. 1-2.
- Ring, D., Kadzielski, J., Malhotra, L., Lee, S.-G. P., Jupiter, J. B. (2005). Psychological Factors Asociated with Idiopathic Arm Pain. *The Journal of Bone and Joint Surgery(American), Vol. 87, No. 2, pp. 374-380*
- Rocha, L. E., & Debert-Ribeiro, M. J. R. d. s. p. (2001). Work and health: a gender study on systems analysts. *35*(6), 539-547.
- Rohilla, E., Hooda, R., & Mehta, M. (2020). Prevalence of Musculoskeletal Disorders Among Garment Industry.
- Roquelaure, Y., LeManach, A. P., Ha, C., Poisnel, C., Bodin, J., Descatha, A., & Imbernon, E. J. O. m. (2012). Working in temporary employment and exposure to musculoskeletal constraints. *62*(7), 514-518.
- Rostamabadi, A., Mazloumi, A., & Foroushan, A. R. J. J. o. o. h. (2014). Work Ability Index (WAI) and its health-related determinants among Iranian farmers working in small farm enterprises. 13-0171-OA.
- Rusli, B., Edimansyah, B., & Naing, L. J. A. o. O. S. (2006). Prevalence and associated factors of stress in dental healthcare workers of a higher institution of learning in Kelantan. *1*, 51-56.
- Sadeghi Yarandi, M., Ghasemi, M., & Ghanjal, A. (2020). The relationship between individual, physical and psychosocial risk factors with musculoskeletal disorders and related disabilities in flight security personnel. *International*

Journal of Occupational Safety and Ergonomics, 1-11. doi:10.1080/10803548.2020.1760528

- Safiri, S., Kolahi, A. A., Hoy, D., Buchbinder, R., Mansournia, M. A., Bettampadi, D., . . . Ferreira, M. L. (2020). Global, regional, and national burden of neck pain in the general population, 1990-2017: systematic analysis of the Global Burden of Disease Study 2017. *Bmj, 368*, m791. doi:10.1136/bmj.m791
- Saha, T. K., Dasgupta, A., Butt, A., & Chattopadhyay, O. (2010). Health Status of
 Workers Engaged in the Small-scale Garment Industry: How Healthy are They?
 Indian journal of community medicine : official publication of Indian
 Association of Preventive & Social Medicine, 35(1), 179-182. doi:10.4103/0970-0218.62584
- Sakthi Nagaraj, T., Jeyapaul, R., & Mathiyazhagan, K. (2019). Evaluation of ergonomic working conditions among standing sewing machine operators in Sri Lanka. *International Journal of Industrial Ergonomics, 70*, 70-83. doi:https://doi.org/10.1016/j.ergon.2019.01.006
- Sargent, L. D., & Terry, D. J. (2000). The moderating role of social support in Karasek's job strain model. *Work & Stress, 14*(3), 245-261. doi:10.1080/02678370010025568
- Schibye, B., Skov, T., Ekner, D., Christiansen, J. U., Sjøgaard, G. J. S. J. o. W., Environment, & Health. (1995). Musculoskeletal symptoms among sewing machine operators. (6), 427-434. doi:10.5271/sjweh.58
- Scotland, P. H. (2021). Income inequality.
- Sealetsa, O. J., & Thatcher, A. (2011). Ergonomics issues among sewing machine operators in the textile manufacturing industry in Botswana. *Work, 38*, 279-289. doi:10.3233/WOR-2011-1131
- Sein, M., Howteerakul, N., & Jirachewee, J. (2010). Job Strain among Rubber-Glove-Factory Workers in Central Thailand. *Industrial health, 48*, 503-510. doi:10.2486/indhealth.MS962
- Shan, C. L., Bin Adon, M. Y., Rahman, A. B. A., Hassan, S. T. S., & Ismail, K. B. (2011). Prevalence of neck pain and associated factors with personal characteristics, physical workloads and psychosocial among male rubber workers in FELDA

settlement Malaysia. *Global journal of health science, 4*(1), 94-104. doi:10.5539/gjhs.v4n1p94

- Sibella, F., Galli, M., Romei, M., Montesano, A., & Crivellini, M. J. C. b. (2003). Biomechanical analysis of sit-to-stand movement in normal and obese subjects. *18*(8), 745-750.
- Silvia Monteiro, M., Maria Costa Alexandre, N., Ilmarinen, J., Mendes Rodrigues, C. J. I.J. o. O. S., & Ergonomics. (2009). Work Ability and Musculoskeletal DisordersAmong Workers From a Public Health Institution. *15*(3), 319-324.
- SIU, O. L., Cooper, C. L., & Donald, I. J. S. M. (1997). Occupational stress, job satisfaction and mental health among employees of an acquired TV company in Hong Kong. 13(2), 99-107.
- Skov, T., Borg, V., & Orhede, E. (1996). Psychosocial and physical risk factors for musculoskeletal disorders of the neck, shoulders, and lower back in salespeople. *Occupational and Environmental Medicine*, *53*(5), 351-356. doi:10.1136/oem.53.5.351
- Skovlund, S. V., Bláfoss, R., Sundstrup, E., & Andersen, L. L. (2020). Association between physical work demands and work ability in workers with musculoskeletal pain: cross-sectional study. *BMC Musculoskeletal Disorders,* 21(1), 166. doi:10.1186/s12891-020-03191-8
- Soe, K. T., Laosee, O., Limsatchapanich, S., & Rattanapan, C. (2015). Prevalence and risk factors of musculoskeletal disorders among Myanmar migrant workers in Thai seafood industries. *Int J Occup Saf Ergon, 21*(4), 539-546. doi:10.1080/10803548.2015.1096609
- Søgaard, K., Sjøgaard, G. J. E., & reviews, s. s. (2017). Physical activity as cause and cure of muscular pain: evidence of underlying mechanisms. *45*(3), 136.
- Soori, H., Rahimi, M., & Mohseni, H. J. E.-E. M. H. J., 14 , 697-703,. (2008). Occupational stress and work-related unintentional injuries among Iranian car manufacturing workers.
- Sorić, M., Golubić, R., Milosević, M., Juras, K., & Mustajbegović, J. (2013). Shift work, quality of life and work ability among Croatian hospital nurses. *Coll Antropol, 37*(2), 379-384.

- Steinisch, M., Yusuf, R., Li, J., Rahman, O., Ashraf, H., Strümpell, C., . . . Loerbroks, A. (2013). Work stress: Its components and its association with self-reported health outcomes in a garment factory in Bangladesh-Findings from a cross-sectional study. *Health & place, 24C*, 123-130. doi:10.1016/j.healthplace.2013.09.004
- Storheim, K., & Zwart, J.-A. (2014). Musculoskeletal disorders and the Global Burden of Disease study. *Annals of the Rheumatic Diseases, 73*(6), 949. doi:10.1136/annrheumdis-2014-205327
- Sundquist, J., & Johansson, S.-E. (2000). High demand, low control, and impaired general health: working conditions in a sample of Swedish general practitioners. *Scandinavian Journal of Public Health, 28*(2), 123-131. doi:10.1177/140349480002800208
- Taib, M. F. M., Bahn, S., Yun, M. H., & Taib, M. S. M. J. W. (2017). The effects of physical and psychosocial factors and ergonomic conditions on the prevalence of musculoskeletal disorders among dentists in Malaysia. 57(2), 297-308.
- Tantawy, S. (2019). Work-related musculoskeletal symptoms among employees with different tasks: Ahlia University case study. *Biomedical Research*.
- Tokuda, Y., Ohde, S., Takahashi, O., Shakudo, M., Yanai, H., Shimbo, T., . . . Fukui, T. J. R. i. (2007). Musculoskeletal pain in Japan: prospective health diary study. *28*(1), 7-14.
- Urwin, M., Symmons, D., Allison, T., Brammah, T., Busby, H., Roxby, M., . . . Williams,
 G. (1998). Estimating the burden of musculoskeletal disorders in the community: the comparative prevalence of symptoms at different anatomical sites, and the relation to social deprivation. *Annals of the Rheumatic Diseases, 57*(11), 649. doi:10.1136/ard.57.11.649
- van den Berg, T., Elders, L., de Zwart, B., Burdorf, A. J. O., & medicine, e. (2008). The effects of work-related and individual factors on the Work Ability Index: a systematic review.
- van den Berg, T. I., Alavinia, S. M., Bredt, F. J., Lindeboom, D., Elders, L. A., Burdorf, A. J. I. a. o. o., & health, e. (2008). The influence of psychosocial factors at work

and life style on health and work ability among professional workers. *81*(8), 1029-1036.

- Van den Heuvel, S. G., Geuskens, G. A., Hooftman, W. E., Koppes, L. L., & Van den Bossche, S. N. J. J. o. o. r. (2010). Productivity loss at work; health-related and work-related factors. 20(3), 331-339.
- Van der Doef, M., Maes, S. J. W., & stress. (1999). The job demand-control (-support) model and psychological well-being: A review of 20 years of empirical research. *13*(2), 87-114.
- Vander Elst, T., De Witte, H., De Cuyper, N. J. E. J. o. W., & Psychology, O. (2014). The Job Insecurity Scale: A psychometric evaluation across five European countries. *23*(3), 364-380.
- Viester, L., Verhagen, E. A., Hengel, K. M. O., Koppes, L. L., van der Beek, A. J., & Bongers, P. M. J. B. m. d. (2013). The relation between body mass index and musculoskeletal symptoms in the working population. 14(1), 1-9.
- Waddell, G., & Burton, A. K. (2006). Is work good for your health and well-being?
- Wang, P.-C., Harrison, R. J., Yu, F., Rempel, D. M., & Ritz, B. R. (2010). Follow-up of neck and shoulder pain among sewing machine operators: The Los Angeles garment study. *American Journal of Industrial Medicine*, *53*(4), 352-360. doi:https://doi.org/10.1002/ajim.20790
- Wang, P.-C., Rempel, D. M., Harrison, R. J., Chan, J., & Ritz, B. R. (2007). Workorganisational and personal factors associated with upper body musculoskeletal disorders among sewing machine operators. *Occupational and Environmental Medicine, 64*(12), 806. doi:10.1136/oem.2006.029140
- Wang, P. C., Rempel, D. M., Harrison, R. J., Chan, J., & Ritz, B. R. (2007). Workorganisational and personal factors associated with upper body musculoskeletal disorders among sewing machine operators. *Occupational and Environmental Medicine*, 64(12), 806-813. doi:10.1136/oem.2006.029140
- WHO. (2001). International classification of functioning, disability, and health. ICF Introduction 2001, World Health Organization, Geneva.

Work, E. A. f. S. H. a. (2009). OSH in figures: stress at work — facts and figures.

- Yang, T.-C., Matthews, S. A., & Chen, V. Y.-J. J. I. j. o. b. m. (2014). Stochastic variability in stress, sleep duration, and sleep quality across the distribution of body mass index: insights from quantile regression. *21*(2), 282-291.
- Zajacova, A., & Lawrence, E. M. J. A. r. o. p. h. (2018). The relationship between education and health: reducing disparities through a contextual approach. *39*, 273-289.
- Zimmerman, E. a. S. H. W. (2014). Between Education and Health. NAM Perspectives. Discussion Paper, National Academy of Medicine, Washington, DC. doi:https://doi.org/10.31478/201406a
- Žmauc, T., Železnik, D., & Težak, O. (2019). Work ability index in Slovenian hospital nurses aged over fifty years. *Archives of Industrial Hygiene and Toxicology, 70*, 265-272. doi:10.2478/aiht-2019-70-3291
- Zwart, B., Frings-Dresen, M., & Duivenbooden, J. (2002). Test-retest reliability of the Work Ability Index questionnaire. *Occupational medicine (Oxford, England), 52*, 177-181.



APPENDICES

APPENDIX A: Certificate of ethical approval by Chulalongkorn University

<u>A</u>	AF 02-12 The Research Ethics Review Committee for Research Involving Human Research Participants, Group I, Chulalongkorn University
9	✓ Jamjuree 1 Building, 2nd Floor, Phyathai Rd., Patumwan district, Bangkok 10330, Thailand, Tel: 0-2218-3202, 0-2218-3049 E-mail: <u>eccu@chula.ac.th</u>
	COA No. 201/2021
	Certificate of Approval
Study	Y TITLE No. 156.1/64 : THE WORK-RELATED MUSCULOSKELETAL DISORDERS, PSYCHOSOCIAL FACTORS, WORK PRODUCTIVITY AND WORK ABILITY AMONG GARMENT FACTORY WORKERS IN MYANMAR
Princ	ipal Investigator : TUN WIN OO
Place	e of Proposed Study/Institution : College of Public Health Sciences, Chulalongkorn University
	The Research Ethics Review Committee for Research Involving Human Research Participants,
Grour	L Chulalonekorn University Thailand has approved constituted in accordance with Belmont
Deed	t 1070. Destantion of Halaidi. 2012. Courtil for Istantical Operations of Madian
керо	IT 1979, Declaration of Helsinki 2013, Council for International Organizations of Medical
Scien	ces (CIOM) 2016, Standards of Research Ethics Committee (SREC) 2017, and National Policy
and g	uidelines for Human Research 2015.
	DE 2 sit Rayon Minholm
Signa	ature: Triba Tasamapred Signature: Navenari Tingureane
(As	sociate Prof. Prida Tasanapradit, M.D.) (Assistant Prof. Raveenan Mingpakanee, Ph.D.)
	Chairman Secretary
	enaminari secietary
Date	of Approval : 28 September 2021 Approval Expire date : 27 September 2022
The a	approval documents including:
1)	Research proposal
2)	Participant Information Sheet and the Edina 156. 1/64
3)	Researcher 28 SEP 2021
4)	Questionnaire
4)	Approval Expire Date
The app	proved investigator must comply with the following conditions:
1.	It's unethical to collect data of research participants before the project has been approved by the committee.
Ζ.	to the expired date with submission of progress report.
3.	Strictly conduct the research/project activities as written in the proposal.
4.	Using only the documents that bearing the RECCU's seal of approval: research tools, information sheet, consent form, invitation
	letter for research participation (if applicable).
	Report to the RECCU for any serious adverse events within 5 working days.
5.	
5. 6.	Report to the RECCU for any amendment of the research project prior to conduct the research activities.
5. 6. 7.	Report to the RECCU for any amendment of the research project prior to conduct the research activities. Report to the RECCU for termination of the research project within 2 weeks with reasons.
5. 6. 7. 8.	Report to the RECCU for any amendment of the research project prior to conduct the research activities. Report to the RECCU for termination of the research project within 2 weeks with reasons. Final report (AF 01-15) and abstract is required for a one year (or less) research/project and report within 30 days after the completion of the research/project.
5. 6. 7. 8. 9.	Report to the RECCU for any amendment of the research project prior to conduct the research activities. Report to the RECCU for termination of the research project within 2 weeks with reasons. Final report (AF 01-15) and abstract is required for a one year (or less) research/project and report within 30 days after the completion of the research/project. Research project with several phases; approval will be approved phase by phase, progress report and relevant documents for the
5. 6. 7. 8. 9.	Report to the RECCU for any amendment of the research project prior to conduct the research activities. Report to the RECCU for termination of the research project within 2 weeks with reasons. Final report (AF 01-15) and abstract is required for a one year (or less) research/project and report within 30 days after the completion of the research/project. Research project with several phases; approval will be approved phase by phase, progress report and relevant documents for the next phase must be submitted for review.

APPENDIX B: Questionnaires (English)

Personal factors Qu	lestionnaire
---------------------	--------------

No.	Item		Option to choose
1.	Age (in Full year)	years	
2.	Gender	Male	1
		Female	2
3.	Education Level	primary school	1
		secondary school	2
		high school and	3
		above	4
4.	Marital status	Single	1
	จุฬาลงกรณ์มห	Married	2
	CHULALONGKORN	Separated	3
		Widowed	4
5.	Height	cm	
	Weight	kg	
6.	Monthly income	kyats	
7.	Working hour per day	hours	

8.	Duration of work	years	

BMI calculated in SPSS using height and weight data.

Cornell Musculoskeletal Discomfort Ouestionnaire

The diagram below shows the approximate position of the body parts referred to in the questionnaire. Please answer by marking the appropriate box.			During the last work <u>week</u> how often did you experience ache, pain, discomfort in:					If you experienced ache, pain, discomfort, how uncomfortable was this?			If you experienced ache, pain, discomfort, did this interfere with your ability to work?			
			Never	1-2 times last week	3-4 times last week	Once every day	Several times every day	Slightly uncomfortable	Moderately a uncomfortable	Very uncomfortable	Not at all	Slightly interfered	Substantially interfered	
	\cap	Neck												
	SC	Shoulder	(Right) (Left)											
C		Upper Back												
λ	$\sqrt{1}$	Upper Arm	(Right) (Left)											
		Lower Back												
17	I Y UC	Forearm	(Right) (Left)											
(W)		Wrist	(Right) (Left)											
	$\langle \rangle$	Hip/Buttocks												
	1-1-1	Thigh	(Right) (Left)											
	MN	Knee	(Right) (Left)											
		Lower Leg	(Right) (Left)											
<u></u>	mell University, 2003	Foot	(Right) (Left)											

The diagram below shows the approximate position of the body parts referred to in the questionnaire. Please answer by marking the appropriate box.			Durit how ache,	ng the often , pain,	e last v did ye , disco	work <u>w</u> ou exp omfort	eek erience in:	If you experienced ache, pain, discomfort, how uncomfortable was this?			If you experienced ache, pain, discomfort, did this interfere with your ability to work?			
			Never	1-2 times last week	3-4 times last weck	Once every day	Several times every day	Slightly uncomfortabl	Moderately e uncomfortable	Very uncomfortable	Not at all	Slightly interfered	Substantially	
\cap	Neck													
	Shoulder	(Right) (Left)												
(no cont	Upper Back													
ATAT	Upper Arm	(Right) (Left)												
()) = () + () + () + () + () + () + () +	Lower Back													
$\left \left(\begin{array}{c} \dot{\mathbf{x}} \\ \dot{\mathbf{x}} \\ \end{array}\right)\right $	Forearm	(Right) (Left)												
M R	Wrist	(Right) (Left)												
	Hip/Buttocks													
	Thigh	(Right) (Left)												
	Knee	(Right) (Left)												
WW	Lower Leg	(Right) (Left)												
(Condillairosiy, 1983)	Foot	(Right) (Left)												



Work productivity and Activity Impairment Ouestionnaire

The following questions ask about the effect of your health problems on your ability to work and perform regular activities. By health problems we mean any physical or emotional problem or symptom. *Please fill in the blanks or circle a number, as indicated.*

 1) Are you currently employed (working for pay)?
 _____NO

YES

If NO, check "NO" and skip to question 6.

The next questions are about the **past seven days**, not including today.
2) During the past seven days, how many hours did you miss from work because of <u>your health. Problems</u>? Include hours you missed on sick days, times you went in late, left early, etc., because of your health problems. Do not include time you missed to participate in this study.

HOURS

3) During the past seven days, how many hours did you miss from work because of any other reason, such as vacation, holidays, time off to participate in this study?

HOURS

During the past seven days, how many hours did you actually work?
_____HOURS (If "0", skip to question 6.)

5) During the past seven days, how much did health problems affect your productivity <u>while you were working</u>?

Think about days you were limited in the amount or kind of work you could do, days you accomplished less than you would like, or days you could not do your work as carefully as usual. If health problems affected your work only a little, choose a low number. Choose a high number if health problems affected your work a great deal.

Consider only how much <u>health problems</u> affected productivity <u>while you were working</u>.

Health					H	ealth	prot	olems	5		
had no effect on my					CC	omple	etely	prev	ent	from	working
work	0	1	2	3	4	5	6	7	8	9	10

CIRCLE A NUMBER

Work Ability Index

ls your work						
Psychologically dem	anding?					_
Physically demandin	ę?					0
Physically and psych	ologically demanding	?				0
						0
1. Current work al	pility compared to I	nighest work ability	ever:			
Assume that your work work ability? (0 mean	kability at its best has a s s that you currently c	value of 10 points. How annot work at all) (10	many points w) work ability	vould you at its bes	give your c st)	urrent
0 1	2 3 4	4 5 6	7	8	9	1 0
0 0			0	0	0	0
2. Work ability in	relation to demand	s				
How do you rate yo	our current work ability	y with respect to the	physical dem	ands of y	our work	2
Very good (5)	Rather good (4)	Moderat Moderat e (3)	Rather poo (2)	or	Ve	ry poor (1)
0	0	N UNIVERSITY	0			0
How do you rate y	our current work abi	lity with respect to t	he mental de	emands	of your w	ork?
Very good (5)	Rather good (4)	Moderat e (3)	Rather (2)	r poor)	Ve	ery poor (1)
0	0	0	0			0

3. Current diseases

In the	following list, mark your current diseases or injuries. Also			
indica	te whether a physician has diagnosed or treated these	Yes,	Yes,	Ν
		own	physician's	0
diseas	5es.	opinio	diagnosis	(0
		n (2)	(1))
01	Injury due to an accident	0	0	0
	Musculoskeletal disease in back, limbs or other part of the		0	0
02	body (e.g. repeated pain in joint muscle, sciatica,	0	0	0
	rheumatism, arthritis)			
	Cardiovascular disease			
03	(e.g. hypertension, coronary heart disease)	0	0	0
	Respiratory disease			
04	(e.g. repeated infections of the respiratory tract.	0	0	0
	emphysema)			
	Mental disorder			
05	(e.g. depression, "burn-out", anxiety or insomnia)	0	0	0
	Neurological or sensory disease			
06	(e.g. hearing or visual disease migraine enileosy)	0	0	0
	Disasting disease (and iting			
07	Digestive disease / condition	0	0	0
	(e.g. gastritis, gall stones, liver or pancreatic disease,			
08	Genitourinary disease	0	0	0
00	(e.g. infection in urinary tract, gynecological disease or			
	prostate)			
	Skin disease	0	0	0
09	(e.g. allergic or other rash, varicose veins)			
		0	0	0

10	Tumour or cancer			
11	Endocrine or metabolic disease (e.g. diabetes, severe obesity or gout)	0	0	0
12	Blood diseases (e.g. anemia, other blood disorder or defect)	0	0	0
13	Birth defects	0	0	0
14	Other disorder or disease	0	0	0

4. Estimated work impairment due to diseases									
Is your illness or injury a hindrance to your current job? Check more than one alternative if needed.									
There is no hindrance / I have no diseases.									
I am able to do my job, but it causes some symptoms.									
I must sometimes slow down my work pace or change my work methods.									
I must often slow down my work pace or change my work methods.									
Because of my condition, I feel I am able to do only part time work.									
In my opinion I am en	In my opinion I am entirely unable to work.								
5. Illness within last	year (12 months)								
During the last 12 mor	nths:								
how many whole days	s have you been off v	work because of illne	SS:						
None (5)	Max. 9 days (4)	10 - 24 25 - 99 days 100 - 354 days (2) days (3) (1)							

0	0		0		0	0		
6. Estimation of owr	n work ability	in 2 ye	ears					
Do you believe, accor	ding to your pre	esent s	tate of health,					
hat you will be able t	to do your curre	ent job	two years from nov	v?				
Unlikely (1)	Unlikely (1)				Rel cer	latively tain (7)		
0		NH/	112 - 0			0		
		Q	2					
. Mental capacities	1							
.1 Considering the la	ast three month	ns: Hav	e you been able to	enjoy y	our regular da	ily activities?		
Often (4)	Rather often (3)		Sometimes rat (2) selo		rather Never seldom (0)			
0	0				0	0		
.2 Considering the la	ast three month	ns: Hav	e you been active a	nd alert	?			
Often (4)	Rather often (3)		Sometimes (2)		rather seldom (1)	Never (0)		
0	0	1	าวิทยาลัย 		0	0		
C	HULALONG	KORN	UNIVERSITY					
7.3 Considering the la	ast three month	ns: Hav	e you felt yourself to	o be ful	ll of hope abo	out the future?		
Often (4)	Rather often (3)		Sometimes (2)		rather seldom (1)	Never (0)		
0	0		0		0	0		

l

	Item	Strongly	Disagree	Agree	Strongly
		disagree			Agree
1.	My work require me to learn new things.				
2.	There are a lot of repetative task in my work.				
3.	My work require me to be creative.				
4.	My work allow me to make my own decision.	. A A			
5.	My work require a high level of skills.				
6.	On my work, I have a very little freedom to				
	decide how I do on my work.	3			
7.	I can do a variety of different things on my job.	ลัย			
8.	On things that happened at work ,my opinion	RSITY			
	are influencial.				
9.	My work provide rooms for me to develop my				
	own talent.				
10.	My work requires me to do things very quickly.				
11.	My work requires me to do be very hard				

	working.		
12.	My workload is not considered excessive.		
13.	I have enough time to accomplish my work.		
14.	On my job, I am not asked by different people		
	to do things that are contradictory.		
15.	My work require rapid physical activity.		
16.	My work make awkward body position.	A	
17	My work is a steady work.		
18	My work is a security work.		
19	In the future, I am likely to be lay off.	2	
20	My supervisor gives concern to staff's welfare.	RSITY	
21.	My supervisor listen to my opinion.		
22.	My supervisor provides assistance to staff.		
23.	My supervisor organizes well the staff member		
	as to facilitate tasks at work.		
24.	My coworkers attend their duties well.		

25.	My coworker show their care for me.		
26.	My coworker are friendly.		
27.	My coworker are helpful.		

Calculation formula and possible score of JCQ

		
Scale	Formula	Possible
		score
Job Control (9)	= skill discretion + decision authority	24–96
Skill discretion (6)	[Q1+Q3+Q5+Q7+Q9+(5-Q2)]×2	12–48
Decision authority (3)	[Q4+Q8+(5-Q6)]×4	12–48
Psychological job demand(5)	3×(Q10+Q11)+2×[15–(Q12+Q13+Q14)]	12–48
Workplace social support(8)	= supervisor support + coworker support	8-32
Supervisor support (4)	Q20+Q21+Q22+Q23	4–16
Co-worker support (4)	Q24+Q25+Q26+Q27	4–16
Physical job demand (2)	Q15+Q16	2–8
Job insecurity (3)	Q19+[10-(Q17+Q18)]	3-12

APPENDIX C: Participant information sheet and consent form (English)

1. Introduction

My name is Tun Win Oo, a postgraduate student at Chulalongkorn University in Bangkok, Thailand. As a requirement to fulfil academic requirements of the university, we are required to do a research and submit a thesis. I have decided to do my study on the association of psychosocial risk factors, work productivity and work ability on work related musculoskeletal disorders of sewing machine workers in Garment factory in Myanmar

Name of the researcher: Mr. Tun Win Oo

Phone number: +95943020529

Email address: <u>tunwinoo430@gmail.com</u>

Objectives of the research

i. To find out personal factors, work productivity, work ability, psychosocial factors and work related musculoskeletal disorders.

งหาลงกรณ์มหาวิทยาลัย

ii. To find the association between personal factors, work productivity, work ability, psychosocial factors to work related musculoskeletal disorders.

2. How to provide information by whom and how to obtain consent

You will be requested to participate in this study by the researcher by giving the questionnaire form and explained to research participant and let them self-administered. This study is entirely voluntary and if you are eligible of our inclusion criteria you can participate in this study.

3. Detail of the participants

In this study, which will include 400 sewing machine operators in Garment factory in Bago Township, Bago Region, Myanmar. You can continue on reading this information sheet and after weighing out the benefits and risk if any, you are free to decide whether to continue the study or not. If you need further clarification on anything, please do not hesitate to ask the researcher both before, during and after the study.

4. Details of screening process of inclusion/exclusion criteria or qualifications inclusion and exclusion criteria will provided to factory and they will perform these criteria and select research participants. Moreover, during recruitment, before explaining research process, and questionnaire, confirm again these criteria by researcher. After screening process, researcher will invite the potential participant to research projects through factory authorities. Inclusion Criteria - Those who worked at least 1 year as sewing operators - Those who are more than 18 year of age- Those who free from disability - Those who suffered from injury, accident, or undergo surgery during the last 3 months - Those who were pregnant (female respondent) - Those who were taking any prescribed medicine by self (or) from a doctor

5. Procedure upon participants วณ์มหาวิทยาลัย

The research involves filling out a self – administered questionnaire which will take about 30 minutes. There are five sections of questionnaire which are (1) Personal factors (2) Work related musculoskeletal disorders related to body pain due to work (3) Work productivity related to hours (4) Psychosocial Questionnaire (5) Work ability Questionnaire related to worker efficiency. The whole questionnaire uses filling blank, tickling in multiple choice and circling in boxes for the ease of the participants in response to questionnaire. Your answers are anonymous.

6. Risks and Harms

There may be a slight risk that you may share some personal or confidential information by chance or that you may feel uncomfortable about talking about some of the topics about psychosocial factors. However, we do not wish this to happen, and you may refuse to answer.

7. Benefits

The study will provide benefit to you, as the study about sewing workers. The study aim to find out the psychosocial problem, work ability, work productivity and work related musculoskeletal disorders among sewing workers. Therefore, the result of the study can provide essential research data that is needed to get better working condition, better well-being of sewing workers by occupational health intervention and better labor law.

8. Confidentiality

Any information that is linked to you will be kept confidentially. Your names or other identifying information will not be mentioned in the report or summaries of the study. The final report can be available from the researcher and this report will be used for only fulfilment of the academic requirement of the Master of Public health degree. All the data will be kept confidential and will not disclose to anyone

9. How researcher will manage with personal data after research project is completed.

The information papers will be burn and delete the input data in my computer forever that I have used within one year after completion of the study period.

10. Compensation There is nothing to give participants but only heartfelt thanks.

11. Voluntary Participation

After reading the information sheet and receiving further explanation on the unclear part of the questionnaire from the researcher if necessary, the participant will be required to give a written inform consent of the selected participant according to inclusion and exclusion criteria. The consent forms will be kept safely by researcher. If you do not want to participate in this study, you do not need to give consents and you do not need to explain anything as a reason. You can withdraw from the study at any time as you wish with no need to give reason and it will not have any negative impact upon the participants

12. Right of the participant and consent I have read the objectives of this research, what I will engage in details, benefits and risk (if there is any) of this research, and the rights and duties of the participants. I have been given the contact details of the researcher. I have read the information sheet and the researcher has explained me and guaranteed to act as indicated in the information sheet. I clearly understand with satisfaction. I willingly agree to participate in this research will not contain any name or identifying information of me. It will take about 30 minutes to answer questionnaire. You can also ask anything you want to know before, during and after the study conduct any time. The principal researcher can be reached at all time with given address mentioned above. If the researcher has new information regarding benefit or risk/harm, you will be informed as soon as possible.

If researcher does not perform upon you as indicated in the participant information sheet and consent form, you can report the incident to the Research Ethics Review Committee for Research Involving Human Research Participants, Group I, Chulalongkorn University (RECCU) Jamjuree 1 Bldg., 254 Phyathai Rd., Patumwan district, Bangkok 10330, Thailand, Tel./Fax. 0-2218-3202, 0-2218-3049 E-mail: eccu@chula.ac.th". I have read the information in this consent form. Furthermore, I have received a copy of participant's information sheet and informed consent form. I, who signed here consent to participate in this study (____)

Sign of the participant Tun Win Oo (Principal researcher)

Associate Professor Nutta Taneepanichskul (Thesis Advisor)



Chulalongkorn University

APPENDIX C: Questionnaire (Myanmar)

ကြွက်သားနှင့်ကျောရိုးမဆိုင်ရာမအီမသာဖြစ်ခြင်း၊ လုပ်ငန်းခွင်အတွင်းစိတ်ပိုင်း၊လူမှုရေးပိုင်းဆိုင်ရာ သွင်ပြင်လက္ခဏာများ၊အလုပ်လုပ်နိုင်စွမ်း၊လုပ်ငန်းခွင်ထုတ်လုပ်နိုင်စွမ်းနှင့် လုပ်ဆောင်မှုထိခိုက်ခြင်းမေးခွန်းလွှာ။

အပိုင်း	ာ ဖြေဆိုသူ၏နောက်ခံအချက်အလက်		
စဉ်	အကြောင်းအရာ	ရွေးချယ်ခြင်း	မှတ်ချက်
1.	အသက် (ပြည့်ပြီးနှစ်)	<u>ه</u> و هه	
2.	ශීදි	ကျား 1	
		Θ 2	
3.	ပညာရေးအဆင့်	သူငယ်တန်းမှ ၄တန်းအထိ 1	
		၅ တန်းမှ စတန်းအထိ 2	
		၉တန်းနှင့်အထက် 3	
4.	အိမ်ထောင်ရေးအခြေအနေ	အိမ်ထောင်မရှိ 1	1
		အိမ်ထောင်ရှိ 2	
		အိမ်ထောင်ကွဲ/ကွာရှင်း 3	
		အိမ်ထောင်ဖက်ကွယ်လွန် 4	
5.	စက်ရုံမှ လက်ရှိ ရရှိသော လစဉ်ဝင်ငွေ (စုစုပေါင်း)	თებ	
6.	တစ်ရက်စက်ချုပ်အလုပ်လုပ်ချိန် (နာရီ၊ဗိနစ်)	နာရီမြနစ်	
7.	စက်ချုပ်သမားအဖြစ်လုပ်ခဲ့သော စုစုပေါင်းနှစ်ကာလ (နှစ်၊လ)	နစ် လ	
8.	သင်၏ Body Mass Index (BMI) (သုတေသီမှဖြ	ည့်စွက်ရန်)	
Α.	အရပ်အမြင့် Height (cm)		
В.	ကိုယ်အလေးချိန် Weight (kg)		
		kg	
C.	BMI		
		.	

အဝိုင်း၂ ကြွက်သားနှင့်ကျောရှိ	းမဆိုင်ရာမေးခွန်းလွှာ (အမျိုးသမီးေ	ဖြဆိုသူ	များအဝ	၃က်)						901 - 631		
အောက်ပါပုံကမေးခွန်းပါကိုယ်နွော၏ ပါသည်။ သင့်လျော်သောအကွက်များအတွင်း ((အမျိုးသမီးဖြေဆိုသူများအတွက်)	ပံပိုကမေးခွန်းပဲကိုယ်ခန္ဓ၁၏ယေဘူယျအစိတ်အပိုင်းများကိုရည်ညွှန်း ၊ ဂူဝ်သောအကွက်များအတွင်း (=) အမှတ်အသားပြု၍ဖြေဆိုပါ။ သမီးဖြေဆိုသူများအတွက်)		ပြီးခဲ့သောအလုပ်တစ်ပတ်အတွင်းသင် မည်မျှ နာကျင်၊ ကိုက်ခဲ၊ မအီမသာ ဖြစ်ခဲ့သလဲ။				သင် နာကျင်၊ ကိုက်ခံ၊ ဗအီမသာ ဖြစ် ခဲ့ပါက မည်မျှဆိုးဝါးခဲ့သလဲ။			သင် နာကျင်၊ ကိုက်ခဲ၊ မအီမသာ ဖြစ် ခဲ့ပါက သင်၏အလုပ်လုပ်နိုင်စွမ်း အပေါ် အနောက်အယှက် ဖြစ်ခဲ့ပါ သလား။		
0		မဖြစ်ပါ။	ဖြီးခဲ့သော အပတ် ၁–၂ ကြိမ်	ဖြီးခဲ့သော အမတ် ၃-၄ ကြိမ်	တစ်နေ့တကြိမ်	နေဝိုင်း အကြိမ်များစွာ	အနည်းငယ်	အသင့်အတင့်	းခဲ့အမင်း	ရော့တဲ့။	အနည်းငယ်ဖြစ်	အတော်အတန် ဖြစ်
	- လည်ပင်။											
AT	ု ပခုံး (ညာ) (ဘယ်)											
IV VI	- ကျော (အပေါ်ဘက်)											
IAA	လက်မောင်း (ညာ) (ဘယ်)											
111 -111	- ကျော (အောက်ဘက်)											
21215	လက်ဖျံ (ညာ) (ဘယ်)											
1 10	လက်ကောက်ဝတ် (ညာ) (ဘယ်)	8										
IX	တင်ပါး/ တင်ပါးဆုံ											
1-1-1	ပေါင် (ညာ) (ဘယ်)											
	ု ဒူး(ခေါင်း) (ညာ) (ဘယ်)											
	ခြေသလုံး (ညာ) (ဘယ်)											
	_ ခြေထောက် (ညာ) (ဘယ်)											

1967	11	11	CONTRACTOR OF A		
1	11	Mr.		P 1 1 1 1 10.	-
	11			8 III I II	

အပိုင်း၂ ကြွက်သားနှင့်ကျောရိုးမဆိုင်ရာမေးခွန်းလွှာ (အဖ	မျိုးသားဖြေး	ဆိုသူများး	ജറുന്)								
ခောက်ပါပုံကမေးနွန်းပါကိုယ်ခန္ဓာ၏ယေဘူယျအစိတ်အပိုင်းများကို၊ ည်ညွှန်းပါသည်။ သင့်လျော်သောအကွက်များအတွင်း (=) အမှတ်အသားပြု၍ဖြေဆို (အမျိုးသားဖြေဆိုသူများအတွက်)	ရ ပြီးခဲ့ေ မည်ရွှ ပါ။ မအိမ္မ	ပြီးခဲ့သောအလုပ်တစ်ပတ်အတွင်းသင် မည်ရွှ နာဂဂျင်၊ ကိုက်ခဲ၊ မအီမသာဖြစ်ခဲ့သလဲ။				သင် နာကျင်၊ ကိုက်ခဲ၊ မအီသော ဖြစ် ခဲ့ပါက မည်မျှဆိုးဝါးခဲ့သလဲ။			သင် နာကျင်၊ ကိုက်ခဲ၊ မအီမသာ ဖြစ် ခဲ့ပါက သင်၏အလုပ်လုပ်နိုင် စွမ်း အပေါ် အနှောက်အယှက် ဖြစ်ခဲ့ ပါသလား။		ခဲ၊ က င် စွမ်း ယှက်
\cap	မဖြစ်ပါ။	ပြီးခဲ့သော အပတ် ၁–၂ ကြိုန်	မြီးခဲ့သော အပတ် ၃-၄ ကြို့န်	တစ်နေ့တကြိမ်	နေတိုင်း အကြိမ်များစွာ	အနည်းငယ်	အသင့်အတင့်	သူမဆန်ပူးဆ	ရဖြစ်ပါ။	အရည်းလမ်ဖြစ်	အတော်အတန် ဖြစ်
လည်ပင်း											
ပခုံ၊ (ညာ)											
(ဘယ်)											
ကျော (အပေါ်ဘက်)											
A A CONSERVER (200)										<u> </u>	
	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>
ကျော (အောကဘက)	- <u>+</u> - <u>+</u> - <u>+</u> -	<u> </u>	<u> </u>	<u> </u>	_ <u>H</u> _	<u> </u>	_ <u></u>	<u> </u>	<u> </u>	_ <u></u>	<u> </u>
		H	H	H	H		H	H		H	H
4 0 wn/emp/000 (200)		<u> </u>		_				<u> </u>	-	_ <u>_</u>	— <u> </u>
())	1 =	Ē	Ē		ē			Ē		Ē	
တင်ပါ။/ တင်ပါ။ဆုံ											
ပေါင် (ညာ)											
(ဘယ်)											
ျူး(ခေါင်း (ညာ)											
(300)			<u> </u>	<u> </u>	<u> </u>	<u> </u>			<u> </u>		
(ဆယ်)		H	H	H	H		H	H		님	님
ခြေထောက် (ညာ)		8				-		Ē	8		

အဝို	င်း ၃ လုပ်ငန်းခွင်အတွင်းစိတ်ပိုင်းလူမှုရေးပိုင်းဆိုင်ရ	ႃၣသွင်ပြင်လ	က္ခဏာများ		
ညွှန်	က္ဘားချက်။ သင်လက်ခံထားမှုနှင့်နီးစပ်သည်ဟုသင်ယူဆ	ာသောအခြေဒ	စနေကို (✔) [ခြစ်ပါ။	
စဉ်	အကြောင်းအရာ	အလွန်	သဘော	သဘော	အလွန်
		သဘော	မတူ	മ	သဘော
		မတူ			တူ
1.	ကျွန်ုပ်၏အလုပ်သည် အသစ် အသစ်သောများကို				
	သင်ကြားရန်လိုအပ်သည်။				
2.	ကျွန်ုပ်၏အလုပ်တွင် ထပ်ခါတလဲလဲလုပ်ရသော				
	အလုပ် များ အများအပြားပါဝင်သည်။				
3.	ကျွန်ုပ်၏အလုပ်သည် တီထွင်ဖန်တီးမှုများပါဝင်ပါ				
	သည်။				
4.	ကျွန်ုပ်၏အလုပ်တွင်ကိုယ်ဝိုင် ဆုံးဖြတ်ချက် များ				
	ချမှတ်နိုင်သည်။				
5.	ကျွန်ုပ်၏အလုပ်က ကျွမ်းကျင်မှု များစွာ				
	လိုအပ်ပါသည်။				
6.	ကျွန်ုပ်၏အလုပ်တွင် ဓိဓိအလုပ်နှင့်ပတ်သက်၍				
	ကိုယ်တိုင်ဆုံးဖြတ်၍ လွတ်လပ်ခွင့် များ အနည်းငယ်				
	သာပါဝင်ပါသည်။				
7.	ကျွန်ုပ်၏အလုပ်တွင်အလုပ်အမျိုးအစားများစွာပါဝင်				
	အလုပ်လုပ်နိုင်ပါသည်။				
8.	ဓိဓိ၏အလုပ်နှင့်ပတ်သက်၍တစ်စုံတစ်ခုဖြစ်ပျက်ခဲ့				
	ပါက မိမိ၏ ပြောရေးဆိုခွင့်မှာအခရာကျပါသည်။				
9.	ကျွန်ုပ်၏အလုပ်က ဓိဓိ၏ကိုယ်ပိုင် အရည် အသွေး				
	များ တိုးတက်စေရန်အတွက် အခွင့်အလမ်း ကို				
	ပေးအပ်ထားသည်။				
10.	ကျွန်ုပ်၏အလုပ်က အလုပ်ကိုလျှင်လျှင်မြန်မြန်ပြီးစီး				
	အောင်လုပ်ရန် လိုအပ်သည်။				
11.	ကျွန်ုပ်၏အလုပ်က အလုပ်ကို အလွန်ကြိုးကြိုး				
	စားစား လုပ်ရန် လိုအပ်သည်။				
12.	ကျွန်ုပ်၏အလုပ်သည် အလွန်များပြားလွန်းသည်ဟု				
	မယူဆပါ။				
13.	ကျွန်ုပ်၏အလုပ်ပြီးမြောက်ရန်အချိန်အလိုအလောက်				
	ရှိသည်။				
14.	ကျွန်ုပ်၏အလုပ်တွင် လူအမျိုးမျိုးက တစ်ခုနှင့်တစ်ခု				
	ဆန့်ကျင်ဘက် ဖြစ်သော အရာများကိုလုပ်ရန်				
	ဓဓကုခုင်းစေခြင်းမျိုးမရှိပါ။				-
15.	ကျွနုပ၏အလုပ်က မြန်ဆနံသောကိုယ်ကာယ ေ၀၀၀ ေ				
	လှုပရှားမှု မျိုးကိုလုံအပသည်။				
16.	ကျွနုပ၏အလုပ်က သက်တောင့်သက်သာမရှိသော				

	ကိုယ်ခန္ဓာအနေအထားကိုဖြစ်စေသည်။	
17.	ကျွန်ုပ်၏အလုပ်က ပုံမှန်အေးအေးဆေးဆေး	
	အလုပ်မျိုး ဖြစ်သည်။	
18.	ကျွန်ုပ်၏အလုပ်ပြုတ်ခြင်းနှင့် ပတ်သက်၍ မစိုးရိမ်	
	ရပါ။	
19.	အနာဂတ်တွင်ကျွန်ုပ်သည်အလုပ်မှ အနားယူချင်ယူ	
	(အလုပ်ထွက်) လိမ့်မည်။	
20.	ကျွန်ုပ်၏ကြီးကြပ်သူသည်အလုပ်သမားများ၏	
	အရေး ကိစ္စကိုအလေးထားသည်။	
21.	ကျွန်ုပ်၏ကြီးကြပ်သူသည်မိမိ၏ထင်မြင်ချက်များကို	
	နားထောင်သည်။	
22.	ကျွန်ုပ်၏ကြီးကြပ်သူသည်အလုပ်သမားများကို	
	ကူညီ ပေးသည်။	
23.	ကျွန်ုပ်၏ကြီးကြပ်သူသည် အလုပ်များ ပြီးမြောက်	
	စေရန် အတွက် အလုပ်သမားအဖွဲ့ဝင်များကို	
	ကောင်း မွန်စွာ စည်းရုံးသည်။	
24.	ကျွန်ုပ်၏လုပ်ဖော်ကိုင်ဖက်များသည်သူတို့အလုပ်ကို	
	ကောင်းမွန်စွာလုပ်ကိုင်ကြသည်။	
25.	ကျွန်ုပ်၏လုပ်ဖော်ကိုင်ဖက်များသည် မိမိကို ဂရုစိုက်	
	ကြောင်းထုတ်ဖော်ပြသကြသည်။	
26.	ကျွန်ုပ်၏လုပ်ဖော်ကိုင်ဖက်များသည်ဖော်ရွေကြ	
	သည်။	
27.	ကျွန်ုပ်၏လုပ်ဖော်ကိုင်ဖက်များသည်ရိုင်းပင်းကူညီ	
	ကြသည်။	

အဝိုစ်	င်း ၄ အဂ	ပုပ်လုပ်နိုင်	စွမ်းအ	သွှန်းကိန်း		1				
စဉ်	အကြော	င်းအရာ				ရွေးချယ်ခြင်း				မှတ်ချက်
	သင့်အဂ	ပုပ်သည်								
•	စိတ်ဝိုင်း	ဆိုင်ရာဖိစီး	မှုများရှိဖ	ပါသလား။		ရှိပါသဥ မရိပါ	ည်	1		
•	ຄຽຽວະເອ	ဝိင်ရာ (ခန	ეოზაზ	င်းဆိုင်ရာ) (ກວເມ	ရိပါသူ	ຽ	- 1		
	30000	ည်မများရှိသ				မရှိပါ	_	2		
•	စိတ်ဝိင်း	ဆိုင်ရာနှင့်	ခန္ဘာကိုဖ	ນໍຽີ້ີ່ ເວັ້າສີ່	,	ရိပါသူ	- 			
	နှစ်ခုစလ	းဖိစ်းမှုများ	င့် ခန္ဓာကိုယ်ပိုင်းဆိုင်ရာ ဂူးရိသလား။				-	2		
1.	လက်ရှိဒ	လုပ်လုပ်န	နိုင်စွမ်းရ	သိနှင့် အမြင့်	မားဆုံးလုပ်နိ	င်စွမ်းရဉ	ည်ကိုနိုင်	င်းယှဉ်ခြင်	:	
2	သင်၏ဒ	ကောင်းဆ	မ်းလုပ်လု	ပ်နိုင်စွမ်းကို	အမှတ် (၁၀)	ဟုဆိုပါ	စို့။ သင်	၏လက်ရှိ	300	လုပ်လုပ် နိုင်စွမ်းကို
	(ဝ မှ ၁ဝအထိ) ပေးရမည်ဆိုပါက သင်အမှတ်မည်မျှ ပေးမည်နည်း။									
	0 1 2 3 4 5						7	8	9	10
	0									10
	လုံးဝအလုပ်ကောင်း							အကောင်	င်းဆုံး	လုပ်နိုင်ပါသည်။
	ကောင်းမလုပ်နိုင်ပါ။									
	0 0		c c	<u>ېر</u>	ပတတစခုကိုး	သာကွင်း	Olii			
2.	လိုအပဲခု	က်အရအဖ	<u>v</u> ovo	နိုင်စွမ်းရည်		00	00			· · · · · · · ·
Α.	သင်၏ဝ	ഗന് ദ് ജസ് ം ം	ပလုပ်နို	င်စွမ်းကို သ	င့်အလုပ်၏က	ာက္ဝင္း	ဆိုင်ရာ	အားထုတ	မှုနှင့်	ု နူငးယုဉ လျှင
6	သငမည	သူသတမှင	ာမညန	ມະ ແ	S	\$2		\$	s	<u>8-9</u> 1
	အလွနေခ	ကာငးမွန	നെ	၁ငးမွနစ္စာ ၇၉၈၃၈၃။	အသင့်အဝ	ာင္ငံသာ	ແລະໃຊ້ເພື່			အလွနဆုးဝါးနေ
	မွာလုပ	နင်သည်။	လုပ	နင်သည်။	00403	သည။	မလုပ်နိုင်ပစ			ປາລາວ 1
B	သင်း၏	3 030000	Second	4 80.2029	3	ပ်၏စိတ်ပိုင်းဆိုင်ရာဖိုစီးမနှင့် နိုင်းယဉ် လျှင်				2400 240029
ь.	SDC6II	COC) glass	Sugar	နိုင်စိုမ်းကို	သင့်အလိုပ	GILOOOC	ုပ်ဆု	ငရာဖစ်းရှိ	ရှင့်	မှိုင်းယှဉ် လျှင
5	Jucon	82200	ကမည်ရ နည်း	ມະຫ 	ကသင်ကက	2.83	8~	Sacion		ကားနိုက်ကုန်းခဲ
	အလွေန	စတချမ်း လုပ်ထိုင်	000	ရမ်းသာမှာ	30,300	သူတို့။	စတဆင်းရစွာ			အလွနစတဆင်းရ
	2860	ပြာသိ။	L CQOq	CONTR	ရဖာသာပါ	ມ _ິ ພ"	လုပ်နေရပါသည်။		<u> </u>	မားပုံလာပ်နေရ ပါသည်။
-	quo	5 5	-	4	3			2	_	1
3.	ရောဂါအ	- ခြေအနေန	င်လပ်နိ	င်စမ်းအားက	ပါဆင်းမ					-
-	သင်၏စ	ရာဂါဝေဒန	ာ သိမဟ	စ တ် ထိခိက်း	u ။ ဒဏ်ရာရမက	သင်၏ဂ	ပက်ရှိ	အလပ်ကိ	8	က် နေ ပါသလား။
	အောက်	ပါအခြေအဖ	နေများဂ	ှ ဝစစ်ဆေးကြ	ည်ပါ။		л			
	ညွှန်ကြား	ချက်။ သင်o	လက်ခံထ	ားမှုနှင့်နီးစပ်ခ	ပည်ဟုသင်ယူး	ဆသောဒ	ခြေအ	နေတစ်ခု (တစ်ခု	ခုထက် ဝိုနိုင်သည်)
	ကို ကွင်း	ဝဝါ။		100 8	55967 - 835		1991	al 15		an tana tana

•	ကျွန်ုပ်တွင်ရောဂါမ	ရှိပါ။ ကောင်းစွာအလ	ပ်လပ်နိုင်သည်။			6				
•	ကျွန်ုပ်အလုပ်လုပ်နို	။ ၀၀၈ ကိုသည်။သို့သော်ရောဂ	ာါလက္ခဏာများခံစား	နေရသည်။		5				
•	တစ်ခါတစ်ရံကျန်ပ်၏အလုပ်ပံစံကြောင်တစ်ခါတစ် လေအလုပ်လုပ်နိုင်နန်း ကို လောခ၊									
	ရပါသည်။									
•	ရံဖန်ရံခါကျွန်ုပ်၏အလုပ်လုပ်နိုင်နှန်းကိုလျှောချ လိုက်ရ၍ အလုပ်လုပ်သည်ပုံစံကို									
	ပြောင်းလဲပစ်လိုက်ရ	ရပါသည်။				3				
•	ကျွန်ုပ်၏ရောဂါအခ	ခြအနေများကြောင့်က	ဂျွန်ုပ်သည်အချိန်ပိုင်း	အလုပ်များသာလုပ်နိ	င်သည်ဟု					
	ခံစားရပါသည်။					2				
•	ကျွန်ုပ်၏အခြေအခေ	နအရကောင်းမွန်စွာအ	လုပ်လုပ် နိုင်ခြင်းမ	ပါ။		1				
4.	ပြီးခဲ့သောနှစ်ကနေ	မကောင်းဖျားနာမှုမျာ	း (၁၂ လအတွင်း)							
	ကျွန်ုပ်သည်လွန်ခဲ့ေ ပျက်ကွက်ခဲ့သနည်း	သာ (၁၂) လအတွင်း ။	နေမကောင်း၊ဖျားနာ	ရောဂါများကြောင့် အ	လုပ်ရက် မ	ည်မျှ				
	မရှိပါ။ အများဆုံး ၉ ရက် ၁၀ မှ ၂၄ ရက် ၂၅ မှ ၉၉ ရက် ၁၀၀–၃၅									
	5	4	3	2	1	3				
5.	လာမည့် (၂) နှစ်ကာ	ာလ အလုပ်လုပ်နိုင်စွမ်	န်းခန့်မှန်းချက်		<u>i</u> .					
	ယခုလက်ရှိကျန်းမာ	ာရေးအခြေအနေအရ	နောက် (၂) နှစ်ကြာ	ခဲ့လျှင် ယခုလက်ရှိ အ	လုပ်ကို					
	ဆက်လက်လုပ်ကိုင်	နိုင်မည်လား။	C GL - GY PY PAL 949	P CAR PRON THE	100 850					
	မလုပ်နိုင်	ပါ။	မသေချာပါ။	မလုပ်နိုင်ပါ။ မသေချာပါ။ လုပ်နိုင်ပါမည်။						
	1 4 7				ι L					
	1		4		7					
6.	1 စိတ်ပိုင်းဆိုင်ရာစွမ်း	ဆောင်နိုင်ခြင်း	4		7					
6. A.	1 စိတ်ပိုင်းဆိုင်ရာစွမ်း လွန်ခဲ့သော (၃) လ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် :	4 အလုပ် များကို စိတ်ဖ	ပျာ်ရွှင်စွာလုပ်ကိုင်နိုင်	7 ငိပါသလား။					
6. A.	1 စိတ်ပိုင်းဆိုင်ရာစွမ်း လွန်ခဲ့သော (၃) လ ခဏခဏ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် : မကြာမကြာ	4 အလုပ် များကို စိတ်ဖ တစ်ခါတစ်လေ	- ပျာ်ရွှင်စွာလုပ်ကိုင်နိုင် ဖြစ်တောင့်ဖြစ်ခဲ	7 နိပါသလား။ ဘယ်သော	ာအခါမျှ				
6. A.	1 စိတ်ပိုင်းဆိုင်ရာစွမ်း လွန်ခဲ့သော (၃) လ ခဏခဏ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် : မကြာမကြာ ဆိုသလို	4 အလုပ် များကို စိတ်ရ တစ်ခါတစ်လေ	- ပျာ်ရွှင်စွာလုပ်ကိုင်နို ဖြစ်တောင့်ဖြစ်ခဲ	7 ငိပါသလား။ ဘယ်သေး	ာအခါမျှ				
6. A.	1 စိတ်ပိုင်းဆိုင်ရာစွမ်း လွန်ခဲ့သော (၃) လ ခဏခဏ 4	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် : မကြာမကြာ ဆိုသလို 3	4 အလုပ် များကို စိတ်ရ တစ်ခါတစ်လေ 2	ပျာ်ရွှင်စွာလုပ်ကိုင်နို ဖြစ်တောင့်ဖြစ်ခဲ 1	7 7 င်ပါသလား။ တယ်သေး	ာအခါမျှ				
6. A. B.	1 စိတ်ပိုင်းဆိုင်ရာစွမ်း လွန်ခဲ့သော (၃) လ ခဏခဏ 4 လွန်ခဲ့သော (၃) လ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် မကြာမကြာ ဆိုသလို 3 အတွင်း သင်တက်တ	4 အလုပ် များကို စိတ်ရ တစ်ခါတစ်လေ 2 က်ကြွကြွ လန်းလန်း	ပျာ်ရွှင်စွာလုပ်ကိုင်နို ဖြစ်တောင့်ဖြစ်ခဲ 1 ဆန်းဆန်းရှိပါသလ	7 ເປັນເຊິ່ງ ອັນເຊິ່ງ ອັນເຊິ່ງ ເຊິ່ງ ເຊິ່ງ	ာအခါမျှ				
6. A. B.	1 ຈິတົງຊີ င်းဆိုင်ရာစွမ်း လွန်ခဲ့သော (၃) လ ອဏອဏ 4 လွန်ခဲ့သော (၃) လ ອဏອဏ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် : မကြာမကြာ ဆိုသလို 3 အတွင်း သင်တက်တ မကြာမကြာ	4 အလုပ် များကို စိတ်ရ တစ်ခါတစ်လေ 2 က်ကြွကြွ လန်းလန်း တစ်ခါတစ်လေ	ပျာ်ရွှင်စွာလုပ်ကိုင်နိုး ဖြစ်တောင့်ဖြစ်ခဲ 1 ဆန်းဆန်းရှိပါသလာ ဖြစ်တောင့်ဖြစ်ခဲ	7 ອ້ບໃວແດນາະຫ ອາເມີຣວນ: 0 ກະຫ ອາເພ	ာအခါမျှ				
6. A. B.	1 ຈິတົຈິໃ້ເວັາຊີໂຣ້ຄາຊິຣໍ່ສິ ດູງຊໍ້ອຸ້ອີລນາ (၃) ເບ ອຫາອຫກ 4 ດູງຊໍ້ອຸ້ອີລນາ (၃) ເບ ອຫາອຫກ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် : မကြာမကြာ ဆိုသလို 3 အတွင်း သင်တက်တ မကြာမကြာ ဆိုသလို	4 အလုပ် များကို စိတ်စ တစ်ခါတစ်လေ 2 က်ကြွကြွ လန်းလန်း တစ်ခါတစ်လေ	းပျာ်ရွှင်စွာလုပ်ကိုင်နို ဖြစ်တောင့်ဖြစ်ခဲ 1 ဆန်းဆန်းရှိပါသလာ ဖြစ်တောင့်ဖြစ်ခဲ	7 ອັດເມືອງ ອັດເມືອງ 0 ນະແຫຼງ ອັດເມືອງ	ာအခါမျှ ဘာခါမျှ				
6. A. B.	1 စိတ်ပိုင်းဆိုင်ရာစွမ်း လွန်ခဲ့သော (၃) လ ခဏခဏ 4 လွန်ခဲ့သော (၃) လ ခဏခဏ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် မကြာမကြာ ဆိုသလို 3 အတွင်း သင်တက်တ မကြာမကြာ ဆိုသလို 3	4 အလုပ် များကို စိတ်ရ တစ်ခါတစ်လေ 2 က်ကြွကြွ လန်းလန်း တစ်ခါတစ်လေ 2	ပျာ်ရွှင်စွာလုပ်ကိုင်နို ဖြစ်တောင့်ဖြစ်ခဲ 1 ဆန်းဆန်းရှိပါသလ ဖြစ်တောင့်ဖြစ်ခဲ 1	7 ວ່າພີຂອງ ອາເພີຂອງ 0 ອາເພີຂອງ 0 ອາເພີຂອງ 0 ອາເພີຂອງ 0 ອາເພີຂອງ 0 ອາເພີຂອງ 0	ာအခါမျှ သအခါမျှ				
6. A. B.	1 စိတ်ပိုင်းဆိုင်ရာစွမ်း လွန်ခဲ့သော (၃) လ ခဏခဏ 4 လွန်ခဲ့သော (၃) လ ခဏခဏ 4 လွန်ခဲ့သော (၃) လ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် မကြာမကြာ ဆိုသလို 3 အတွင်း သင်တက်တ မကြာမကြာ ဆိုသလို 3 အတွင်းသင်၏အနာက	4 အလုပ် များကို စိတ်ရ တစ်ခါတစ်လေ 2 ကိကြွကြွ လန်းလန်း တစ်ခါတစ်လေ 2 ကာနှင့် ပတ်သက်၍ရ	ပျာ်ရွှင်စွာလုပ်ကိုင်နို ဖြစ်တောင့်ဖြစ်ခဲ	7 ວິເພີ່ວເພີ່ອງ ອາເພີ່ອງ ອາເພີອງ ອາເພື່ອງ ອາເພື່ອງ ອາເພື່ອງ ອາເພື່ອງ ອາເພື່ອງ ອາເພື່ອງ ອາເພື່ອງ ອາເພື່ອງ ອາເພື່ອງ	ာအခါမျှ ာအခါမျှ				
6. A. B.	1 ຈິດຈິຊິຣໍະລາຊິຣ໌ຄາຊູອ໌ະ ດູຣູຊ໌ອຸ້ອລາກ (၃) ດ ຈາກອາດກ 4 ດູຣູຊ໌ອຸ້ອລາກ (၃) ດ ຈາກອາດກ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် မကြာမကြာ ဆိုသလို 3 အတွင်း သင်တက်တ မကြာမကြာ ဆိုသလို 3 အတွင်းသင်၏အနာဂ မကြာမကြာ	4 အလုပ် များကို စိတ်ရ တစ်ခါတစ်လေ 2 က်ကြွကြွ လန်းလန်း တစ်ခါတစ်လေ 2 တစ်နှင့် ပတ်သက်၍တေစ်လေ	ပျာ်ရွင်စွာလုပ်ကိုင်နို ဖြစ်တောင့်ဖြစ်ခဲ	7 ເບິ່ງ ແມ່ນ ອາເມີຄວາ ອາເມີຄວາ	ာအခါမျှ ာအခါမျှ ။ ာအခါမျှ				
6. A. B.	1 ຈິດຈິຊິຣໍະລຸຊິຣ໌ຄາອູອ໌ະ ເບູຣ໌ອຸ້ອະသາ (၃) ເບ ອဏອဏ 4 ເບູຣ໌ອຸ້ອະນາ (၃) ເບ ອဏອဏ ຊ ເບູຣ໌ອຸ້ອະນາ (၃) ເບ ອဏອဏ	ဆောင်နိုင်ခြင်း အတွင်း နေ့စဉ်ပုံမှန် : မကြာမကြာ ဆိုသလို 3 အတွင်း သင်တက်တ မကြာမကြာ ဆိုသလို 3 အတွင်းသင်၏အနာက မကြာမကြာ ဆိုသလို	4 အလုပ် များကို စိတ်ရ တစ်ခါတစ်လေ 2 က်ကြွကြွ လန်းလန်း တစ်ခါတစ်လေ 2 တစ်နှင့် ပတ်သက်၍ရေ	မျာ်ရွှင်စွာလုပ်ကိုင်နိုး ဖြစ်တောင့်ဖြစ်ခဲ 1 ဆန်းဆန်းရှိပါသလာ ဖြစ်တောင့်ဖြစ်ခဲ 1 မျှာ်လင့်ချက်ပြည့်ဝစွ ဖြစ်တောင့်ဖြစ်ခဲ	7 ວ້ເປິນເလາະແ ອາເພີຣວາ:	ာအခါမျှ ာအခါမျှ ။။ ဘာအခါမျှ				

7.	လက်ရှိရောဂါများ			
	သင်တွင်အောက်ဖော်ပြပါရောဂါများရှိပါသလား။	ရိပါသည်။	ရှိပါသည်။	မရှိပါ။
	ဆရာဝန်ပြောကြား၍ကသမများပါဝင်ပါသည်။	မိမိအထင်။	ဆရာဝန်မ	л
	သင်လက်ခံထားမနှင်နီးစပ်သည်ဟသင်ယုဆသောအခြေ	(ဆရာဝန်	ပြောကြား	
	အနေတစ်ခုကို (🗸) ခြစ်ပါ။	နှင့်မပြပါ)	ထားပါသည်	
		2	1	0
•	(၁) မတော်တဆထိခိက် ဒဏ်ရာရခြင်း		_	
•	(၂) ခန္ဓာကိုယ်နေရာအနှံ(ခြေ၊လက်၊ကျောနှင်	2		
	အခြားနေရာများ) နာကျင်ကိုက်ခဲ့သည်ရောဂါ (၁ပမာ			
	အဆစ်အမြစ်၊ ကြွက်သားများ အဆက် မပြတ်ကိုက်ခဲ ယောင်			
	ယမ်းခြင်း၊ လေးဘက်နာ အဆစ်အမြစ် ရောဂါ)			
•	(၃) နှလုံးနှင့်သွေးကြောဆိုင်ရာရောဂါ (ဥပမာ၊ သွေးတိုး၊	C2		
	နှလုံး သွေးကြောကျဉ်း)			
•	(၄) အသက်ရှုလမ်းကြောင်းဆိုင်ရာရောဂါများ			
	(ဥပမာ၊အသက်ရှုလမ်းကြောင်းပိုးဝင်ခြင်း၊ အဆုတ်ပွခြင်း)			
•	(၅) စိတ်ပိုင်းဆိုင်ရာပြဿနၥများ (ဥပမာ၊ စိတ်ဓာတ် ကျခြင်း၊			
	ခြေကုန်လက်ပန်းကျခြင်း၊စိုးရိမ်စိတ်လွန်ကဲ ခြင်း၊အိပ်မပျော်ခြင်း)			
•	(၆) အာရုံကြောဆိုင်ရာရောဂါများ (ဥပမာ၊ အကြား အမြင်			
	ချို့တွဲခြင်း၊ခေါင်းတစ်ခြမ်းကိုက်ခြင်း၊ အတက်ရောဂါ))			
•	(၇) အစာလမ်းကြောင်းရောဂါများ (ဥပမာ အစာ			
	မကြေခြင်း၊သည်းခြေကျောက်တည်ခြင်း၊ အသည်း သို့မဟုတ်			
	ပန်ကရိယရောဂါများ၊ အဆက်မပြတ် ဝမ်းချုပ်ခြင်း)			-
•	(၈) မျိုးပွားနှင့်ဆီးလမ်းကြောင်းရောဂါများ (ဥပမာ၊			
	ဆီးလမ်းကြောင်းပိုးဝင်ခြင်း၊အမျိုးသမီးရောဂါ၊			
	ဆီးကျိတ်ရောဂါ)			
•	(၉) အရေပြားရောဂါများ (ဓာတ်မတည့်ခြင်း ၊ အနီစက်			
	များထွက်ခြင်း)			
•	(၁၀) အကျိတ် (သို့မဟုတ်)			
	ကင်ဆာရောဂါ			
•	(၁၁) အတွင်းထုတ်ဂလင်း သို့မဟုတ် ၆ဝကမ္မ ဆိုင်ရာ			
	ရောဂါများ (ဥပမာ၊ ဆီးချိုရောဂါ၊ အဝလွန်ရောဂါ သို့မဟုတ်			
	ဂေါက်ရောဂါ)			
•	(၁၂) သွေးရောဂါများ (ဥပမာ၊ သွေးအားနည်းခြင်း၊			
	အခြားသွေးရောဂါများ သို့မဟုတ် ချို့ယွင်းမှုများ)	0		
•	(၁၃) မွေးရာပါချို့ယွင်းမှုများ			
•	(၁၄) အခြားပုံမမှန်မှုများ သို့မဟုတ်			
	ရောဂါများ			

အပိုင်း ၅ လုပ်ငန်းခွင်ထုတ်လုပ်နိုင်စွမ်းနှင့်လုပ်ဆောင်မှုထိခိုက်ခြင်းမေးခွန်းလွှာ။

အဝိုင်း ၅ လုပ်ငန်းခွင်ထုတ်လုပ်နိုင်စွမ်းနှင့်လုပ်သောင်မှုထခုက်ခြင်းမေးခွန်းလွှာ။ အထွေထွေကျန်းမာရေး အောက်ပါမေးခွန်းများသည်သင်၏အလုပ်လုပ်နိုင်စွမ်းနှင့်ပုံမှန်လှုပ်ရှားမှုများအပေါ်သင်၏ကျန်းမာရေးပြဿ နာများ၏သက်ရောက်မှုအကြောင်းမေးမြန်းသွားမည်ဖြစ်သည်။ကျန်းမာရေးပြဿနာဆိုသည်မှာမည်သည့်ရုပ် ဝိုင်းဆိုင်ရာ၊စိတ်ခံစားမှုပိုင်းဆိုင်ရာပြဿနာ၊ဝေဒနာများကိုရည်ညွှန်းပါသည်။ လိုအပ်သလိုကွက်လပ်များ ကို ဖြည့်ပါ။နံပါတ်များကိုပိုင်းပါ။

စဉ်	အကြောင်းအရာ	ရွေးချယ်ခြင်း	မှတ်ချက်			
1.	လက်ရှိသင်အလုပ်ရှိပါသလား။	မဟုတ်ပါ။ 1	"မဟုတ်ပါ။" ဟု ဖြေ			
	(လုပ်ခငွေကြေးအတွက်အလုပ်လုပ်နေပါသ	ဟုတ်ပါသည်။ 2	ဆိုပါက စစ်ဆေးပြီး			
	လား။)		မေးခွန်း 6. သို့သွားပါ။			
	အောက်ပါမေးခွန်းများသည်လွန်ခဲ့သော (၇) ရက်	(ယနေ့မပါ) အတွင်းမှအဓ	ကြောင်းများ ဖြစ်သည်။			
2.	ပြီးခဲ့သော(၇) ရက်အတွင်း သင်၏ကျန်းမာရေး [ကြောင့်သင်လုပ်ငန်းခွင်မှ နာရီပေါင်း မည်မျှ ပျက် ခဲ့သလဲ။ သင်နေအကောင်းသဖြင့် ခွင့်ယူခဲ့သောရ အလုပ်ထဲနောက်ကျ ရောက် ရှိခြင်းများ၊ အလုပ်မှ ပြန်ခြင်းများ စသည့် နာရီ များအားလုံးပါဝင်သည့် (ဤသုတေသနတွင် ပါဝင်ရ၍ပျက်ကွက်ရသည့် : များမပါဝင်ပါ။)	ြဲဿနာ ဝ်ကွက် က်များ၊ – – – – – – စ စော နာရီပေါင်း ၅။ အချိန်	_1			
3.	ပြီးခဲ့သော (၇) ရက်အတွင်း ဥပမာ အားလပ်ရက် အလုပ်ပိတ်ရက်များ၊ ဤသုတေသနတွင်ပါဝင်ရန် စသည့် အခြားသောအကြောင်းများကြောင့် သင် ခွင်မှ နာရီပေါင်း မည်မျှပျက်ကွက်ခဲ့သလဲ။	များ။ ခြင်း ၂ လုဝ်ငန်း နာရီပေါင်း	1			
4.	ဗြီးခဲ့သော (၇) ရက်အတွင်းသင်အမှန် တကယ် န ပေါင်း မည်မျှအလုပ်လုဝ်ခဲ့ရသလဲ။	ာရီ နာရီပေါင်း	အကယ်၍ "ဝ" (သုည) ဖြေပါက – ၊ ဗေးခွန်း 6. သို့ သွားပါ။			
5.	ပြီးခဲ့သော (၇) ရက်အတွင်း သင်၏ကျန်းမာရေး[ထုတ်လုပ်နိုင်မှုအပေါ်မည်မျှထိခိုက်မှုရှိသလဲ။အဝ သောအလုပ်အမျိုးအစားနှင့်ပမာဏအကန့်အသပ သင်လုပ်နိုင်သည့်နေ့အရေအတွက်ထက် လျော့န ခါတိုင်းကဲ့သို့ သေသေချာချာအာရုံစူးစိုက်၍ မလု၊ ကသင်၏အလုပ်အပေါ်အနည်းငယ်သာထိခိုက်မှု ကျန်းမာရေးပြဿနာများက သင်၏အလုပ်အပေ ရွေးပါ။ <u>ကျန်းမာရေးပြဿနာများ</u> ကု န်ထုတ်လုပ်နိုင်စွမ်းရည်ကိုမည်	ပြဲဿနာများက သင်အလုပ် လုပ်လုပ်ရသည့် နေ့များအင တ်ဖြစ်ခြင်းကိုစဉ်းစားပါ။ ည်းစွာ လုပ်ကိုင်နေရမှု၊ အ ပ်နိုင်ခြင်းစသည်။အကယ်၍ ရှိပါက နံပါတ်အငယ်များကို ဖြာလွန်အမင်းထိခိုက်မှုရှိပ <u>၊းက သင်အလုပ်လုပ်နေစဉ်</u> မျှထိနိုက်နေသည်ကိုသာစ	ခွင်အတွင်း သင်၏ တွင်း သင်လုပ်နိုင် လုပ်ချိန်အတွင်း သင် ရဲ့ကျန်းမာရေး ပြဿနာ ရေးပါ။ အကယ်၍ ါက နံပါတ်အကြီး ကို ဦးစားပါ။			
	0 1 2 3 4 5	6 7 8	9 10			

		0 ကျန်းမာဖ များမိမိဒ အကျိုးသ	ရေးပြဿနာ စလုပ်အပေါ စက်ရောက်မှ မရှိပါ							ကျန်း၀ များမိ မလုပ် ဆီးထ	10 မာရေးပြဿနာ ဒိကိုလုံးဝအလုပ် နိုင်အောင်တား ားသည်။
6	(8.2			2 2	နံပါ သူတန်း	တ်တစ်ခု မာတပြ	ကိုသာက္	ဒွင်းပါ။	S		
	ပုံမှန် ထိန်း အစာ ပြဿ ရွေးပ ထိနိုး	လှုပ်ရှားမှ ခြင်း၊လေ့ အနှင့်ပမာ စနာကသ ဘမ္ဒရိပါဂ <u>ကျန်းမာ</u> ထိ ခိုက်မှု	ုဆိုသည်မှ ကျင့်ခန်းလ ဏအပေါ်း လင်၏လှုပ်ရ သိ၍ကျန်းလ က နံပါတ်အ သ ေရးပြဿ ချားကိုသာ	ာသင်စေ လုပ်ခြင် အကန့်း ဓားမှုမျာ မာရေးဖြ ဓကြီးကို နာများ စဉ်းစာ	နေ့စဉ်နေ းစာဖတ် အသတ်[မြိဿနာရ ပြိုသာနာရ ပြိုရွေးပါ။ မကြောင့် းပါ။ လုပ်	အိမ်တွင် ခြင်းစသ ဖြစ်မှုများ အနည်း၊ ချားက သ သင်၏နို ငန်းခွင်ဒ	နေထိုင်စ ည်။ သင် ငယ်သာ သင်၏လှုင် စွဲခူဝ ုံမှ န် ဓတွင်းအ	ဉ်လုပ်နေ လုပ်နိုင်ရ င်ခြင်းစၥ ထိနိုက်မှု လိုရားမှု မု န်အလုပ်ရ လုပ်လုပ်	နရသော သော ပုံမှ သည်။ အ ရှိပါက ခ များ အပေ များလုပ်န် သ်သည့်အ	ဈေးဝယ် န် လှုပ်ရှ ကယ်၍ နံပါတ် အ ြဲ အလွန် ရြိအရေး ခြေအရေး	ခြင်း၊ ကလေး ဘးမှု အမျိုး ကျန်းမာရေး မငယ်များကို အမင်း ရည်အပေါ် န မဟုတ်ဝါ။
	0	1	2	3	4	5	6	7	8	9	10
	0 ကျန်းမာရေးပြဿနာများ မိမိ၏နေ့စဉ်လှုပ်ရှားမှုများ အပေါ် အကျိုးသက်ရောက်မှု မရှိပါ။		:					ကျန် နေ့ စ	းမာရေးဖြ ဉ်လှုပ်ရှာ	10 ဒိဿနာများမိမိ၏	
		အကျိုးသ	အပေါ က်ရောက်မှ မရှိပါ	2					ကိုင် ပါသ	၍မရအေ ည်။	းမှုများလုံးဝလုပ် ၁င်တားဆီးထား

ပါဝင်ဖြေဆိုခြင်းအတွက်ကျေးဇူးတင်ပါသည်။

APPENDIX D: Participant information sheet and consent form (Myanmar)

သုတေသနတွင်ပါဝင်သူများကိုအသိပေးစာနှင့်သဘောတူညီချက်ပုံစံ

အပိုင်း ၁ သုတေသနနှင့်ပတ်သက်သောအသိပေးစာ

<u>၁။ နိဒါန်း</u>

ကျွန်တော်သည် ထိုင်းနိုင်ငံ၊ ဘန်ကောက်မြို့၊ချူလာလောင်ကွန်းတက္ကသိုလ်တွင် ပြည်သူ့ကျန်းမာရေး မဟာဘွဲ့ကို ဆည်းပူးနေသောကျောင်းသားတစ်ယောက်ဖြစ်ပါသည်။ တက္ကသိုလ်၏ပညာရေးလိုအပ်ချက်များအရကျွန်တော် သည်သုတေသနတစ်ခုလုပ်ပြီးစာတမ်းကျမ်းတစ်စောင်တစ်ခုတင်ပြရန်လိုအပ်သည်။ကျွန်တော်၏ မဟာဘွဲ့ အတွက်ပြုစုနေသောစာတမ်းကျမ်းမှာ မြန်မာနိုင်ငံရှိအထည်ချုပ်စက်ရုံမှ စက်ချုပ် လုပ်သားများ ၏ စိတ်ပိုင်းလူမှုရေးပိုင်းဆိုင်ရာသွင်ပြင်လက္ခဏာများ၊လုပ်ငန်းခွင်ထုတ်လုပ်နိုင်စွမ်း၊လုပ်ငန်းခွင် အလုပ်လုပ်နိုင်စွမ်း နှင့်ကြွက်သားနှင့်ကျောရိုးဆိုင်ရာမအိမသာဖြစ်ခြင်းတို့၏ဆက်နွယ်မှု အကြောင်း ဖြစ်ပါသည်။

သုတေနပြုလုပ်သူအမည် – မစ္စတာထွန်းဝင်း ဦး

ဖုန်းနံပါတ်: ၀၉–၄၃၀–၂၀–၅၂၉

အီးမေးလ်လိပ်စာ: tunwinoo430@gmail.com

လေ့လာမှု၏ခေါင်းစဉ် –

မြန်မာနိုင်ငံရှိအထည်ချုပ်စက်ရုံမှ စက်ချုပ်လုပ်သားများ၏စိတ်ပိုင်းလူမှုရေးပိုင်းဆိုင်ရာ သွင်ပြင် လက္ခဏာများ၊ လုပ်ငန်းခွင်ထုတ်လုပ်နိုင်စွမ်း၊လုပ်ငန်းခွင်အလုပ်လုပ်နိုင်စွမ်းနှင့်ကြွက်သားနှင့်ကျောရိုးဆိုင်ရာမအီမသာဖြစ်ခြင်း တို့၏ဆက်နွယ်မှု

ဤသုတေသနတွင်ပါဝင်ရန်သင့်ကိုလေးစားစွာဖိတ်ခေါ်ပါသည်။ ဤသုတေသနတွင် မြန်မာနိုင်ငံရှိ စက်ချုပ် လုပ်သားပေါင်း (၄၀၀) ခန့်ပါဝင်မည်ဖြစ်ပါသည်။ အောက်တွင် ဖော်ပြမည့် အချက်အလက်များကို ဆက်လက် ဖတ်ရှု၍ မရှင်းလင်း သည် ကိုမေးမြန်းနိုင်ပါသည်။ဤသုတေသနတွင် သင်ပါဝင်မည်/မပါဝင်မည်ကို သင် လွတ်လပ်စွာ ဆုံးဖြတ်နိုင်ပါသည်။ သင်မည်သည့်အရာကိုထပ်မံရှင်းလင်းလိုလျှင်၊ လေ့လာမှုမပြုလုပ်မီနှင့်၊ အပြီး၌သုတေသနပြုလုပ်သူအား မေးမြန်းရန်မတွန့်ဆုတ်ပါနှင့်။

၂။ မေးခွန်းလွှာတွင်ပါဝင်သောအကြောင်းအရာများ

ဤမေးခွန်းလွှာတွင်မိမိဖာသာဖြေဆိုရမည့်အဝိုင်း (၅)ဝိုင်းပါဝင်၍ကြာမြင့်ချိန် မိနစ် (၃၀)ခန့်ဖြစ်ပါသည်။ အဝိုင်း (၅)ဝိုင်းမှာ

(၁) ဖြေဆိုသူ၏နောက်ခံအချက်အလက်များ

(၂) ကြွက်သားများနှင့်ကျောရိုးဆိုင်ရာမအီမသာဖြစ်မှုမေးခွန်းလွှာများ

(၃) စိတ်ဝိုင်း၊လူမှုရေးဝိုင်းဆိုင်ရာသွင်ပြင်လက္ခဏာမေးခွန်းလွှာများ

(၄) ထုတ်လုပ်နိုင်စွမ်းနှင့်ပတ်သက်သောမေးခွန်းလွှာများ

(၅) အလုပ်လုပ်နိုင်စွမ်းနှင့်ပတ်သက်သောမေးခွန်းလွှာများ ပါဝင်ပါမည်။

155

မေးခွန်းလွှာတစ်ခုလုံးသည် ဖြည့်စွက်ရခြင်း၊ ရွေးချယ်စရာများအပေါ်အမှတ်ခြစ်ပေးရခြင်း၊ ဖြေဆိုသူ အဆင်ပြေ စေရန် ပေးထားသည့် အဖြေများပေါ်တွင်ဝိုင်းပေးရခြင်းများပါဝင်ပါမည်။ သင်၏ဖြေဆိုမှုများကို မည်သူဖြေကြား မှန်း မသိအောင် လျှို့ဝှက်ပေးထားမည်ဖြစ်ပါသည်။

<u>၃။ သုတေသန၏ရည်ရွယ်ချက်</u>

(၁) စက်ချုပ်လုပ်သားများ၏နောက်ခံအချက်အလက်များ၊ ထုတ်လုပ်နိုင်စွမ်း၊ အလုပ်လုပ်နိုင်စွမ်း၊ စိတ်ပိုင်းဆိုင်ရာ လူမှုရေးဆိုင်ရာအချက်များနှင့် ကြွက်သားများနှင့် ကျောရိုးဆိုင်ရာမအီမသာဖြစ်မှုများကိုရှာဖွေ ဖော် ထုတ်ရန်။

(၂) ပုဂ္ဂိုလ်ရေးအချက်များ၊ ထုတ်လုပ်နိုင်စွမ်း၊ အလုပ်လုပ်နိုင်စွမ်း၊ စိတ်ပိုင်းဆိုင်ရာ လူမှုရေးဆိုင်ရာ အချက်များနှင့် ကြွက်သားများနှင့်ကျောရိုးဆိုင်ရာမအီမသာဖြစ်မှုများ အကြား ဆက်စပ်မှုကိုသိရှိရန်။

<u>၄။ လုပ်ထုံးလုပ်နည်း</u>

ဤအသိပေးစာကိုဖတ်ရှုဒြီး၊ လိုအပ်ပါက မေးခွန်းလွှာမှမရှင်းလင်းသည့်အချက်များကို သုတေသန ပြုလုဝ်သူ မှထဝ်မံရှင်းလင်း ပြောကြားပေးပြီးသည့်နောက် ပါဝင်ခြင်းနှင့် ဖယ်ထုတ်ခြင်း ဆိုင်ရာ စံသတ်မှတ်ချက်များအရ ပါဝင်မည့်သူများသည်မိမိတို့ပါဝင် ရွေးချယ်ခံရမှု အတွက် သဘောတူညီ ကြောင်းကိုစာဖြင့် အသိပေး ရန်လိုအပ်လိမ့်မည်။ သဘောတူညီကြောင်းစာကို သုတေသနပြုလုဝ်သူက လုံခြုံစွာသိမ်းဆည်း ထား မည်ဖြစ် သည်။

သင်သည်ဤလေ့လာမှုတွင်မပါဝင်လိုပါက သဘောတူညီချက်ပေးရန်မလိုအပ်ပါ။ အကြောင်းပြချက် တစ်ခုခု ပေးရန်လည်းမလိုအပ်ပါ။ အကြောင်းပြချက်ပေးရန်မလိုဘဲသင်ဆန္ဒရှိသည့် အတိုင်းအချိန်မရွေး သင်နှုတ်ထွက် နိုင်ပြီးယင်းသည်ပါဝင်သူများအပေါ်မည်သည့်ဆိုးကျိုးတစ်စုံတစ်ရာသက်ရောက်မှုမျှရှိလိမ့်မည်မဟုတ်ပေ။

၅။ အကျိုးကျေးဖူးများ

ဤလေ့လာမှုသည်စက်ချုပ်လုပ်သားများအကြောင်းလေ့လာမှုဖြစ်၍သင့်အတွက်အကျိုးကျေးစူးရှိပါလိမ့်မည်။ အဆိုပါလေ့လာမှုသည်စက်ချုပ်လုပ်သားများအကြားစိတ်ဝိုင်းလူမှုရေးဝိုင်းဆိုင်ရာပြဿနာများ၊ထုတ်လုပ်နိုင်စွမ်း၊ အလုပ်လုပ်နိုင်စွမ်းနှင့် အလုပ်နှင့်ဆက်စပ်သောကြွက်သားများနှင့် ကျောရိုးဆိုင်ရာ မအီမသာဖြစ်မှုများ ကိုရှာဖွေရန်ရည်ရွယ်သည်။ ထို့ကြောင့်လေ့လာမှုရလဒ်သည်စက်ချုပ်လုပ်သားတို့အတွက် ဝိုခိုကောင်းမွန်သော လုပ်ငန်းရွင်အခြေအနေ၊ ကျန်းမာရေးအတွက်လုပ်ငန်းရွင်လာမှုအလက်များကိုပေးစွမ်းနိုင်မည်ဖြစ်သည်။

<u>၆။ အချက်အလက်များကိုလျှို့ဝှက်ထိန်းသိမ်းထားမှု</u>

သင်နှင့်ဆက်စပ်သောနေသောမည်သည့်သတင်းအချက်အလက်ကိုမဆိုလျှို့ဝှက်ထားပါမည်။ သင်၏နံမည်များ (သို့မဟုတ်) တစ်စုံတစ်ယောက်ဟုသိရှိနိုင်သော အခြားအချက်အလက်များကိုအစီရင်ခံစာ (သို့မဟုတ်) လေ့လာမှု၏ အနှစ်ချုပ်တွင်ဖော်ပြမည်မဟုတ်ပါ။ နောက်ဆုံးအစီရင်ခံစာကိုသုတေသနပြုလုပ်သူထံမှရရှိနိုင်ငြီး ဤအစီရင်ခံစာကိုပြည်သူ့ကျန်းမာရေးမဟာဘွဲ့၏ပညာရေးလိုအပ်ချက်ကိုဖြည့်ဆည်းရန်သာသုံးလိမ့်မည်။ အချက်အလက်အားလုံးကိုလျှို့ဝှက်ထိန်းသိမ်းထားရှိမည်ဖြစ်ပြီးမည်သူတစ်ဦးတစ်ယောက်ကိုမျှထုတ်ဖော်ပြော ကြားသွားမည်မဟုတ်ပါ။

<u>ဂု။ ပါဝင်ခွင့်နှင့်သဘောတူညီချက်</u>

ကျွန်ုပ်သည်ဤသုတေသန၏ရည်ရွယ်ချက်များ၊ ဤသုတေသန၏ အကြောင်းအရာများ၊ အကျိုးကျေးဇူးများ နှင့်အန္တရာယ်အလားအလာများ/ထိခိုက်နစ်နာမှုများ (တစ်စုံတစ်ရာရှိခဲ့ပါက) ပါဝင်သူများ၏ အခွင့်အရေးနှင့် တာဝန်များကို ဖတ်ရှုပြီးဖြစ်ပါသည်။ ကျွန်ုပ်သည်သုတေသနပြုလုပ်သူနှင့် ဆက်သွယ်နိုင်မည့်အချက်အလက် အသေးစိတ်ကိုလည်း ရရှိထားသည်။ ကျွန်ုပ်သည်အသိပေးစာကို ဖတ်ရှုအပြီး သုတေသနပြုလုပ်သူက ကျွန်ုပ်ကိုရှင်းပြပြီးနောက် အသိပေးစာတွင်ဖော်ပြထားသည့်အတိုင်းဆောင်ရွက်ရန် ကတိပြုပါသည်။ ကျွန်ုပ်သည် ကောင်းမွန်ကျေနပ်စွာနားလည်သဘောပေါက်ပါသည်။ ဤသုတေသနတွင်ပါဝင်ရန် ကျွန်ုပ်ဆန္ဒရှိကြောင်း သဘောတူပါက ဤလေ့လာမှုတွင် ကျွန်ုပ်၏အမည်နှင့်အခြားတစ်စုံတစ်ယောက်ဟု သိရှိနိုင်သော မည်သည့် အချက်အလက်များမှပါဝင်မည်မဟုတ်ပါ။ ဤသည်မှာ ဓိနှစ် (၃၀) ခန့်ကြာမြင့်မည်ဖြစ်သည်။

လေ့လာမှုမစဓီ၊ သင်ပါဝင်လုပ်ဆောင်နေစဉ်နှင့်ပါဝင်မှု အပြီးတွင် သင်သိလိုသည့် မည်သည့် အကြောင်း အရာကိုမဆိုမေးမြန်းနိုင်သည်။ အဓိကသုတေသနပြုလုပ်သူ သည်အထက်ဖော်ပြပါ လိပ်စာနှင့် အတူ အချိန်မရွေး ဆက်သွယ်နိုင်ပါသည်။ သုတေသနပြုလုပ်သူသည်အကျိုးကျေးဇူး (သို့မဟုတ်) အန္တရာယ်/ ထိခိုက်နှစ်နာမှုနှင့် ပတ်သက်သောသတင်းအချက်အလက်အသစ်များရှိပါက သင်ကိုအမြန်ဆုံး အကြောင်းကြားပါလိမ့်မည်။

ပါဝင်သူများအတွက်သတိပေးစာ နှင့် သဘောတူညီချက်ပုံစံတွင်ဖော်ပြထားသည့် အတိုင်းသုတေသနပြုလုဝ်သူက မလုဝ်ဆောင်ပါက၊ သင်သည်အဖြစ်အပျက်ကို ထိုင်းနိုင်ငံ၊ ဘန်ကောက်မြို့၊ချူလာလောင်ကွန်းတက္ကသိုလ်ရှိ **သုတေသနကျင့်ဝတ်သုံးသပ်ရေးကော်မတီ** Research Ethics Review Committee for Research Involving Human Research Participants (RECCU), Group I, Chulalongkorn University (RECCU) Jamjuree 1 Bldg., 254 Phyathai Rd., Patumwan district, Bangkok 10330, Thailand, တယ်လီဖုန်းအမှတ်/ဖက်(စ်) ဝ–၂၂၁၈–၃၂ဝ၂၊ ဝ–၂၂၁၈–၃ဝ၄၉ အီးမေးလ်၊ eccu@chula.ac.th" သို့အကြောင်းကြားတိုင်ကြားနိုင်ပါသည်။

ဤသဘောတူညီချက်ပုံစံမှအချက်အလက်တွေကိုကျွန်ုပ်ဖတ်ရှုပြီးပါပြီ။ ထို့ပြင်၊ ကျွန်ုပ်သည်ပါဝင်သူ၏ အသိပေးစာ နှင့် သဘောတူညီချက်ပုံစံဓိတ္တူကိုလက်ခံရရှိခဲ့သည်။

3

ကျွန်ုပ်သည်ဤလေ့လာမှုတွင်ပါဝင်ရန်သဘောတူသည်။

_)

သုတေသနတွင်ပါဝင်သူ၏လက်မှတ်

ထွန်းဝင်း ဦး (အဓိကသုတေသနပြုလုဝ်သူ) တွဲဖက်ပါမောက္ခ Nutta Taneepanichskul (စာတမ်းအကြံပေး)

VITA

NAME Tun Win Oo

DATE OF BIRTH 9.7.1991

PLACE OF BIRTH Yangon, Myanmar

INSTITUTIONS ATTENDED College of Public Health Science, Chulalongkorn University

HOME ADDRESS

No. (29), Aungzayar 1st street Yangonthit Quarter,

Thingangyun Township, Yangon, Myanmar



จุฬาลงกรณ์มหาวิทยาลัย Chulalongkorn University